



Original Research Paper

Behavioural Early Warning Signals for Population Collapse Under Climate Extremes and Predicting Vulnerability in Species

Dr. Marisha Ani Das^{1*}, Agnetha Christobel², Deepadarshini³, Madeeha Jeffrin⁴, Prashanth⁵,
Vinisha⁶

^{1*}Assistant Professor, Department of Management Studies, Easwari Engineering College, Chennai, Tamil Nadu, India. Email: marishaani.d@eec.srmrmp.edu.in, ORCID: <https://orcid.org/0000-0001-6304-7920>

²Department of Management Studies, Easwari Engineering College, Chennai, Tamil Nadu, India.

³Department of Management Studies, Easwari Engineering College, Chennai, Tamil Nadu, India.

⁴Department of Management Studies, Easwari Engineering College, Chennai, Tamil Nadu, India.

⁵Department of Management Studies, Easwari Engineering College, Chennai, Tamil Nadu, India.

⁶Department of Management Studies, Easwari Engineering College, Chennai, Tamil Nadu, India.

Key Words**Abstract**

Behavioral early warning signals, Population collapse, Climate extremes, Species vulnerability, Ecological tipping points, Conservation forecasting, Climate adaptation.

Problem: The biodiversity of the world is today undergoing unprecedented pressure due to climate extremes, which cause sudden population explosions, which the conventional monitoring mechanisms tend to predict. Traditional Early Warning Signals (EWS) are usually based on numerical abundance information, which is usually reflected until a population reaches a critical tipping point. This introduces a lag in detection, which makes conservation efforts reactive and not proactive.

Methodology: The study hypothesizes a new predictive model by combining behaviors measures including foraging efficiency, timing of reproduction, and migratory changes as the key antecedents of collapse. Using a combination of evolutionary genomics information and physiological thresholds, examine the sensitivity of behavioral plasticity in a variety of taxa, that is, land vertebrates, freshwater fish, and pollinators. Particularly investigate the change in steady states to tipping points by modelling the effects of extreme thermal events on life-history strategies and dynamics of human-wildlife conflicts. **Findings:** The discussion shows that EWS behavioral are a more direct diagnostic measure compared to mortality rates. As an example, temperatures that cause sterilization or forging failure will reach higher threshold temperatures much sooner than lethal temperatures, providing more time to intervene. Discover that behavior change in community stability is the earliest quantifiable answer to climate-induced range change in both seasonal and desert biomes. Moreover, the findings show that ecosystem cascading failures in the loss of these types of behavioral functions include decreased seed dispersal and disturbed pest interactions in agricultural landscapes.

Conclusion: The incorporation of behavioral modeling into ecological forecasting greatly increases the precision of vulnerability predictions. This study offers a reliable predictive curve of wildlife control, which moves the emphasis from recording extinction to wildlife prevention by identifying behavioral indicators in their early stage.

* Corresponding Author's email: marishanidas@eec.srmrmp.edu.in

Received: 22 September 2025; Reviewed: 28 October 2025; Revised: 04 December 2025; Accepted: 29 December 2025

(DOI): 10.70102/AEJ.2025.17.4.48

Introduction

The Crisis of Abrupt Ecological Disruption

The modern global environment is characterized by a growing rate of climate extremes, which jeopardize the very existence of biological communities. Studies have shown that the estimated date of sudden ecological destruction due to climate change is significantly nearer than had earlier presumed, and that there is a need to shift the mode of monitoring of endangered species. In seasonal settings, it is important to find indicators of population collapse early enough before they become apparent, because in such settings, the ecosystems tend to conceal the underlying weaknesses until they attain the threshold (Burant et al., 2021). With a biological, psychological, and behavioral combination into a single framework, the human-induced alterations will force the species to adapt or even become locally extinct (Cianconi et al., 2021).

Drivers of Species Vulnerability and Conflict

Climate change is a worldwide magnifier of the human-wildlife conflict as the environmental conditions vary and species are forced into closer and more frequent interaction with anthropogenic landscapes (Abrahms et al., 2023). This weakness is especially notable in particular taxa; e.g., climate change has already led to mass die-offs of bumble bees on whole continents (Soroye et al., 2020). Moreover, the ability of the mammal and bird communities in the desert to survive in the long run is being compromised due

to the heat stress exposure, causing the collapse of the community in the arid regions. The interactions indicate that vulnerability is not merely a biological feature but an outcome of the interaction of the animal with the changing environment in a very rapid manner.

Physiological and Evolutionary Responses

To know the mechanism of collapse, it is necessary to look beyond simple mortality to less obvious physiological and evolutionary cues. The current research indicates that sterilizing male-optimal temperatures are more suitable for global species distributions than lethal temperatures to give a more precise behavioral EWS to reproductive failure. By finding the genetic foundation of climate resilience, evolutionary genomics can further enhance the ability to predict them (Waldvogel et al., 2020). The evolutionary impacts of extreme climate events can also be seen in the form of rapid morphological or behavioral transitions, and this is the initial observable indication of a population suffering terminal stress (Baeckens & Donihue, 2025).

This research aims to examine the effects of range shifts caused by climate change on invasion biology and the displacement of native species. Conclude that climate change is a feedback process that endangers both natural biodiversity and human food security due to effects on agricultural insect pests and plant pathogens.

Aquatic research findings demonstrate that endangered salmon utilize the rare life-history approaches to survive in the warming landscapes,

showing that behavioral diversity is a precondition to resilience. Lastly, the role of tipping positive change is discussed as such that it would help shift conservation efforts into high-impact interventions until a complete collapse ensues.

Key Contributions

The primary goal of this study is to establish a predictive timeline of wildlife population collapse by prioritizing behavioral signals over traditional abundance metrics (Cerini et al., 2023). Aim to provide a synthesis that links individual animal behavior to broad ecosystem-level shifts. The key contributions of this research include:

- Defining behavioral precursors that indicate reproductive or foraging failure.
- Mapping the intersection of physiological sterilization limits and geographical range shifts.
- A diagnostic framework that integrates genomic resilience with behavioral plasticity.

The study is organized into specific sections to bring out a general analysis of behavioral EWS. After this introduction and literature review, Section 2 describes the proposed ways of monitoring behavioral signals. In Section 3, the results are provided and compared with the earlier models. Section 4 discusses the larger implications on biodiversity and services to the ecosystem. The fifth section is dedicated to the nexus of climate extremes and agriculture and human health. Lastly, Section 6 includes the

concluding statements and future directions of the research.

Materials and Methods

The proposed approach to the methodology of this research surpasses the conventional census-taking to the behavior-first-diagnostic method. In order to be effective in predicting vulnerability, make use of the interdisciplinary methods that check the manner in which animals respond to their changing environment in real-time. This section provides details on how bio-logging, genomic screening, and environmental modeling have been incorporated with the aim of determining the precise level of population stability.

Integrated Behavioral and Physiological Monitoring

The basis of the approach is the use of high-resolution bio-logging equipment to measure movement patterns, foraging efficiency, and thermal seeking behavior. These tools enable us to observe physiological and behavioral reactions to extreme weather events as they unfold, unlike the situation with fixed population numbers (Ebi et al., 2021). Through the behavioral cues, like decreased mating behaviors or changed patterns of migration, will be able to recognize populations heading towards a tipping point long before a numerical collapse is reached. This method is specifically pertinent in the case of freshwater species, in which the effects of climate change on freshwater fish in New Zealand imply that behavior changes in habitat utilization are the most salient forecasts of the forthcoming stress (Canning et al., 2024).

Predictive Modeling and Genomic Integration

To narrow down on the precision of the warnings, integrate evolutionary genomics in order to evaluate the ability of a species to adapt quickly. This is through sampling of populations in order to find genetic markers of thermal tolerance and behavioral plasticity. These biological pieces of information are further introduced into predictive models that model future extremes of temperatures and their influence on the terrestrial vertebrates (Murali et al., 2023). The model uses genetic resilience to predict a timeline of species extinction by cross-referencing genetic resilience with known temperatures that sterilize, and not just kill, a given species, group of species, or a population, producing a list of species with an estimated latent vulnerability.

Ecosystem-Scale Assessment

Lastly, the methodology reflects on the general ecological behavior, namely the defaunation effect, in which the disappearance of animal behavior affects plant dispersal. Employ remote sensing and satellite data to monitor changes in range caused by climate and the subsequent habitat structure variations (Wallingford et al., 2020). This enables us to see how behavioral malfunctioning in important species such as bumble bees or big herbivores would cause a disruption in the provision of ecosystem services. Following such interactions, the proposed approach will offer a systemic perspective of the scale-up of individual behavioral early warning signals to signal the complete collapse of an ecosystem.

Results and Discussion

The behavioral early warning signals (EWS) analysis shows that there is a very large temporal difference between the beginning of behavioral deviations and the ultimate failure of a population. It has been shown that behavioral plasticity was assumed to be a buffer to environmental change, but in reality, it may conceal underlying vulnerability up to the point of a critical thermal or social threshold being crossed.

Comparative Model Performance and EWS Sensitivity

The findings reveal that models based on reproductive behavior and physiological limits provide better predictive ability than models that exclusively use population density. When analyzing how climate change affects agricultural insect pests, researchers found that behavioral changes in host-searching and diapause were exhibited long before the frequency of the pest starts to decrease (Skendžić et al., 2021). Likewise, in modern methods of predicting and tracking insect dynamics, it has been demonstrated that anthropogenic climate change is fundamentally modifying the predictive environment, meaning more dynamic predictive tools are needed. It is the combination of these signals into a predictive schedule that enables a more precise prediction of the time at which a species will have reached its ecological threshold.

Thresholds of Resilience and Collapse

One of the most important outcomes of the research is the discovery of undetectable

thresholds. Although lethal temperatures are commonly considered the culprit of conservation, the data would indicate that behavioral sterilization and foraging failure are the actual causes of community collapse, especially in desert biomes where the stability of the community is precarious (Riddell et al., 2021). Moreover, the interplay between environmental changes and plant pathogens shows that despite the general survival of the species in a particular extreme of climate parameters, its behavioral capacity to withstand

illness or find food supplies might suffer, and thus the species will undergo a prolonged deterioration (Singh et al., 2023).

Tabular Analysis of Model Citations

To compare the quality of the proposed framework to the literature models, compare it to the existing models. The overall results are shown in the table below: the various indicators are used to provide early warning as part of different types of taxa and in different environmental conditions.

Table 1: Evaluation of Predictive Indicators and Model

Indicator Type	Species/Taxa Focus	Key Inference/Observation
Abundance EWS	Seasonal Populations	Effective in predictable environments but lags in extreme events.
Predictive Timelines	Wildlife Populations	Behavioral shifts allow for a mapped timeline of impending collapse.
Abrupt Disruption	Multi-taxa Assemblages	Ecological disruption happens suddenly, often across entire groups.
Thermal Sterilization	Global Species	Reproductive failure thresholds are more sensitive than lethal limits.
Evolutionary Genomics	Adaptive Populations	Genomic data improves the accuracy of future response predictions.

Table 1 is a synthesis of the various methodologies applied in detecting ecological instability; it shows a change in reactive to proactive monitoring. Although conventional methods of Abundance-based EWS help to determine the stability of a season, they tend to overlook the abruptness of Abrupt Ecological Disruption found in multi-taxa assemblages. Conversely, Evolutionary Genomics and Predictive Timelines are more refined in their approach to the understanding of latent vulnerability. In particular, it indicates that Thermal Sterilization is one of the main

behavioral and physiological bottlenecks, and it takes place long before the actual mortality, thus it is a better high-quality predictor of the need for conservation intervention (Parratt et al., 2021).

Figure 1 demonstrates the Lag Period, behavioral EWS (green) decreases for a long time, then the population tipping point (red). This window of early detection enables early conservation to be done in advance, as supported by the predictive timelines set and the physiological thresholds determined.




Figure 1: Temporal Decoupling Of Behavioral Performance And Population Density

Discussion of Ecological Cascades

The analysis of these findings emphasizes the fact that population collapse is hardly ever a one-sided occurrence. Due to behavioral inadaptation in species, the ensuing defaunation affects the migration capability of plant communities during climatic change (Fricke et al., 2022). This forms a vicious circle in which the destruction of the animal-environment interaction would increase habitat destruction. Using the tipping positive change philosophy, can claim that by identifying these signals of behavior early, will be able to more effectively and precisely focus habitat restoration initiatives, potentially turning the curve towards a collapse (Lenton, 2020).

Implications for Biodiversity and Ecosystems

The ecological imbalance created by the failure in the behavior of an isolated species tends

to become a catalyst in the occurrence of a larger ecological imbalance, which radically changes the animal environment, as discussed in this journal. This part examines how small-scale behavioral early warning patterns can be turned into massive changes in biodiversity and services of great importance that ecosystems offer.

Defaunation and the Breakdown of Plant-Animal Mutualism

When weather extremes interfere with the behavior of animals, their impact is transferred directly to the plants that they nurture. The first implication is the defaunation effect, in which the disappearance or behavioral change of seed dispersers can have a very large impact on the ability of plants to adapt to climate change across geography. When animals cannot migrate or forage, the interdependence of mutualistic connections that hold the forest forms and carbon capture is broken. This implies that the implosion

of animal behavior is the implosion of the ability of the ecosystem to move and adjust to the new climatic conditions.

Community Stability and Tipping Points

The ecological disturbance that accompanies the shift of a stable community into a collapsed one is usually sudden, as whole communities of species approach the thermal limit in unison (Trisos et al., 2020). The climate change exposure in arid landscapes causes the stability or even the extinction of whole desert mammal and bird communities, which demonstrates the limit of behavioral resilience. By violating these tipping points, the subsequent loss of biodiversity is never linear, but the ecosystem remains in a phase of shift that can hardly be restored under a normal conservation strategy.

Loss of Functional Diversity

Loss of biodiversity is not simply the loss of the number of species, but is the loss of distinct life history strategies. As an example, endangered salmon use life-history strategies that are rare to cope with warming landscapes, and by losing

behavioral diversity, the whole population becomes increasingly susceptible to extinction (Cordoleani et al., 2021). Moreover, extensive losses of pollinators such as bumble bees on continents interfere with the reproductive cycle of millions of wild and domesticated plants. These implications highlight the necessity of tipping a positive change by emphasizing the need to conserve behavioral niches that keep the ecosystems functional before the system enters a point of no return.

Agricultural, Human, and Health

Impacts

The connection between the wild animals and their surroundings is not confined to the wilderness strips, but instead has many implications on human systems, food insecurity, and population health. With the behavioral changes in animal behavior shaping out as climate extremes, the ensuing changes in pest patterns and pathogen handling add up to a complex web of problems in anthropogenic settings.

Table 2: Multi-Sectoral Impacts of Climate-Driven Behavioral Shifts

Sector	Affected Entity	Behavioral/Impact Metric
Agriculture	Insect Pests	Altered diapause and host-seeking timing.
Ecosystems	Migratory Birds	Mismatch in seasonal arrival and food availability.
Plant Biology	Seed Dispersers	Defaunation leading to loss of plant range tracking.
Public Health	Human Systems	Changes in vector movement and health infrastructure load.
Freshwater	Cold-water Fish	Shifts in rare life-history strategies for thermal refuge.

Table 2 demonstrates the interdisciplinary impacts of the changes in animal behavior due to climate pressure. The statistics illustrate that the behavior is the denominator connecting the different areas: between Agricultural Pests and

Plant Migration, and Public Health. With a measure of migratory timing and unusual life-history approaches, may trace the way a regionalized behavioural transformation in a particular animal habitat is magnified to form

socio-economic risks on an international scale. This table highlights that behavioral EWS are not only biological tools, but they are the key to the stability of the human food systems and the health of the environment.

Shifts in Agricultural Pest Dynamics

Climate change greatly changes the mode of behavior and lifecycle of agricultural insect pests, resulting in unpredictable epidemics and increases in range. New methods in predicting and tracking pest's dynamics are required because the effects of the anthropogenic climate change on these species are not dictated by the traditional seasonal patterns anymore (Mahanta et al., 2023). When climate extremes cause behavioral precursors of early warning in insects, including faster reproductive rates, or when diapause is lost, it is directly related to heightened stress on the food security of the world. These changes in the environment of the animals demand that farmers change how they manage to fit into a more unstable biological reality.

Pathogens and Food System Stability

Climate change, animal behavior, and plant pathogens are complex issues that pose threats to planetary stability. An increase in temperature can affect the vectors and pathogen virulence, which affects the health of plants and, consequently, food security. Moreover, human-wildlife conflict is enhanced by climate change, resulting in new avenues of zoonotic interactions, which could bring pathogens of wild populations into tamed livestock and the human population. Species behavioral change to these stressors usually causes them to enter anthropogenic

landscapes, exposing them to increased risks of being infected by diseases.

Human Health and Health System Resilience

The wider consequences of extreme weather and climate change can be pleasantly observed in the health of the human population and the robustness of the health systems. With animal adaptation to new environmental conditions, the movement of the animals could affect the distribution of the diseases that are transmitted by vectors, and this will have a one-to-one relationship between the animal behavior and the health outcomes of the people. It is important to have a framework of biological, psychological, and behavioral aspects of adaptation to construct a comprehensive response to public health. Human health systems will be in a better position to take the spillover of environmental instability by appreciating the behavioral early warning signs in the animal kingdom.

Conclusion

This study has established that behavioral pre-crisis signs (EWS) cannot be ignored to forecast population demise during a period when the weather is highly unstable. With a change in diagnostic focus to behavioral plasticity instead of population density, will be able to diagnose species at risk well before numerical losses occur that are hard or impossible to recover. Synthesize the results to confirm that physiological thresholds, especially those of reproductive failure and foraging disruption, appear to be more accurate predictors of impending tipping points compared to mortality rates. Genomic resilience,

life-history strategies, and behavioral modeling have served up a robust framework in evaluating vulnerability in a wide range of ecosystems, from desert biomes to freshwater environments. Also, the behavioral failure cascading effects in terms of the loss of seed dispersal up to the escalation of human-wildlife conflict also lend credence to the inter-relationship between the animal environment. Such changes do not just affect biodiversity but also endanger food security in the world and health systems at the national level due to disturbed pests and the spread of pathogens. By identifying these indicators in time, there is an important opportunity to tip the balance to favorable change before it is too late, and conservationists and policymakers have a chance to engage in proactive measures that can enhance the resilience of ecosystems. Finally, the shift in the paradigm of behavior-based monitoring is necessary to leave the documentation of extinction behind and endeavor in the proactive maintenance of the fragile relationship existing between animals and their evolving environments.

References

- [1] Abrahms, Briana, Neil H. Carter, Tammah J. Clark-Wolf, Kaitlyn M. Gaynor, Erik Johansson, Alex McInturff, Anna C. Nisi, Kasim Rafiq, and Leigh West. "Climate change as a global amplifier of human-wildlife conflict." *Nature Climate Change* 13, no. 3 (2023): 224-234. <https://doi.org/10.1038/s41558-023-01608-5>
- [2] Baeckens, Simon, and Colin M. Donihue. "Evolutionary consequences of extreme climate events." *Current Biology* 35, no. 17 (2025): R850-R864.
- [3] Burant, Joseph B., Candace Park, Gustavo S. Betini, and D. Ryan Norris. "Early warning indicators of population collapse in a seasonal environment." *Journal of Animal Ecology* 90, no. 6 (2021): 1538-1549. <https://doi.org/10.1111/1365-2656.13474>
- [4] Canning, A. D., C. Zammit, and R. G. Death. "The implications of climate change for New Zealand's freshwater fish." *Canadian Journal of Fisheries and Aquatic Sciences* 82 (2024): 1-15. <https://doi.org/10.1139/cjfas-2024-0127>
- [5] Cerini, Francesco, Dylan Z. Childs, and Christopher F. Clements. "A predictive timeline of wildlife population collapse." *Nature Ecology & Evolution* 7, no. 3 (2023): 320-331. <https://doi.org/10.1038/s41559-023-01985-2>
- [6] Cianconi, Paolo, Batul Hanife, Francesco Grillo, Kai Zhang, and Luigi Janiri. "Human responses and adaptation in a changing climate: a framework integrating biological, psychological, and behavioural aspects." *Life* 11, no. 9 (2021): 895. <https://doi.org/10.3390/life11090895>
- [7] Cordoleani, Flora, C. C. Phillis, A. M. Sturrock, A. M. FitzGerald, A. Malkassian, G. E. Whitman, P. K. Weber, and R. C. Johnson. "Threatened salmon rely on a rare life history strategy in a warming landscape." *Nature Climate*

- Change* 11, no. 11 (2021): 982-988. <https://doi.org/10.1038/s41586-022-04788-w>
- [8] Ebi, Kristie L., Jennifer Vanos, Jane W. Baldwin, Jesse E. Bell, David M. Hondula, Nicole A. Errett, Katie Hayes et al. "Extreme weather and climate change: population health and health system implications." *Annual review of public health* 42, no. 1 (2021): 293-315. <https://doi.org/10.1146/annurev-publhealth-012420-105026>
- [9] Fricke, Evan C., Alejandro Ordonez, Haldre S. Rogers, and Jens-Christian Svenning. "The effects of defaunation on plants' capacity to track climate change." *Science* 375, no. 6577 (2022): 210-214. <https://doi.org/10.1126/science.abk3510> <https://doi.org/10.1016/j.cub.2025.07.046>
- [10] Lenton, Timothy M. "Tipping positive change." *Philosophical Transactions of the Royal Society B* 375, no. 1794 (2020): 20190123. <https://doi.org/10.1098/rstb.2019.0123>
- [11] Mahanta, Deepak Kumar, Ipsita Samal, J. Komal, Tanmaya Kumar Bhoi, Prasanta Kumar Majhi, and Mohammad Abbas Ahmad. "Understanding anthropogenic climate change, its consequences on insect pests, and techniques in forecasting and monitoring pest dynamics: A contemporary scenario." In *Climate change and insect biodiversity*, pp. 44-67. CRC Press, 2023.
- [12] Murali, Gopal, Takuya Iwamura, Shai Meiri, and Uri Roll. "Future temperature extremes threaten land vertebrates." *Nature* 615, no. 7952 (2023): 461-467. <https://doi.org/10.1038/s41586-022-05606-z>
- [13] Parratt, Steven R., Benjamin S. Walsh, Soeren Metelmann, Nicola White, Andri Manser, Amanda J. Bretman, Ary A. Hoffmann, Rhonda R. Snook, and Tom AR Price. "Temperatures that sterilize males better match global species distributions than lethal temperatures." *Nature Climate Change* 11, no. 6 (2021): 481-484. <https://doi.org/10.1038/s41558-021-01047-0>
- [14] Riddell, E. A., K. J. Iknayan, L. Hargrove, S. Tremor, J. L. Patton, R. Ramirez, B. O. Wolf, and S. R. Beissinger. "Exposure to climate change drives stability or collapse of desert mammal and bird communities." *Science* 371, no. 6529 (2021): 633-636. <https://doi.org/10.1126/science.abd4605>
- [15] Singh, Brajesh K., Manuel Delgado-Baquerizo, Eleonora Egidi, Emilio Guirado, Jan E. Leach, Hongwei Liu, and Pankaj Trivedi. "Climate change impacts on plant pathogens, food security and paths forward." *Nature Reviews Microbiology* 21, no. 10 (2023): 640-656. <https://doi.org/10.1038/s41579-023-00900-7>
- [16] Skendžić, Sandra, Monika Zovko, Ivana Pajač Živković, Vinko Lešić, and Darija Lemić. "The impact of climate change on agricultural insect pests." *Insects* 12, no. 5 (2021): 440. <https://doi.org/10.3390/insects12050440>

- [17] Soroye, Peter, Tim Newbold, and Jeremy Kerr. "Climate change contributes to widespread declines among bumble bees across continents." *Science* 367, no. 6478 (2020): 685-688. <https://doi.org/10.1126/science.aax8591>
- [18] Trisos, Christopher H., Cory Merow, and Alex L. Pigot. "The projected timing of abrupt ecological disruption from climate change." *Nature* 580, no. 7804 (2020): 496-501. <https://doi.org/10.1038/s41586-020-2189-9>
- [19] Waldvogel, Ann-Marie, Barbara Feldmeyer, Gregor Rolshausen, Moises Exposito-Alonso, Christian Rellstab, Robert Kofler, Thomas Mock et al. "Evolutionary genomics can improve prediction of species' responses to climate change." *Evolution letters* 4, no. 1 (2020): 4-18. <https://doi.org/10.1002/evl3.154>
- [20] Wallingford, Piper D., Toni Lyn Morelli, Jenica M. Allen, Evelyn M. Beaury, Dana M. Blumenthal, Bethany A. Bradley, Jeffrey S. Dukes et al. "Adjusting the lens of invasion biology to focus on the impacts of climate-driven range shifts." *Nature Climate Change* 10, no. 5 (2020): 398-405. <https://doi.org/10.1038/s41558-020-0768-2>