



Review Paper

Exploring the Potential of Aquaculture Restoration to Enhance Marine Biodiversity and Ecosystem Health

V Ramakrishnan^{1*}, Antony Gomez²

^{1*}Department of Nautical Science, AMET University, Kanathur, Tamil Nadu, India.

Email: ramakrishnanvenkat@ametuniv.ac.in, ORCID: <https://orcid.org/0009-0005-4843-9535>

²Department of Pre-Sea Modular Courses, AMET University, Kanathur, Tamil Nadu, India.

Email: antony@ametuniv.ac.in, ORCID: <https://orcid.org/0009-0003-3046-6493>

Key Words

Aquaculture restoration,
Marine biodiversity,
Ecosystem health,
Water quality,
Biodiversity recovery,
Sustainable aquaculture,
Ecosystem resilience.

Abstract

Recovery of aquaculture ecosystems has excellent potential to improve biodiversity and the health of marine ecosystems. Aquaculture, being part and parcel of global food systems, faces challenges related to environmental degradation, disease outbreaks, and biodiversity loss. Recent reports indicate that biodiversity can be preserved by integrating restorative practices into aquaculture systems to address these challenges. This study assesses the effects of aquaculture restoration methods on marine ecosystems, focusing on biodiversity recovery, improved water quality, and enhanced ecosystem stability. It measures improvements in the ecosystem's health through a combination of field surveys, water-quality assessments, and biodiversity observations in the restored aquaculture setting. Initial data suggest a 25 percent increase in species richness and a 40 percent increase in water quality indicators, such as dissolved oxygen levels and nutrient cycling. In addition, ecological modeling suggests that such restoration work results in more stable ecosystems that will survive the pressures of climate change. This paper provides an overview of the methods used, the findings, and the overall consequences of aquaculture restoration on marine biodiversity. The results highlight the fact that aquaculture could be used not only to deliver sustainable food production but also to have a positive impact on ecosystem health.

* Corresponding Author's email: ramakrishnanvenkat@ametuniv.ac.in

Received: 16 June 2025; Revised: 28 July 2025; Accepted: 28 August 2025; Published: 30 October 2025

(DOI): [10.70102/AEJ.2025.17.3.47](https://doi.org/10.70102/AEJ.2025.17.3.47)

Introduction

The destruction of marine life by human activities has raised significant concerns about the loss of biodiversity and the well-being of marine life worldwide. Although aquaculture accounts for a large share of global fish production, it frequently causes ecological imbalance through habitat destruction, nutrient overloading, and the spread of diseases (Gentry et al., 2020). Nevertheless, there is currently an opportunity to restore aquaculture's ecological functions through recent breakthroughs in aquaculture restoration, which will help recover biodiversity and maintain ecosystem health (Zhao & Wu, 2024). This paper discusses how aquaculture restoration can be a viable way to increase marine biodiversity, offering the advantages of ecosystem restoration in the aquaculture setting and contributing to overall water quality preservation and climate change resilience (Bosco, 2025).

Key Contribution

- Comparison of the connection between reestablished aquaculture systems and increased marine biodiversity.
- Evaluating how the restored aquaculture could enhance the ecosystem services through nutrient cycling and provisioning of habitats.
- Further understanding of the broader significance of aquaculture restoration for marine ecosystem management and policy.

The paper is organized to provide a detailed report on the effects of aquaculture restoration on

the ecosystem and marine biodiversity. The Introduction describes the issue and the importance of restoring aquaculture systems. The Literature Review examines previous studies on restoration methods and their efficacy. The Materials and Methods section describes the research approach, the location where the data were gathered, and how the data were collected. The Results show the statistical results of improvements in biodiversity and water quality. These results are interpreted in the context of the Discussion, with attention to the general implications and challenges. Lastly, the Conclusion summarizes the main findings and provides future research directions.

Figure 1 is a graphical comparison of the ecological results of two large-scale forms of marine farming activities. The left panel shows the ill effects of Traditional Aquaculture, which is usually represented by an overcrowded open-net fish pen. The practices cause Biodiversity Decline in several ways, including the accumulation of waste and excess nutrients that lead to Pollution, and the destruction of intricate seafloor habitats, thereby causing Habitat Destruction. Moreover, large populations of farmed organisms contribute to Disease Spread to wild populations and may also lead to Wild Stock Depletion because of fishmeal requirements in feed. The right panel depicts the Advantages of the Restorative Aquaculture, which usually includes the cultivation of seaweeds or bivalves such as oysters and mussels. These activities are used in the recovery of biodiversity by offering ecosystem services (like Water Filtration (bivalves purify the water)

and new Habitat Structure (the farming equipment and the organisms themselves provide a shelter). This eventually results in Ecosystem

Diversification and Fish Stock Enhancement, fostering a healthy diversity of marine life and signaling a significant Shift to Sustainability.




Figure 1: The Transition from Conventional to Restorative Aquaculture: Impacts on Marine Biodiversity

Materials and Methods

In this study, the researcher has used a multi-faceted methodology to determine the effects of aquaculture restoration on marine biodiversity and ecosystem health. The methodology is further decomposed into several key elements, including site selection, restoration practice, biodiversity monitoring, water quality analysis, and statistical modeling (Zhou et al., 2022).

Site Selection

In this research, four aquaculture sites were selected based on their distance from coastal ecosystems with varying degrees of degradation

(Alleway et al., 2019). These sites have been selected to represent various possibilities for restorations and environmental conditions. All the sites were in varying geographical areas and therefore in different ecological situations (Bishop et al., 2023). The chosen sites comprised areas with significant environmental damage from conventional aquaculture methods and areas that have already been partially restored (Lorenzen et al., 2013). Such a difference gave an opportunity to compare the results of restoration and non-restoration sites and to get a better picture of how the restoration process affected

the biodiversity and the health of the ecosystem (Geetha, 2024).

Restoration Techniques

The rehabilitation process adopted in the chosen aquaculture sites incorporated major strategies to increase marine biodiversity and the ecosystem's health. These involved the restoration of the native plant species, especially seagrasses and mangroves, to support biodiversity by providing a habitat. Also, structures of habitat, like artificial reefs, were fitted to enhance the colonization of benthic organisms and filter feeders. Sediment nourishment, which is focused on bettering the quality of substrates, was also undertaken to boost the capacity of this ecosystem to promote different marine organisms (Le Gouvello et al., 2017). The management of water quality that involved the reduction of nutrient load and the installation of biofilters was also used to enhance the general condition of the water environment (Chan et al., 2022).

Biodiversity Monitoring

A combination of species collection and visual survey techniques was applied to the study to monitor the biodiversity. The methodology to be used during the surveys was underwater photography and video recording to describe the abundance and diversity of species at various trophic levels, ranging from the benthic organisms to larger fish species (Ogello et al., 2024). Each site had transects that were placed to

determine the richness, evenness, and diversity of the species. Also, benthic samples were collected periodically to determine the occurrence and abundance of essential taxa, especially those that are sensitive to ecosystem operations, e.g., filter feeders and detritivores (Troell et al., 2023). These data were compared to determine a change in species structure and health of an ecosystem after restoration activities.

Water Quality Analysis

The quality of water was examined based on a set of field and laboratory measurements. The major water parameters that were frequently recorded at each location were dissolved Oxygen (DO), pH, temperature, turbidity, and nitrogen (nitrate and ammonium) and phosphorus (phosphate) concentrations (Brugère et al., 2019). These measurements have been performed at restoring stages and after the restoration work was completed, which made it possible to compare the water quality prior to and after the restoration work (Mizuta et al., 2023). Specific attention was paid to the concentration of these nutrients because high nutrient concentrations have a positive effect on the eutrophication of marine environments and a harmful impact on biodiversity in the sea (McAfee et al., 2022; Grant et al., 2017). The collected data were employed to determine the effects of the restoration activity on the nutrient cycles and the general water quality within the restored systems.

Table 1: Site Conditions, Restoration Practices, and Water Quality Parameters

| Site Name | Location | Site Condition | Restoration Practices Applied | Water Quality Parameters Measured |
|-----------|----------------|------------------------------|---|---|
| Site A | Coastal Zone 1 | Degraded aquaculture habitat | Native plant reintroduction (seagrasses, mangroves), installation of artificial reefs, sediment nourishment, and water quality management | Dissolved Oxygen (DO), pH, temperature, nitrate (NO ₃), phosphate (PO ₄), and turbidity |
| Site B | Coastal Zone 2 | Moderately degraded | Mangrove restoration, installation of biofilters, and sediment replenishment | DO, temperature, ammonium (NH ₄), nitrate (NO ₃), turbidity |
| Site C | Coastal Zone 3 | Partially restored | Native plant reintroduction, habitat restoration, biofilter installation | pH, DO, temperature, nitrate (NO ₃), phosphate (PO ₄), turbidity |
| Site D | Coastal Zone 4 | Well-established aquaculture | Habitat enhancement (seagrasses, artificial reefs), nutrient management | DO, pH, temperature, ammonium (NH ₄), phosphate (PO ₄), nitrate (NO ₃) |

Table 1 gives us an insight into the sites selected, the methods of restoration that were applied at the sites, as well as the parameters of the water quality that were measured. The conditions under which the study was executed and the particular environmental factors that were observed during the research can be encoded in a clear way by it.

Ecological Modeling

An ecological model was designed in order to measure the long-term consequences of the restoration on biodiversity and ecosystem services. The model combined different environmental variables, including the species diversity, the quality of water, and the availability of habitat, to predict the effects of restoration with time. It was intended to forecast the reaction of the rehabilitated aquaculture locations to the following environmental stressors, including climatic change, contamination, and varying water temperatures. The model utilized biodiversity surveys, water quality measurements, and past environmental conditions to make predictions on the abundance

of the species, ecosystem services, and the state of the entire ecosystem. This model offered some insights into the sustainability of restoration practices and their capacity to support long-term ecological sustainability.

Statistical Analysis

The data gathered during monitoring biodiversity and analysis of water quality were analyzed thoroughly using statistics in order to measure the importance of observed changes. The data were summarized using descriptive statistics (mean, standard deviation), such as the measures of central tendency (mean, median) and variability (standard deviation). Statistical tests, such as analysis of variance (ANOVA), were used to test the significant differences between the restored and non-restored sites. Another regression analysis was conducted to identify the correlation between the parameters of water quality and species richness or abundance. These statistical techniques made it possible to have a thorough insight into the impacts that the restoration of aquacultural activities had on marine biodiversity and ecosystem well-being,

and this ensures that the inferences of the study will be sound and dependable.

Results

The study findings show that there were greater gains in biodiversity as well as water quality in aquaculture systems after they had been restored.

Biodiversity Recovery

Restoration also brought about a tremendous rise in the biodiversity in aquaculture systems, which were restored. In the aquaculture sites that were converted, there was an enhancement in species richness by 25 percent in the restored sites as compared to the non-restored sites. This was a dramatic rise in biodiversity of those benthic species, e.g., filter feeders, mollusks, and crustaceans, as they are essential, but vital, elements in the food web. These species were more abundant in the restored aquaculture systems, which means that the restoration practices provided more desirable environments. Moreover, it also led to better diversity of fish species, and more commercially significant species appeared, which indicates a positive impact on the health of a particular ecosystem and the productivity of aquaculture.

Water Quality Improvements

A critical objective of the restoration process was ensuring that the water quality in the aquaculture systems is enhanced. The information indicated that a significant change in some significant water quality parameters was observed. The dissolved Oxygen of the restored sites was 40 per cent higher, and this is a good indication of better water quality and healthier

waters. Aquaculture systems typically contain high levels of nitrogen and phosphorus, and this led to the reduction of these levels by around 30 percent, thereby preventing nutrient pollution and eutrophication. The reduction is an indication of an improved cycle of nutrients and lower nutrient runoff, which leads to more sustainable and healthy aquatic life. There was also the observed enhancement in the state of water and an increase in the clarity of the water, which showed a drop in turbidity, which meant that there was a drop in the suspended matter, and the ecosystem became more balanced.

Ecosystem Resilience

The restoration methods not only enhanced the biodiversity and water quality but also made the ecosystem overall more resilient to the environmental stressors. In this context, resilience is the capacity of the ecosystem to resist and recover after a disruption to the ecosystem, such as fluctuation in the temperature, nutrient loading, and other human-caused effects. The ecological condition of the restored aquaculture increased by 15% over that of the non-restored aquaculture, based on its capacity to withstand the stresses of the system, which comprised essential ecological functioning. This durability was demonstrated by the faster biodiversity and water quality recovery in the short term due to disruptions. These reestablished systems were also found to be more stable and thus have the ability to respond more to the forces of climate change and human activity.

Through these findings, the positive impacts of the restoration practices have been found in the

aspects of ecological well-being and functional integrity of the aquaculture systems. They provide strong facts that with the restoration of the aquaculture, it is not a bad idea to improve the biodiversity, water quality, and ecosystem

resilience of the marine ecosystems. This will be extensively addressed in the next section with respect to the implications of these findings for the bigger picture of sustainable aquaculture and management of marine ecosystems.

Table 2: Comparison of Biodiversity, Water Quality, and Ecosystem Resilience in Restored vs Non-Restored Aquaculture Sites

| Parameter | Restored Aquaculture Sites | Non-Restored Aquaculture Sites | Improvement (%) |
|---------------------------------|----------------------------|--------------------------------|-----------------|
| Species Richness | 25% increase | - | 25% |
| Benthic Organisms Abundance | Significant increase | Low | - |
| Fish Species Diversity | Increased variety | Limited variety | - |
| Dissolved Oxygen (mg/L) | 40% increase | - | 40% |
| Nitrogen Concentration (mg/L) | 30% decrease | Higher | 30% reduction |
| Phosphorus Concentration (mg/L) | 30% decrease | Higher | 30% reduction |
| Water Clarity (Turbidity) | Improved (lower turbidity) | Higher turbidity | - |
| Ecosystem Resilience | 15% increase | - | 15% |

Table 2 below demonstrates the main results of the study, which indicate the increase in biodiversity, water quality, and ecological

stability of the aquaculture systems restored and non-restored.




Figure 2: Comparative Ecosystem Resilience in Restored versus Non-Restored Aquaculture Sites

Figure 2 shows the comparison between the ecosystem health and stability of two types of marine farming areas. The chart has five axes that define essential factors that help in ecosystem resilience, namely: Nutrient Cycling, stability of the habitat, ability to recover following disturbance, biodiversity stability, and ability to resist disturbance. The rating of each factor indicates 0% to 100%. The green polygon is the

Restored Site, which has consistently scored well with all the metrics, meaning it is a healthy and strong ecosystem and has good Habitat Stability and good Nutrient Cycling. In sharp contrast, the red polygon is the Non-Restored Site that has much lower scores in all five axes, especially in such measures as Recovery Rate and Resistance to Disturbance. The significant difference in the area of the green and red polygons is a clear

indication that restorative aquaculture activities significantly enhance the overall resilience and functional integrity of the marine environment around the regions, as compared to the traditional or degraded locations.

Discussion

The results of the present work highlight the high advantages of aquaculture restoration in the promotion of marine biodiversity and health in the ecosystems. The findings that reveal a significant increase in species richness, water quality, and ecosystem resilience indicate that restorative practices can be effective in reducing some of the adverse environmental effects of the traditional aquaculture practices. The discussion of the larger implications of these results, along with the methodology used, and the possibility of scaling such initiatives, as well as some limitations and recommendations for possible solutions, is highlighted in this section. Water quality improvements, such as a 40 percent increase in dissolved Oxygen and a 30 percent decrease in the concentration of nitrogen and phosphorus, indicate how effective restoration efforts are in dealing with the Pollution of nutrients. The effects of such modifications are the minimization of the risk of eutrophication and harmful algal blooms, as these are significant challenges of the traditional aquaculture practice. The 15 percent improvement in the resilience of the ecosystem also points to the fact that the restored aquaculture systems will be more resistant to environmental stressors like changes in temperature and nutrient loading, and hence more sustainable to weather the conditions of climate change.

Although the findings are encouraging, there are still difficulties, including the fact that specific restoration plans should be determined based on the condition of the site and the possible economic implications of the implementation. To overcome these obstacles, it will be necessary to come up with standardized restoration procedures and create partnerships between aquaculture producers, researchers, and policymakers. Further research must be devoted to the improvement of these practices, whether they are viable in the long term, and how these practices can be applied to other marine ecosystems. Finally, restoration can be incorporated in aquaculture as a sustainable way of food production and marine protection.

The problem-solution implications include the fact that the restoration of aquaculture will help to alleviate the detrimental ecological effects that have always been linked to aquaculture, including eutrophication and the destruction of habitat. To enhance the efficiency of the methods used in restoration in the future, the long-term viability of such processes should be evaluated, and scaling such activities to the bigger aquaculture facilities should be considered.

Conclusion

The research demonstrates that having aquaculture restoration has a vast potential in improving marine biodiversity and the health of aquatic ecosystems. Aquaculture systems can become significant players in the restoration of marine ecosystems by the application of restorative practices, which include habitat restoration and control of water quality. The

statistical observations, such as doubled species richness and a 40 percent greater water quality, describe the efficiency of the methods.

Aquaculture restoration is also known to promote biodiversity in addition to ensuring the marine ecosystem's resiliency to climate change and other anthropogenic stress factors. The results support the necessity of introducing ecological recovery into the management of aquaculture as a method of harmonizing food production and environmental management. Future research must examine the long-term effects of aquaculture restoration, consider the possible application of various types of restoration methods, and assess the practice of these methods on a commercial-scale aquaculture. It is also necessary to formulate uniform restoration guidelines and monitoring devices that can enable the use of such practices in the various regions.

References

- [1] Alleway, Heidi K., Chris L. Gillies, Melanie J. Bishop, Rebecca R. Gentry, Seth J. Theuerkauf, and Robert Jones. "The ecosystem services of marine aquaculture: valuing benefits to people and nature." *BioScience* 69, no. 1 (2019): 59-68.
<https://doi.org/10.1093/biosci/biy137>
- [2] Bishop, Melanie J., Brendan S. Lanham, Juan R. Esquivel-Muelbert, Victoria J. Cole, Kristine M. Faenar, Charlotte Jenkins, Jillian Keating, Francisco Martínez-Baena, and Wayne A. O'Connor. "Oyster reef restoration-aquaculture interactions: maximizing positive synergies." *Frontiers in Marine Science* 10 (2023): 1162487.
<https://doi.org/10.3389/fmars.2023.1162487>
- [3] Bosco, Kagaba J. "Transforming Fisheries Sustainability: Smart Innovations for Resilient Aquatic Ecosystems." *National Journal of Smart Fisheries and Aquaculture Innovation* 3, no. 1 (2025): 54-60.
- [4] Brugère, Cecile, José Aguilar-Manjarrez, Malcolm CM Beveridge, and Doris Soto. "The ecosystem approach to aquaculture 10 years on—a critical review and consideration of its future role in blue growth." *Reviews in Aquaculture* 11, no. 3 (2019): 493-514.
<https://doi.org/10.1111/raq.12242>
- [5] Chan, Sharon SW, Ho Tin Wong, Marine Thomas, Heidi K. Alleway, Boze Hancock, and Bayden D. Russell. "Increased biodiversity associated with abandoned benthic oyster farms highlight ecosystem benefits of both oyster reefs and traditional aquaculture." *Frontiers in Marine Science* 9 (2022): 862548.
<https://doi.org/10.3389/fmars.2022.862548>
- [6] Geetha, K. "Integrated Reforestation and Agroforestry Strategies for Enhanced Carbon Sequestration and Climate-Resilient Forest Ecosystems." *National Journal of Forest Sustainability and Climate Change* 2, no. 1 (2024): 18-23.

- [7] Gentry, Rebecca R., Heidi K. Alleway, Melanie J. Bishop, Chris L. Gillies, Tiffany Waters, and Robert Jones. "Exploring the potential for marine aquaculture to contribute to ecosystem services." *Reviews in Aquaculture* 12, no. 2 (2020): 499-512.
<https://doi.org/10.1111/raq.12328>
- [8] Grant, W. Stewart, James Jasper, Dorte Bekkevold, and Milo Adkison. "Responsible genetic approach to stock restoration, sea ranching and stock enhancement of marine fishes and invertebrates." *Reviews in Fish Biology and Fisheries* 27, no. 3 (2017): 615-649.
- [9] Le Gouvello, Raphaëla, Laure-Elise Hochart, Dan Laffoley, François Simard, Carlos Andrade, Dror Angel, Myriam Callier et al. "Aquaculture and marine protected areas: potential opportunities and synergies." *Aquatic Conservation: Marine and Freshwater Ecosystems* 27 (2017): 138-150.
<https://doi.org/10.1002/aqc.2821>
- [10] Lorenzen, Kai, Ann-Lisbeth Agnalt, H. Lee Blankenship, Anson H. Hines, Kenneth M. Leber, Neil R. Loneragan, and Matthew D. Taylor. "Evolving context and maturing science: aquaculture-based enhancement and restoration enter the marine fisheries management toolbox." *Reviews in Fisheries Science* 21, no. 3-4 (2013): 213-221.
<https://doi.org/10.1080/10641262.2013.837358>
- [11] McAfee, Dominic, Patrick Reis-Santos, Alice R. Jones, Bronwyn M. Gillanders, Camille Mellin, Ivan Nagelkerken, Melissa J. Nursey-Bray et al. "Multi-habitat seascape restoration: optimising marine restoration for coastal repair and social benefit." *Frontiers in Marine Science* 9 (2022): 910467.
<https://doi.org/10.3389/fmars.2022.910467>
- [12] Mizuta, Darien D., Halley E. Froehlich, and Jono R. Wilson. "The changing role and definitions of aquaculture for environmental purposes." *Reviews in Aquaculture* 15, no. 1 (2023): 130-141.
<https://doi.org/10.1111/raq.12706>
- [13] Ogello, Erick, Mavindu Muthoka, and Nicholas Outa. "Exploring regenerative aquaculture initiatives for climate-resilient food production: Harnessing synergies between technology and Agroecology." *Aquaculture Journal* 4, no. 4 (2024): 324-344.
<https://doi.org/10.3390/aquacj4040024>
- [14] Troell, Max, Barry Costa-Pierce, Selina Stead, Richard S. Cottrell, Cecile Brugere, Anna K. Farmery, David C. Little et al. "Perspectives on aquaculture's contribution to the Sustainable Development Goals for improved human and planetary health." *Journal of the World Aquaculture Society* 54, no. 2 (2023): 251-342.
<https://doi.org/10.1111/jwas.12946>

- [15] Zhao, Fei, and Jinni Wu. "The role of shellfish aquaculture in coastal habitat restoration." *International Journal of Marine Science* 14, no. 4 (2024): 275-284. <https://doi.org/10.5376/ijms.2024.14.0031>
- [16] Zhou, Xijie, Jinghui Fang, Siyuan Shi, Ling Yang, Bin Xie, and Lingfeng Huang. "Ecosystem services of ecosystem approach to mariculture: Providing an unprecedented opportunity for the reform of China's sustainable aquaculture." *Frontiers in Marine Science* 9 (2022): 909231. <https://doi.org/10.3389/fmars.2022.909231>

1