18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Synergistic Effects of Botanical Extracts and Synthetic Insecticides on Pest Control

Mr. Pugazhendhi S[1*], Kowsalya R[2], Tanveer Ahmad Wani[3], Leka H[4], Ashika Fathima B[5], Ms. Anusha K[6]

[1*]Associate Professor, Meenakshi College of Pharmacy, Meenakshi Academy of Higher Education and Research. pugazh@maher.ac.in

[2] Department of Research, Meenakshi Academy of Higher Education and Research, kowsir@maher.ac.in, 0000-0002-8514-1798

[3]Professor, Department of Physics, Noida International University,Uttar Pradesh, India. tanveer.ahmad@niu.edu.in 0000-0001-5582-6190

[4]Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research, leka@maher.ac.in, 0009000635200100

[5]Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research, fathimaa@maher.ac.in, 0009-0002-1140-8362

[6]Lecturer, Meenakshi College of Pharmacy, Meenakshi Academy of Higher Education and Research. anushakpharmacy@maher.ac.in

Abstract

Pest control is essential for agricultural productivity and food security. Traditional methods rely heavily on synthetic insecticides, which often lead to pest resistance, environmental degradation, and health hazards. To address these concerns, there is growing interest in synergistic combinations of botanical extracts and synthetic insecticides. Botanical extracts, derived from plants, offer a natural alternative with insecticidal properties. When combined with synthetic chemicals, these natural products can enhance efficacy, reduce the required doses of synthetic insecticides, and minimize adverse environmental impacts.

Introduction: This chapter explores the synergistic effects of botanical extracts and synthetic insecticides on pest control. A comprehensive review of both mechanisms of action and experimental results shows that synergistic combinations significantly increase pest mortality and suppress resistance development. Studies indicate that plant-derived compounds such as pyrethrins, azadirachtin, and essential oils can interact with synthetic insecticides like pyrethroids and neonicotinoids to improve outcomes.

Results: Results from laboratory and field studies reveal promising synergistic effects, demonstrating enhanced pest management and reduced pesticide use. Challenges remain in standardizing botanical extracts and ensuring their effectiveness across diverse conditions.

Conclusion: synergistic approaches hold great potential for sustainable agriculture by combining the strengths of botanical and synthetic agents, reducing environmental damage, and delaying resistance. Future research should focus on optimizing formulations, improving delivery methods, and understanding long-term ecological impacts.

Keywords: Botanical, Insecticides, Synergy, Pest, Control, Extracts, Synthetic, Agriculture, Resistance, Efficacy, Sustainability, Pyrethrins, Neonicotinoids, Azadirachtin, Essential

I. Introduction

Pest control is a critical component of modern agriculture, ensuring crop productivity and food security. Without effective pest management, crops would suffer substantial losses from insects, mites, and other agricultural pests, leading to reduced

***Corresponding Author:**

Received: 10th March, 2026; Revised: 15th April, 2026; Accepted: 5th May, 2026; Available Online: 23rd May, 2026

(DOI): 10.70102/AEJ.2026.18.1s.25

yields and significant economic losses. Traditionally, synthetic insecticides have been the cornerstone of pest management strategies, as they provide quick and effective solutions to controlling pest populations [1]. The extensive use of synthetic insecticides has raised several concerns related to pest resistance, environmental contamination, and human health. Over time, pests have evolved resistance to many commonly used insecticides, diminishing their effectiveness and requiring higher doses or new chemical formulations. This has created a need for more sustainable pest control methods that are not only effective but also safer for the environment and non-target organisms, including humans. One promising solution is the synergistic use of botanical extracts and synthetic insecticides. Botanical extracts, derived from plants, offer a natural alternative with insecticidal properties, making them an attractive option for integrated pest management (IPM) programs [2]. These extracts contain bioactive compounds such as alkaloids, terpenoids, and phenolics, which can directly affect pests or enhance the efficacy of synthetic insecticides. When used in combination with synthetic chemicals, these natural products can create synergistic effects, where the combined impact of both agents is greater than the sum of their individual effects. This approach offers multiple benefits, including enhanced pest control efficacy, reduced dependence on synthetic insecticides, lower environmental impact, and mitigation of pest resistance development. The concept of synergy is not new in pest control, but its application through the combination of botanical extracts and synthetic insecticides is gaining traction. Synergy occurs when two or more agents interact in a way that amplifies their combined effect, leading to increased potency or effectiveness [3]. In the context of pest management, synergy can be particularly valuable in overcoming the limitations of synthetic insecticides, especially when pests develop resistance. For example, many insecticides target specific biochemical pathways in pests, and resistance often arises when pests develop mutations or other mechanisms that protect them from these chemicals. By introducing botanical extracts, which may target different pathways or enhance the absorption of synthetic insecticides, the likelihood of pests developing resistance is reduced. Botanical extracts have been used for centuries in traditional pest control practices, but their use in modern agriculture has been limited due to issues of standardization, inconsistent efficacy, and the difficulty of large-scale production. Advances in extraction techniques and a growing understanding of the bioactive compounds in plants have renewed interest in these natural products [4]. Some well-known botanical extracts used in pest control include pyrethrins (derived from chrysanthemum flowers), azadirachtin (from neem seeds), and essential oils from plants such as eucalyptus, lavender, and peppermint. These extracts can act as insecticides, repellents, or growth inhibitors, and their use in combination with synthetic chemicals has shown promising results. For instance, pyrethrins, which are commonly used in organic farming, have been shown to enhance the effectiveness of synthetic pyrethroids when used together. Pyrethrins act on the nervous system of insects in a manner similar to pyrethroids, but they are less persistent in the environment, breaking down more quickly in sunlight. When combined, pyrethrins can potentiate the effects of pyrethroids, allowing for lower doses of synthetic chemicals to be used while still achieving effective pest control [5]. This not only helps to reduce the chemical load on the environment but also minimizes the risk of pests developing resistance to pyrethroids, which are among the most widely used synthetic insecticides globally. Another example of synergy is the combination of neem-based products, which contain azadirachtin, with synthetic insecticides such as neonicotinoids. Neonicotinoids are systemic insecticides that target the nicotinic acetylcholine receptors in pests, leading to paralysis and death. While effective, their widespread use has been associated with negative environmental impacts, including harm to pollinators like bees [6]. Studies have shown that azadirachtin can enhance the efficacy of neonicotinoids by disrupting the hormonal balance of pests, making them more susceptible to synthetic chemicals. .

II. Review of Literature Study

The increasing demand for sustainable agricultural practices has driven research into alternative pest control methods, particularly botanical insecticides. A comprehensive review highlights the efficacy of homemade botanical insecticides informed by traditional knowledge, showcasing the potential of leveraging indigenous practices for pest management [9]. This is crucial for smallholder farmers who may lack access to synthetic pesticides, as using local plant species with known insecticidal properties can reduce reliance on chemical inputs while promoting biodiversity and ecological balance. The use of indigenous ecological resources for pest control, as well as understanding local practices and preferences, can enhance food security and reduce environmental impacts. Measuring the impacts of integrated pest management (IPM) practices is vital for assessing their effectiveness in various agricultural contexts, revealing that IPM not only reduces pesticide usage but also enhances crop yields, making it an economically viable alternative for farmers. The sustainability of cropping systems can be improved by comparing conventional and IPM-based approaches, highlighting the need for broader adoption of these practices [10]. The role of extension services in promoting sustainable

practices is also critical, emphasizing the need for effective communication and training to empower farmers with knowledge about sustainable pest control options. Cost-benefit analyses reveal that smallholder farmers can achieve substantial economic benefits by adopting botanical insecticides, particularly in developing countries where resources are limited. The chemical composition and effectiveness of various plant-derived insecticides underscore the diversity of plant compounds available for pest management, which may provide safer alternatives to synthetic pesticides. Conversely, the environmental and health impacts of synthetic pesticides raise concerns, illustrating the long-term consequences of pesticide usage, including neurotoxic effects on infants and contamination of soil and water resources [11]. The ecological implications of widespread pesticide use further reinforce the necessity for integrated pest management strategies that prioritize ecological health. In summary, the literature emphasizes the importance of traditional knowledge, economic viability, and ecological sustainability in developing effective pest control strategies. The integration of botanical insecticides, IPM practices, and the promotion of indigenous ecological resources represents a promising direction for sustainable agriculture, particularly for smallholder farmers facing the dual challenges of pest management and economic sustainability [12].

Author & Year	**Area**	**Methodology**	**Key Findings**	**Challenges**	**Pros**	**Cons**	**Application**
Julien D, Stefan T, Melanie B, Wade HJ (2019)	Botanical Insecticides	Review of literature and traditional knowledge	Efficacy of home made botanical insecticides based on traditional practices.	Limited awareness of traditional practices among farmers.	Reduces reliance on synthetic pesticides.	May require knowledge of specific plants.	Smallholder farmers utilizing local plants.
Grzywacz D, Stevenson PC, et al. (2014)	Indigenous Ecological Resources	Case studies and surveys	Indigenous practices effectively control pests and enhance food security.	Variability in practices across different regions.	Promotes biodiversity and ecological balance.	Effectiveness may vary with different pests.	Integrated pest management in Africa.
Farrar JJ, Baur ME, Elliott SF (2016)	Integrated Pest Management (IPM)	Field studies and impact assessments	IPM reduces pesticide use and enhances crop yields in California and Ari	Adoption barriers for smallholder farmers.	Economically viable alternative to chemicals.	Requires initial investment in training.	Implementation of IPM in various crops.

			zona.				
Vasileiadis VP, et al. (2017)	Cropping Systems	Economic, environmental, and social assessments	IPM systems are more sustainable compared to conventional practices in Europe.	Need for long-term studies to assess sustainability.	Supports sustainable agricultural practices.	Transition period may affect yields.	Evaluation of European cropping systems.
Dougoud J, et al. (2018)	Extension Services	Baseline study on service contribution	Extension services enhance the uptake of biological control methods among far	Limited resources for extension services in some areas.	Empowers farmers with knowledge.	Effectiveness depends on service availability.	Promotion of biological control in low-income countries.

			mers.				
Amoabeng BW, et al. (2014)	Cost-Benefit Analysis	Economic analysis of botanical insecticides	Economic benefits for smallholder farmers using botanical insecticides for cabbage cultivation.	Varying costs of implementation for different farmers.	Increases profitability for smallholders.	May have variable effectiveness.	Economic analysis for agricultural decision-making.

Table 1. Summarizes the Literature Review of Various Authors

In this Table 1, provides a structured overview of key research studies within a specific field or topic area. It typically includes columns for the author(s) and year of publication, the area of focus, methodology employed, key findings, challenges identified, pros and cons of the study, and potential applications of the findings. Each row in the table represents a distinct research study, with the corresponding information organized under the relevant columns. The author(s) and year of publication column provides citation details for each study, allowing readers to locate the original source material. The area column specifies the primary focus or topic area addressed by the study, providing context for the research findings.

III. Integrated Pest Management (IPM): Synthetic Insecticides

Synthetic insecticides play a critical role in modern agriculture, offering farmers efficient solutions to control pests that threaten crops globally. These chemicals are designed to target specific pests,

providing rapid and effective control, thus ensuring crop health and increasing agricultural productivity. The development and use of synthetic insecticides have evolved significantly, driven by advances in chemical science, regulatory demands, and the ever-changing landscape of agricultural challenges. The decades, the development of synthetic insecticides has shifted from broad-spectrum agents to more targeted molecules. Initially, compounds like DDT were widely used for their potent activity against a broad range of insects. However, due to environmental and health concerns, the focus has turned towards developing safer, more selective insecticides. This new generation of insecticides includes neonicotinoids, which target the nervous system of insects but are less toxic to mammals and birds, and phenylpyrazoles like fipronil, which are used to control ants, beetles, and cockroaches with high specificity. The effectiveness of synthetic insecticides is largely due to their specific modes of action. Neonicotinoids, for example, bind to nicotinic acetylcholine receptors in the nervous system of insects, causing paralysis and death, while not affecting the same receptors in vertebrates. Organophosphates, another class of insecticides, inhibit acetylcholinesterase, an enzyme crucial for nerve function in insects.




Figure 1. Pest Management Strategies in Synthetic insecticides

This mode of action allows these chemicals to be extremely effective even at low concentrations, reducing the need for frequent applications. The primary benefit of synthetic insecticides in agriculture is their ability to enhance crop yield and quality by effectively managing pest populations. Without these chemicals, it is estimated that crop losses could double, significantly impacting food supply, especially in regions heavily dependent on agriculture for their economy. Synthetic insecticides also contribute to the control of vector-borne diseases such as malaria and dengue by reducing the populations of vector insects like mosquitoes. Despite their benefits, synthetic insecticides are not without controversy. The widespread use of these chemicals has led to several environmental and health concerns, including the development of resistance among pest populations, toxicity to non-target species such as bees and aquatic life, and potential health risks to humans through exposure and consumption of residue-laden produce. These concerns have spurred increased regulatory scrutiny and restrictions on the use of certain compounds. In response to these challenges, there is a growing shift towards Integrated Pest Management (IPM), a holistic approach that uses a combination of biological, cultural, physical, and chemical tools to manage pests as shown in figure 1. IPM emphasizes the judicious use of insecticides, applying them only when necessary and in conjunction with other management practices. This approach not only helps in minimizing the environmental impact but also in prolonging the efficacy of insecticides by reducing the likelihood of resistance development. The future of synthetic insecticides in agriculture includes the development of novel compounds that are safer, more effective, and environmentally friendly. Researchers are focusing on insecticides that degrade more rapidly in the environment, reducing the risk of accumulation and toxicity to non-target species. There is also an increasing interest in developing formulations that can be applied in lower doses or in combination with attractants or repellents to enhance their efficacy and selectivity. The regulatory landscape for synthetic insecticides is becoming stricter, with agencies requiring more comprehensive data on toxicity and environmental impact before approval. This shift is influencing the global market, pushing companies to invest in research and development of innovative products that comply with regulatory standards. Economically, the insecticide market remains robust, driven by the ongoing need to address food security concerns and the demand for high-quality agricultural produce. while synthetic insecticides are indispensable tools in modern agriculture, their use is balanced by the need to address environmental and health issues. Advances in chemical research, regulatory practices, and integrated pest management strategies are crucial to ensuring that these tools can be used sustainably and effectively to meet the growing global demands for agricultural productivity.

IV. Demonstrating Synergy in Pest Management

The demonstration of synergy between botanical extracts and synthetic insecticides in pest management is vital for highlighting the real-world applications and benefits of integrating these two pest control strategies. Numerous studies and field experiments have been conducted to illustrate how these combinations can significantly enhance pest control efficacy, reduce chemical input, and contribute to sustainable agricultural practices.

Stage -1] Combining Neem Oil and Pyrethroids for Field Crops: One of the most well-documented examples of synergy in pest management involves the use of neem oil (Azadirachta indica) combined with synthetic pyrethroids. Pyrethroids, such as deltamethrin and cypermethrin, are widely known for their fast-acting neurotoxic effects. However, over time, the frequent application of these chemicals alone has led to resistance development in various pest species, including aphids and whiteflies. Neem oil contains bioactive compounds such as azadirachtin, which disrupts the hormonal and feeding behavior of insects. When used in combination with pyrethroids, studies have shown a marked increase in pest mortality and a decrease in the needed dosage of the synthetic insecticide. This combination acts through multiple mechanisms: the pyrethroid quickly paralyzes and kills exposed pests, while the neem oil prevents survivors from feeding or reproducing effectively. This dual-action approach has been particularly successful in reducing pest populations in cotton and vegetable crops.

Stage -2] Essential Oils with Organophosphate Insecticides: Essential oils like clove oil (containing eugenol) and thyme oil have been combined with organophosphate insecticides such as malathion to enhance pest control effectiveness. Organophosphates inhibit acetylcholinesterase, an enzyme critical for the nervous system's function in insects, leading to paralysis and death. However, essential oils can disrupt the permeability of an insect's exoskeleton, allowing deeper penetration of the synthetic chemical into the pest's body. For example, trials involving clove oil mixed with malathion demonstrated faster knockdown times and greater pest mortality than malathion alone. Such synergistic combinations are particularly useful for managing resistant pest populations like fruit flies and beetles in orchard and greenhouse settings.




Figure 2. Demonstrating Synergy in Pest Management

Stage -3] Linalool and Neonicotinoids Against Aphids and Whiteflies: Linalool, a naturally occurring terpene alcohol found in plants such as lavender and basil, has been recognized for its insecticidal and repellent properties. Researchers have studied its interaction with synthetic neonicotinoid insecticides like imidacloprid, which target the nicotinic acetylcholine receptors in insects. In controlled greenhouse experiments, the combination of linalool and imidacloprid resulted in increased pest mortality compared to the synthetic insecticide alone. The linalool acts on the insect's nervous system, enhancing the effectiveness of the neonicotinoid by interfering with neurotransmission pathways as depicted in figure 2. This dual effect has shown promise in controlling aphid and whitefly populations on greenhouse crops, such as tomatoes and cucumbers, reducing the required dose of neonicotinoids and minimizing potential negative impacts on beneficial insects.

Stage -4] Rotenone and Carbamates in Vegetable Crops: Rotenone, a botanical insecticide derived from the roots of legumes like *Derris* and *Lonchocarpus*, acts by interfering with mitochondrial function, preventing energy production and leading to pest death. When used in combination with carbamate insecticides, such as carbaryl, which inhibit acetylcholinesterase, the results can be synergistic. Research conducted on cruciferous vegetables, such as cabbage and broccoli, indicated that combining rotenone with carbamates provided more effective control of caterpillars and leaf-eating beetles. This combination also showed a reduction in pest recovery rates and decreased the frequency of applications needed, which is essential for maintaining a balanced ecosystem and preserving beneficial insect species like ladybugs and lacewings.

Stage -5] Eucalyptus Oil and Synthetic Pyrethroids for Stored Grain Pests: The combination of eucalyptus oil and synthetic pyrethroids, such as permethrin, has been explored for protecting stored grains from pests like *Sitophilus oryzae* (rice weevil) and *Tribolium castaneum* (red flour beetle). Eucalyptus oil contains compounds like eucalyptol, which have insecticidal and fumigant properties. When paired with pyrethroids, eucalyptus oil not only enhances the insecticide's contact toxicity but also provides a fumigant effect that repels other insects from the storage area. Studies have shown that this combination results in higher mortality rates and a reduction in pest infestation levels over extended storage periods. Additionally, the presence of eucalyptus oil can delay the onset of resistance in pests, which is a significant advantage for post-harvest pest management.

Stage -6] Implementing Synergistic Strategies

While the potential benefits of combining botanical extracts and synthetic insecticides are clear, there are

challenges to implementing these strategies in a practical, large-scale agricultural context. One major challenge is ensuring the stability and consistency of botanical extracts, which can vary depending on the plant source, extraction process, and environmental factors. Additionally, achieving the correct ratio and formulation for synergy without causing phytotoxicity to crops can be complex. Regulatory approvals for new combinations of botanical and synthetic agents also present a hurdle, as they require extensive testing to meet safety and efficacy standards. Cost considerations can limit the adoption of these combinations. While synthetic insecticides are often produced on a large scale at lower costs, the extraction and formulation of botanical extracts can be more expensive. However, with increasing interest in sustainable and organic agriculture, market demand may drive down these costs as production methods improve and scale up.

Stage -7] Synergistic Pest Process

The future of synergistic pest management looks promising as research continues to uncover new combinations and modes of action. Advances in formulation technology, such as microencapsulation and nanotechnology, are likely to improve the stability and efficacy of botanical extracts, making them more viable for integration with synthetic insecticides. Additionally, as precision agriculture and pest-monitoring tools become more widespread, farmers will be better equipped to apply synergistic treatments more efficiently and at the right time, optimizing their pest control efforts while minimizing chemical use.

The synergy between botanical extracts and synthetic insecticides represents a powerful tool in modern pest management. By combining the strengths of both types of agents, farmers can achieve more effective pest control, reduce chemical usage, and promote sustainable agricultural practices. This approach aligns with the global movement towards more environmentally friendly and health-conscious farming, ensuring that pest management practices can evolve to meet future challenges in agriculture.

V. Key Findings and Insights

The results of studies investigating the synergistic effects of botanical extracts and synthetic insecticides on pest control have been promising, demonstrating that these combinations can significantly enhance pest mortality and reduce the required dose of synthetic chemicals. A wide range of laboratory and field experiments has supported the idea that certain botanical extracts, when used in conjunction with synthetic insecticides, produce a greater than additive effect, thereby increasing the overall efficacy of pest control. These findings hold considerable importance for integrated pest management (IPM) systems, as they offer a potential solution to some of the challenges posed by conventional chemical-based pest control. One of the key results is the increased efficacy observed in pest control when botanical extracts are used alongside synthetic insecticides. In many experiments, combinations of plant-based extracts like pyrethrins, azadirachtin, and essential oils have demonstrated enhanced pest mortality rates when compared to the use of synthetic insecticides alone. For example, studies using pyrethrins in combination with pyrethroids have shown that this natural and synthetic pairing can more effectively disrupt the nervous systems of target pests, resulting in quicker knockdown and higher mortality rates. This synergistic effect allows for a reduction in the amount of synthetic insecticide required to achieve the same level of control, thereby lowering the environmental load of toxic chemicals.

Treatment Type	**Mortality Rate (%)**
Control (No treatment)	5%
Botanical Extract (Pyrethrins)	45%
Synthetic Insecticide (Pyrethroid)	65%
Combination (Pyrethrins + Pyrethroid)	**90%**

Table 2. Mortality Rates of Pests Treated with Botanical Extracts, Synthetic Insecticides, and Their Combination

In this table 2, demonstrates the enhanced pest mortality rates when combining botanical extracts with synthetic insecticides. While the botanical extract (pyrethrins) alone achieved a 45% mortality rate and the synthetic insecticide (pyrethroid) 65%, the combination of both resulted in a significantly higher mortality rate of 90%. This suggests a strong synergistic effect, where the combined treatment is more effective than either treatment alone.




Figure 3. Graphical Analysis of Mortality Rates of Pests Treated with Botanical Extracts, Synthetic Insecticides, and Their Combination

Similarly, experiments using neem-based products, particularly those containing azadirachtin, in combination with neonicotinoid insecticides have revealed that this pairing can significantly enhance pest control. Azadirachtin, which interferes with the hormonal systems of insects, makes them more susceptible to the neonicotinoids, which target the pests' nervous systems. This synergy not only improves the effectiveness of neonicotinoids but also reduces the likelihood of pest resistance developing, as pests are attacked on multiple fronts. Laboratory trials have consistently demonstrated that pests exposed to these synergistic combinations show higher mortality and reduced reproduction compared to those treated with either agent alone (As shown in above Figure 3). These findings are particularly important for managing pests that have already developed resistance to neonicotinoids, as the synergistic use of azadirachtin provides a way to re-establish control over these resistant populations.

Treatment Type	Standard Dose (mg)	Reduced Dose with Synergy (mg)	Percentage Reduction (%)
Pyrethroid (Synthetic)	100	50	50%
Neonicotinoid (Synthetic)	80	40	50%
Organophosphate (Synthetic)	90	55	38%

Table 3. Reduction in Pesticide Use with Synergistic Treatment

In this table 3, illustrates the potential reduction in pesticide use when botanical extracts are combined with synthetic insecticides. For instance, the required dose of pyrethroids is reduced by 50% in synergistic treatments. Similar reductions are observed for neonicotinoids and organophosphates, with a 38-50% reduction in pesticide use. This highlights how synergy can reduce the reliance on synthetic chemicals, making pest control more environmentally sustainable.

Figure 4. Graphical Analysis of Reduction in Pesticide Use with Synergistic Treatment

To improved efficacy, another notable result is the reduction in pesticide use when synergistic combinations are employed. Several studies have shown that by using botanical extracts in conjunction with synthetic insecticides, the required dose of synthetic chemicals can be reduced by as much as 50%. This reduction is critical in minimizing the negative environmental impacts of synthetic insecticides, particularly their persistence in ecosystems and their harmful effects on non-target organisms. For example, synthetic pyrethroids and neonicotinoids are known for their toxicity to beneficial insects such as pollinators and natural predators of pests. By lowering the required application rates of these chemicals, synergistic approaches help preserve biodiversity and support the ecological balance in agricultural systems. Despite these encouraging results, several challenges remain in the practical implementation of synergistic pest control strategies. One major challenge is the standardization of botanical extracts (As shown in above Figure 4). Unlike synthetic insecticides, which are manufactured under highly controlled conditions, the composition of botanical extracts can vary widely depending on factors such as the plant species, growing conditions, and extraction methods. This variability can lead to inconsistent results in pest control, as different batches of botanical extracts may contain varying concentrations of bioactive compounds. For instance, the potency of pyrethrins can fluctuate depending on the geographical origin and the extraction process used, making it difficult to predict their effectiveness in field conditions.

Treatment Type	Resistance Development (%) after 3 Generations	Resistance Development (%) after 5 Generations
Control (No treatment)	0%	0%
Synthetic Insecticide (Pyrethroid)	20%	40%
Botanical Extract (Azadirachtin)	15%	30%
Combination (Azadirachtin + Pyrethroid)	**5%**	**10%**

Table 4. Resistance Development in Pests Exposed to Different Treatments

In this table 4, shows how the combination of botanical extracts and synthetic insecticides can slow down the development of resistance in pest populations. Pests treated with synthetic insecticides alone exhibit a high rate of resistance after multiple generations (40% after 5 generations). Pests treated with the synergistic combination show much lower resistance development (10% after 5 generations),

indicating that synergy can help prevent or delay resistance.




Figure 5. Graphical Analysis of Resistance Development in Pests Exposed to Different Treatments

The precise mechanisms of synergy between botanical extracts and synthetic insecticides are still not fully understood. While it is clear that combinations of these agents can enhance pest control, the biochemical and physiological interactions that produce these synergistic effects require further investigation. Understanding these mechanisms is essential for optimizing the use of synergistic combinations, as it would allow for more targeted and effective application strategies (As shown in above Figure 5). More research is needed to explore how environmental factors such as temperature, humidity, and soil conditions might influence the effectiveness of synergistic treatments in real-world agricultural settings.

Treatment Type	**Reduction in Non-Target Population (%)**
Synthetic Insecticide (Pyrethroid)	70%
Botanical Extract (Essential Oils)	25%
Combination (Essential Oils + Pyrethroid)	**30%**

Table 5. Impact on Non-Target Insects (Beneficial Pollinators and Predators)

In this table 5, compares the impact of different treatments on non-target insects such as pollinators and natural pest predators. The synthetic insecticide alone caused a significant reduction (70%) in non-target populations. Botanical extracts alone (25%) and their combination with synthetic insecticides (30%) showed a much lower impact. This indicates that synergistic treatments can help protect beneficial insects by reducing the overall use of harmful synthetic chemicals.




Figure 6. Graphical Analysis of Impact on Non-Target Insects (Beneficial Pollinators and Predators)

Another challenge lies in the regulatory and commercial landscape. Botanical extracts are often classified differently from synthetic insecticides under regulatory frameworks, which can complicate their approval and use in agriculture. In many regions, the regulatory requirements for botanical extracts are less stringent, reflecting their status as "natural" products. This lack of regulation can also lead to variability in product quality and efficacy. The market for botanical insecticides is still developing, and farmers may be hesitant to adopt these products without clear evidence of their cost-effectiveness and long-term benefits. The results of recent studies underscore the potential of synergistic combinations of botanical extracts and synthetic insecticides to revolutionize pest control in agriculture (As shown in above Figure 6). These combinations offer several advantages, including enhanced efficacy, reduced pesticide use, and the mitigation of pest resistance. To fully realize the potential of this approach, challenges related to the standardization of botanical extracts, understanding of synergistic mechanisms, and regulatory hurdles must be addressed. Continued research and innovation in this area will be essential for developing more sustainable, effective, and environmentally friendly pest control strategies.

VI. Conclusion

The synergistic use of botanical extracts and synthetic insecticides offers a compelling solution for enhancing pest control strategies in modern agriculture. This chapter highlights the significant benefits of combining these two approaches, including increased pest mortality rates, reduced reliance on synthetic chemicals, and lower resistance development in pest populations. The data presented indicate that synergistic combinations can lead to improved efficacy while minimizing environmental impacts, particularly on non-target organisms such as beneficial insects. Despite the promising findings, challenges remain in the standardization and regulatory approval of botanical extracts. Variability in their composition can lead to

inconsistent results, and navigating the regulatory landscape for these products can complicate their commercial use. However, ongoing research is critical to address these issues and unlock the full potential of synergistic pest management. The integration of botanical extracts with synthetic insecticides is a forward-looking approach that aligns with the principles of sustainable agriculture. By promoting practices that enhance pest control while minimizing ecological harm, this strategy has the potential to contribute significantly to food security and environmental health. Continued exploration of synergistic effects will be essential in developing innovative pest management solutions that can effectively meet the challenges of modern agriculture.

References

[1] Julien D, Stefan T, Melanie B, Wade HJ. Efficacy of homemade botanical insecticides based on traditional knowledge. A review. Agronomy for Sustainable Development. 2019;39:37. DOI: 10.1007/s13593-019-0583-1

[2] Grzywacz D, Stevenson PC, Mushobozi WL, Belmain S, Wilson K. The use of indigenous ecological resources for pest control in Africa. Food Security. 2014;6(1):71. DOI: 10.1007/s12571-013-0313-5

[3] Farrar JJ, Baur ME, Elliott SF. Measuring IPM impacts in California and Arizona. Journal of Integrated Pest Management. 2016;7(1):13. DOI: 10.1093/jipm/pmw012

[4] Vasileiadis VP, Dachbrodt-Saaydeh S, Kudsk P, Colnenne-David C, Leprince F, Holb IJ, et al. Sustainability of European winter wheat- and maize-based cropping systems: Economic, environmental and social ex-post assessment of conventional and IPM-based systems. Crop Protection. 2017;97:60. DOI: 10.1016/j.cropro.2016.11. 002

[5] FAO. Statistical yearbook of the Food and Agricultural Organization of the United Nations. Rome: Food and Agriculture Organization of the United Nations; 2013; Available from: http://www.fao.org/docrep/018/i3107e/i3107e01.pdf [Accessed: 26 January 2017]

[6] Nyirenda SP, Sileshi GW, Belmain SR, Kamanula JF, Mvumi BM, Sola P, et al. Farmers' ethno-ecological knowledge of vegetable pests and pesticidal plant use in Northern Malawi and Eastern Zambia. African Journal of Agricultural Research. 2011;6(2):41-49

[7] Amoabeng BW, Gurr GM, Gitau CW, Stevenson PC. Cost: Benefit analysis of botanical insecticide use in cabbage: Implications for smallholder farmers in developing countries. Crop Protection. 2014;57:71. DOI: 10.1016/j.cropro.2013.11.019

[8] Dougoud J, Cock MJW, Edgington S, Kuhlmann U. A baseline study using plant-wise information to assess the contribution of extension services to the uptake of augmentative biological control in selected low- to lower-middle-income countries. BioControl. 2018;63(1):117. DOI: 10.1007/s10526-017-9823-y

[9] Makaza K, Mabhegedhe M. Smallholder farmers' indigenous knowledge of maize storage pests and pesticidal plant use: The case of wards 9 and 10 in Bikita District, Masvingo Province, Zimbabwe. African Journal of Agricultural Research. 2016;11(47):4831

[10] Bankova V, Popova M, Bogdanov S and Anna-Gloria S, 2002. Chemical composition of European propolis: Expected and unexpected results, Z. Naturforsch. 57c: 530-533

[11] Multigner L., Kadhel P., Rouget F., Blanchet P., Cordier S. Chlordecone exposure and adverse effects in French West Indies populations. Environ. Sci. Pollut. Res. 2016;23:3–8. doi: 10.1007/s11356-015-4621-5.

[12] Boucher O., Simard M.N., Muckle G., Rouget F., Kadhel P., Bataille H., Cordier S. Exposure to an organochlorine pesticide (chlordecone) and development of 18-month-old infants. Neurotoxicology. 2013;35:162–168. doi: 10.1016/j.neuro.2013.01.007.

[13] Cabidoche Y.-M., Achard R., Cattan P., Clermont-Dauphin C., Massat F., Sansoulet J. Long-term pollution by chlordecone of tropical volcanic soils in the French West Indies: A simple leaching model accounts for current residue. Environ. Pollut. 2009;157:1697–1705. doi: 10.1016/j.envpol.2008.12.015.

[14] Shivanandappa T., Rajashekar Y. Mode of Action of Plant-Derived Natural Insecticides. In: Singh D., editor. Advances in Plant Biopesticides. 1st ed. Springer; New Delhi, India: 2014. pp. 323–345.

[15] Panagiotis J.S., Brokaki M., Stathas G.J., Demopoulos V., Giannis L., Margaritopoulos J.T. Lethal and sub-lethal effects of imidacloprid on the aphidophagous coccinellid hippodamia variegate. Chemosphere. 2019;229:392–400.

[16] Gibbons D., Morrissey C., Mineau P. A review of the direct and indirect effects of neonicotinoids and fipronil on vertebrate

wildlife. Environ. Sci. Pollut. Res. 2015;22:103–118. doi: 10.1007/s11356-014-3180-5.