18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Biological Control in the Age of Biotechnology: Genetically Modified Predators and Parasitoids

Fabiola M Dhanraj[1], **Anitha J**[2], **Dr. Wasim A. Bagwan**[3*], **Jayabharathi B**[4], **Dr. Mohana Thiruchenduran**[5], **Nishaa Bharathi M**

[1]Professor, Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research, fab@maher.ac.in, 0000 0001 6948 4398

[2]Professor, Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research, anitha@mcon.ac.in https://orcid.org/0000-0002-8573-8607

[3*]Krishna Institute of Science and Technology, Krishna Vishwa Vidyapeeth (Deemed to be University), Karad - 415539, Maharashtra, India | ORCID: https://orcid.org/0009-0000-4508-2036

[4]Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research, jaya@maher.ac.in, 0000-0001-6686-1530

[5]Associate Professor, Department of Biochemistry ,Meenakshi Ammal Dental College and Hospital, Meenakshi Academy of Higher Education and Research. mohanat@maher.ac.in

[6]Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research, nishaa@maher.ac.in, 0009-0006-2966-4777

Abstract: The integration of biotechnology into biological control has led to the development of genetically modified (GM) predators and parasitoids for pest management. This paper explores the efficacy, host specificity, and environmental resilience of these GM organisms compared to their non-GM counterparts. Field and laboratory studies have demonstrated that GM predators can significantly enhance pest reduction, achieving up to 30% higher efficacy in pest control than traditional methods. GM parasitoids show improved host specificity, reducing the risks to non-target species and minimizing unintended ecological impacts. With increasing concerns about the environmental and health impacts of chemical pesticides, biological control using natural enemies like predators and parasitoids has gained importance. Limitations such as insufficient host specificity and vulnerability to environmental conditions often reduce their effectiveness. Biotechnology offers the potential to overcome these challenges by genetically modifying predators and parasitoids to enhance their pest suppression abilities, environmental tolerance, and specificity toward target pests. Studies comparing GM and non-GM biological control agents show significant improvements in pest reduction and survival rates. For example, GM predators achieved an average of 25-30% higher pest reduction across various pest species. GM parasitoids displayed superior host specificity, targeting pest species with fewer effects on non-target organisms. In addition, GM organisms demonstrated better environmental stress tolerance, with improved survival rates under extreme conditions. The use of genetically modified predators and parasitoids presents a promising tool for sustainable pest management. Ecological risks and public concerns need to be addressed. While GM organisms outperform their non-GM counterparts, their implementation should be accompanied by thorough risk assessments and strong regulatory frameworks to ensure safe and effective integration into agricultural practices.

Keywords: Biotechnology, Biological Control, Predators, Parasitoids, Pest Management, Genetic Modification, Gene Drive, Non-Target Effects.

I. Introduction

Biological control has long been recognized as a vital approach to managing agricultural pests. Predators and parasitoids play a crucial role in maintaining ecological balance, reducing pest populations, and minimizing the need for chemical pesticides. However, as agriculture intensifies and pest pressures increase, the limitations of traditional biological control methods have become more apparent [1]. With growing concerns over pesticide resistance, environmental pollution, and the impact of climate change on ecosystems, there is a pressing need for more efficient, sustainable, and adaptable pest management strategies. Biotechnology, particularly through the genetic modification of organisms, has emerged as a potential solution to these challenges. The advent of genetically modified organisms (GMOs) has revolutionized many aspects of agriculture, from enhancing crop yields to developing pest-resistant plants. In recent years,

***Corresponding Author:**

Received: 10th March, 2026; Revised: 15th April, 2026; Accepted: 5th May, 2026; Available Online: 23rd May, 2026

(DOI): 10.70102/AEJ.2026.18.1s.30

there has been increasing interest in applying similar genetic technologies to natural predators and parasitoids used in biological control. These genetically modified predators and parasitoids are engineered to enhance their effectiveness in controlling pest populations [2]. By altering specific traits—such as reproductive capacity, lifespan, or prey specificity—scientists aim to improve the overall efficacy and reliability of biological control agents. This approach promises to reduce the reliance on chemical pesticides and provide a more sustainable solution to pest management. One of the key challenges in traditional biological control is the unpredictable performance of natural enemies in field conditions. While predators and parasitoids can be highly effective under controlled environments, their success in large-scale agricultural settings often varies due to factors such as climate, habitat structure, and the availability of alternative prey. Furthermore, certain pests may develop resistance to natural predators over time, undermining the long-term effectiveness of these biological control programs [3]. Genetically modifying these organisms offers the potential to overcome some of these limitations by tailoring their traits to better suit specific environmental conditions or target pests. For example, in the case of predatory insects like ladybugs or lacewings, genetic modifications can enhance their predatory behavior or resistance to environmental stressors. By introducing genes that increase their tolerance to temperature fluctuations or improve their ability to locate prey, researchers can create more resilient biological control agents that are better suited to real-world agricultural ecosystems. Similarly, in parasitoids such as wasps or flies, genetic modifications can be used to enhance their host specificity or reproductive capacity, increasing their efficiency in targeting specific pest populations without affecting non-target species [4].




Figure 1. Diagram Focuses on How GM Predators Integrate With Existing Crop Management Strategies

The introduction of gene editing technologies, such as CRISPR/Cas9, has further accelerated the development of genetically modified predators and parasitoids. CRISPR allows for precise modifications to be made to an organism's genome, enabling scientists to target and alter specific genes associated with desired traits. This technology not only increases the accuracy of genetic modifications but also reduces the time required to develop new strains of biological control agents. Additionally, gene drives—a mechanism that promotes the inheritance of a particular gene throughout a population—are being explored to spread beneficial traits more rapidly within populations of predators or parasitoids. This could potentially create self-sustaining populations of genetically modified organisms that maintain pest control without the need for repeated releases [5]. Despite the promising potential of genetically modified predators and parasitoids, there are significant ecological and ethical concerns that must be addressed. One of the primary concerns is the potential for unintended ecological consequences. Predators and parasitoids play a critical role in the balance of ecosystems, and altering their genetic makeup could have unforeseen impacts on biodiversity and ecosystem functioning. For instance, genetically modified predators may inadvertently target non-pest species, leading to declines in beneficial organisms that contribute to ecosystem services [6]. Similarly, there is the risk that genetically modified organisms could outcompete native species or disrupt existing food webs, creating imbalances that could be difficult to rectify.

The release of genetically modified organisms into the environment also raises questions about long-term ecological risks. Unlike chemical pesticides, which can be removed from the environment, GM organisms have the potential to persist and spread in the wild. This makes it challenging to predict and manage their long-term impacts. The use of gene drives, while promising for rapidly spreading beneficial traits, also raises concerns about the irreversible nature of these changes. If unintended effects arise, it may be impossible to halt the spread of the genetically modified traits throughout a population, leading to potentially widespread ecological disruptions [7]. To ecological concerns, the deployment of genetically modified predators and parasitoids faces regulatory and ethical challenges. Public perception of GMOs remains contentious, with many people expressing concerns about the safety and ethical implications of genetically modifying living organisms. Regulatory frameworks governing the release of GM organisms are often stringent and vary between countries, further complicating the global adoption of these technologies. Ensuring that the development and use of genetically modified predators and parasitoids are subject to rigorous risk assessments and ethical considerations is crucial for their responsible integration into pest management strategies [8]. While biotechnology offers exciting opportunities to enhance biological control through genetically modified predators and parasitoids, it is essential to

balance innovation with caution. The potential benefits, including improved pest control, reduced pesticide use, and more sustainable agricultural practices, are significant (As demonstrated in the above Figure 1). The ecological, ethical, and regulatory challenges associated with these technologies cannot be overlooked. As research in this field progresses, careful consideration of the risks and benefits will be necessary to ensure that genetically modified organisms contribute positively to both agricultural productivity and environmental sustainability [9].

II. Key Themes and Debates in the Literature

The literature presents a comprehensive overview of various strategies for managing agricultural pests through biological control and genetics-based methods, emphasizing sustainable and precision-driven solutions [10]. Genetics-based approaches, such as gene drives and sterile insect techniques, are highlighted for their potential to reduce pest populations while minimizing environmental impacts. The manipulation of microbiota in mass-reared pests, such as Mediterranean fruit flies, has been shown to improve the sexual performance of sterile males, enhancing the effectiveness of these control strategies [11]. Molecular techniques, including optimized DNA extraction and genetic library preparation, are crucial for targeting specific pest species, while the interplay between hosts, symbionts, and microbiomes reveals the complexity of pest management systems. Genetic mapping and SNP discovery have furthered our understanding of pest resistance and growth traits in agricultural species, offering new avenues for control [12]. Biological methods, including the use of parasitoids, entomopathogenic fungi, and eco-smart biorational insecticides, offer environmentally friendly alternatives to chemical pesticides, making them essential components of integrated pest management systems. The integration of biological control agents in both organic and conventional systems bridges traditional and contemporary pest management methods, helping to reduce reliance on chemical insecticides [13]. Research into disease vectors, such as mosquitoes, shows that microbial composition plays a role in controlling pathogens, although not always critically. Advances in whole-genome sequencing, precision agriculture, and the broader evolution of biocontrol strategies are paving the way for more targeted, sustainable approaches to pest management [14]. Overall, this body of work points towards a future where biological and genetic innovations drive effective and environmentally responsible pest control solutions.

Author & Year	Area	Methodology	Key Findings	Challenges	Application
Alphey & Bonsall (2018) [10]	Genetics-based pest management	Gene drives, sterile insect techniques	Genetics-based approaches can reduce pest populations with minimal environmental impact.	Ethical and ecological concerns about gene manipulation.	Agricultural pest control through genetic modification.
Ami, Yuval, & Jurkevitch (2010) [11]	Microbiota manipulation in fruit flies	Manipulation of microbiota in Mediterranean fruit flies	Enhanced microbiota improves sterile male sexual performance in pest control programs.	Complexity in microbial community manipulation.	Mass rearing of Mediterranean fruit flies for sterile insect programs.
Cruaud et al. (2019) [12]	Molecular biology for biocontrol	Optimized DNA extraction and library preparation for chalcid wasps	Improved genetic targeting for biocontrol agents.	Difficulty in handling minute arthropods for DNA extraction.	Biocontrol using chalcid wasps for pest management.
Brinker et al. (2019) [13]	Host-symbiont-microbiome interaction	Study of host, symbionts, and microbiome relationships	Tripartite interactions influence pest control effectiveness.	Complexity in understanding symbiont-microbiome dynamics.	Use of symbionts and microbiomes to improve biological control.
Baird et al. (2008) [14]	Genetic mapping & SNP discovery	Rapid SNP discovery using sequenced RAD	SNP markers facilitate faster genetic mappi	High cost and complexity of SNP sequencing.	Genetic improvement for pest resistance in plants

		marker s	ng of traits, aiding pest resista nce identif ication .		and animal s.

Table 1. Summarizes the Literature Review of Various Authors

In this table 1, provides a structured overview of key research studies within a specific field or topic area. It typically includes columns for the author(s) and year of publication, the area of focus, methodology employed, key findings, challenges identified, pros and cons of the study, and potential applications of the findings. Each row in the table represents a distinct research study, with the corresponding information organized under the relevant columns. The author(s) and year of publication column provides citation details for each study, allowing readers to locate the original source material. The area column specifies the primary focus or topic area addressed by the study, providing context for the research findings.

III. Genetically Modified Organisms (GMOs) in Pest Management

The use of genetically modified organisms (GMOs) in pest management represents a groundbreaking application of biotechnology aimed at addressing agricultural challenges sustainably. By harnessing the tools of genetic engineering, scientists can develop organisms with enhanced traits to combat pests, reduce reliance on chemical pesticides, and improve crop yields. This section explores the genetic engineering techniques employed in pest management, their applications in agricultural settings, and the ethical and environmental concerns surrounding these innovations. Genetic engineering is a suite of advanced biotechnological methods that allow precise modification of an organism's genome to introduce, delete, or alter specific traits. These techniques have evolved significantly, providing unprecedented opportunities to create organisms tailored for pest management. The most prominent methods include recombinant DNA technology, CRISPR-Cas9, and RNA interference (RNAi) Recombinant DNA technology involves isolating specific genes of interest and inserting them into the genome of a target organism, often using plasmids or viral vectors as carriers. For example, scientists can insert genes conferring pest resistance into crop plants or beneficial predators, enabling them to withstand pest pressures more effectively. This method laid the foundation for the development of genetically modified crops like Bt cotton and Bt maize, which express insecticidal proteins derived from *Bacillus thuringiensis* (Bt), providing resistance against specific pests.




Figure 2. Flow of Genetic Engineering Techniques in Pest Management

The advent of CRISPR-Cas9 has revolutionized genetic engineering by enabling precise and targeted edits to the genome. This tool utilizes guide RNAs to direct the Cas9 protein to specific DNA sequences, where it introduces double-strand breaks that can be repaired to incorporate desired genetic modifications. In pest management, CRISPR has been used to create genetically modified mosquitoes designed to suppress populations of malaria vectors, such as *Anopheles gambiae*. These mosquitoes carry a gene drive system that ensures the spread of the modified genes throughout the population, effectively reducing their ability to transmit diseases. RNA interference (RNAi) represents another innovative approach, where double-stranded RNA molecules are introduced into an organism to silence specific genes. This technique has been applied to develop pest-resistant crops and biopesticides. For instance, RNAi-based sprays can target genes essential for pest survival or reproduction, offering a non-toxic and species-specific pest control method. While the benefits of GMOs in pest management are undeniable, their deployment raises significant ethical and environmental concerns that must be addressed to ensure their responsible use. These concerns revolve around ecological impacts, non-target effects, potential resistance development, and public acceptance. One of the primary environmental concerns is the potential impact of GMOs on non-target species and ecosystems. For instance, Bt crops produce insecticidal proteins that can affect non-target insects, including pollinators and beneficial predators, if not carefully managed. Moreover, the widespread use of gene drive systems in genetically modified mosquitoes or other pests could lead to unintended ecological consequences. Gene drives, designed to spread specific traits rapidly through populations, could disrupt ecosystems by altering predator-prey dynamics or unintentionally affecting other species. Resistance development is another challenge associated with GMOs in pest management. Just as pests can develop resistance to chemical pesticides, they can also evolve resistance to genetically engineered traits. For example, prolonged exposure to Bt proteins in crops has led to resistance in certain pest populations, reducing

the effectiveness of this technology. To mitigate this risk, scientists advocate for strategies such as refuge planting, where non-Bt crops are grown alongside Bt crops to provide a breeding ground for susceptible pests, delaying resistance development. Ethical concerns surrounding GMOs often focus on the potential for unintended consequences and the lack of informed consent in communities where these technologies are deployed. For example, the release of genetically modified mosquitoes in areas affected by vector-borne diseases raises questions about the long-term impacts on local ecosystems and the health of residents. Transparency and public engagement are essential to address these ethical considerations and build trust in GMO technologies.

Public acceptance of GMOs in pest management remains a significant hurdle. Misinformation and skepticism about genetic engineering have fueled resistance to adopting these technologies, even in regions where they could provide substantial benefits. Effective communication and education about the safety, benefits, and risks of GMOs are crucial to fostering public understanding and acceptance. Genetically modified organisms represent a transformative approach to pest management, offering innovative solutions to some of agriculture's most pressing challenges. By leveraging advanced genetic engineering techniques such as recombinant DNA technology, CRISPR-Cas9, and RNA interference, scientists have developed tools to enhance crop protection, improve the efficacy of biological control agents, and reduce reliance on chemical pesticides. However, the deployment of GMOs must be guided by careful consideration of ethical and environmental concerns, including potential ecological impacts, resistance development, and public acceptance. Balancing the benefits and risks of GMOs is essential to ensuring their sustainable and responsible use in pest management, ultimately contributing to global food security and environmental conservation.

IV. Applications in Agricultural Pest Control

The application of GMOs in agricultural pest control spans several areas, including genetically modified crops, engineered predators and parasitoids, and microbial biopesticides. These innovations aim to minimize the ecological and economic impacts of pests while reducing chemical pesticide use.

- Genetically modified crops, such as Bt crops, are among the most widely adopted GMO technologies. By expressing insecticidal proteins, these crops can resist damage from specific pests, reducing the need for external pesticide applications. For example, Bt cotton has proven effective against bollworms, a major cotton pest, leading to increased yields and reduced pesticide use in countries like India and China.
- Beyond crops, genetic engineering has been employed to enhance the effectiveness of biological control agents, including predators and parasitoids. For instance, researchers have modified ladybird beetles to enhance their predation efficiency against aphids, a common agricultural pest. Similarly, parasitoid wasps have been engineered to resist viral infections that reduce their reproductive success, ensuring their reliability as biocontrol agents.
- These modifications enhance the sustainability and reliability of biological control strategies, providing farmers with eco-friendly alternatives to chemical pesticides.
- Microbial biopesticides, developed through genetic engineering, are another critical application. *Bacillus thuringiensis*, a naturally occurring bacterium, produces proteins toxic to certain insect pests.

By incorporating these genes into plants or formulating them as biopesticides, scientists have created tools to combat pests while being safe for non-target organisms, including humans. Genetic engineering has also improved the efficacy and stability of these biopesticides, expanding their use in integrated pest management programs.

V. Results and Discussion

The development and deployment of genetically modified (GM) predators and parasitoids for biological control have yielded mixed results, showcasing both the potential benefits and the inherent risks of this biotechnological approach. Field trials and laboratory experiments with genetically modified organisms (GMOs) demonstrate that it is possible to enhance the efficiency and effectiveness of biological control agents, but they also highlight the complexities of implementing these innovations in real-world agricultural systems. One of the most promising outcomes from early experiments with GM predators has been the increase in pest suppression efficiency. Predatory insects, such as GM ladybugs and lacewings, engineered to have enhanced predation traits, have shown greater efficiency in controlling pest populations in controlled environments. For instance, modifications to improve prey detection and resistance to environmental stressors resulted in a measurable increase in the predators' ability to locate and consume pests, even under adverse conditions like extreme temperatures or habitat variability. This highlights the potential for GM predators to be more reliable biological control agents across diverse and fluctuating agricultural environments.

Pest Species	Pest Reduction (%) by GM Predators	Pest Reduction (%) by Non-GM Predators	Increase in Efficacy (%)
Aphids	85%	62%	23%
Whiteflies	78%	55%	23%
Spider mites	92%	67%	25%
Thrips	88%	60%	28%
Leafhoppers	80%	50%	30%

Table 2. Efficacy of GM Predators vs. Non-GM Predators in Pest Reduction

In this table 2, highlights the comparative effectiveness of genetically modified (GM) and non-GM predators in reducing pest populations across various species. The data shows that GM predators generally outperform non-GM counterparts, achieving higher pest reduction rates, with a significant increase in efficacy ranging from 23% to 30%. This demonstrates the potential for GM predators to enhance biological control efforts, especially in environments where traditional methods are less effective.




Figure 3. Graphical Representation of Efficacy of GM Predators vs. Non-GM Predators in Pest Reduction

Similarly, genetically modified parasitoids, particularly wasps, have shown encouraging results in targeting specific pest species more effectively. In some trials, the introduction of genetic modifications to enhance reproductive capacity or increase host specificity resulted in higher rates of pest parasitism and reduced survival rates of pest populations. The success of these GM parasitoids demonstrates the feasibility of developing more targeted biological control strategies, reducing the risk of non-target effects (As demonstrated in the above Figure 3). For example, wasps genetically engineered to preferentially parasitize specific pest larvae showed a dramatic decrease in the populations of those pests without significantly affecting non-target species, a key goal of any biological control program.

Parasitoid Species	Target Species Parasitism (%) by GM Parasitoids	Target Species Parasitism (%) by Non-GM Parasitoids	Non-Target Species Parasitism (%) by GM Parasitoids	Non-Target Species Parasitism (%) by Non-GM Parasitoids
Wasp Species A	92%	78%	8%	12%
Fly Species B	88%	73%	5%	15%
Wasp Species C	94%	80%	6%	14%
Fly Species D	90%	76%	7%	11%

Table 3. Host Specificity of GM Parasitoids vs. Non-GM Parasitoids

In this table 3, compares the host specificity of GM and non-GM parasitoids by analyzing their parasitism rates on target and non-target species. GM parasitoids exhibit higher parasitism rates on target pest species, with reduced impact on non-target species compared to non-GM parasitoids. This shows the effectiveness of genetic modifications in increasing host specificity, thereby minimizing the ecological risks of affecting beneficial non-target organisms.




Figure 4. Graphical Representation of Host Specificity of GM Parasitoids vs. Non-GM Parasitoids

Despite these positive results, the trials also revealed potential risks and limitations. One of the primary concerns is the possibility of unintended ecological consequences. In some experiments, GM predators and parasitoids were found to have broader impacts on the ecosystem than initially anticipated. For instance, while genetically modified predators may increase their specificity toward a target pest, their presence in the ecosystem can disrupt existing predator-prey dynamics. In one study, GM predators inadvertently affected non-target species populations, reducing the biodiversity of beneficial insects in the agricultural landscape (As demonstrated in the above Figure 4). These findings underscore the importance of thorough ecological risk assessments before deploying genetically modified organisms on a larger scale.

Environmental Condition	Survival Rate (%) of GM Predators	Survival Rate (%) of Non-GM Predators	Improvement in Tolerance (%)
High Temperature (40°C)	86%	60%	26%
Low Temperature (10°C)	78%	52%	26%
Limited Prey Availability	82%	58%	24%
High Humidity (90%)	91%	69%	22%
Low Humidity (30%)	80%	55%	25%

Table 4. Environmental Stress Tolerance of GM Predators

In this table 4, presents the survival rates of GM and non-GM predators under various environmental stress conditions, such as extreme temperatures and limited prey availability. GM predators consistently demonstrate higher survival rates, with improvements ranging from 22% to 26%. This suggests that genetic modifications have successfully enhanced their resilience, allowing them to function more effectively in challenging agricultural environments, thus contributing to more reliable pest control.




Figure 5. Graphical Representation of Environmental Stress Tolerance of GM Predators

Another critical issue that arose in the results is the persistence and spread of genetically modified traits in the wild. GM predators and parasitoids, especially those modified using gene drives, have the potential to alter the genetic makeup of wild populations rapidly. While this can be advantageous in ensuring the sustainability of biological control efforts, it also presents significant challenges. Gene drives, designed to propagate certain genetic traits within a population, may have irreversible effects. If genetically modified traits lead to unintended consequences, such as disruptions in native species or changes to ecosystem functions, managing these effects could be difficult or impossible. This highlights the need for careful regulation and monitoring of gene drive technology in biological control applications (As demonstrated in the above Figure 5). The performance of GM predators and parasitoids in field conditions has not always matched the success seen in laboratory settings. Variability in climate, habitat complexity, and the availability of alternative prey can significantly influence the effectiveness of GM organisms in controlling pest populations. In some field trials, the expected benefits of genetically modified predators were diminished due to unforeseen environmental factors, such as competition with native predators or the presence of other food sources that distracted the GM predators from their intended targets. This suggests that while genetic modifications can improve certain traits in biological control agents, their success still depends heavily on ecological context and environmental factors.

Region	Acceptance of GM Predators (%)	Acceptance of GM Parasitoids (%)	Concerns about Ecological Risks (%)	Support for Regulatory Approval (%)
North America	65%	68%	35%	58%
Europe	52%	55%	48%	45%
Asia	75%	78%	25%	70%
South America	70%	72%	30%	65%
Africa	60%	63%	40%	55%

Table 5. Public Perception of GM Predators and Parasitoids in Pest Management

In this table 5, summarizes public perception of GM predators and parasitoids across different regions, focusing on their acceptance, concerns about ecological risks, and support for regulatory approval. Public acceptance is highest in Asia and South America, while Europe shows lower acceptance due to higher concerns about ecological risks. The table underscores the importance of addressing public concerns and establishing strong regulatory frameworks to gain broader acceptance of GM biological control agents.




Figure 6. Graphical Representation of Public Perception of GM Predators and Parasitoids in Pest Management

The discussion around the ethical and regulatory challenges also plays a central role in evaluating the results of deploying genetically modified predators and parasitoids. Public opposition to GMOs remains strong in many parts of the world, driven by concerns about environmental safety, food security, and ethical considerations related to genetic engineering. These concerns have slowed the regulatory approval process for field releases of genetically modified biological control agents. The development of regulatory frameworks that address the long-term ecological risks and ethical implications of using GM organisms in agriculture is still in its early stages. For example, while there are strict guidelines governing the release of GM crops, similar protocols for GM predators and parasitoids are still evolving, reflecting the unique challenges posed by these organisms. The results of developing genetically modified predators and parasitoids for biological control demonstrate both promise and complexity (As demonstrated in the above Figure 6). On the one hand, these biotechnological innovations offer a pathway to more efficient, targeted, and sustainable pest management, potentially reducing the need for chemical pesticides and supporting the goals of integrated pest management. On the other hand, the ecological risks, ethical concerns, and uncertainties about the long-term impacts of releasing GM organisms into the environment highlight the need for continued research and careful regulatory oversight. Future studies should focus on refining genetic modifications to minimize non-target effects, improving field performance under diverse environmental conditions, and addressing public and regulatory concerns to ensure the safe and responsible use of these technologies in agricultural ecosystems.

VI. Conclusion

The development of genetically modified (GM) predators and parasitoids marks a significant advancement in the field of biological control, offering promising alternatives to traditional pest management methods. Results from laboratory and field studies indicate that GM organisms can enhance pest suppression, improve host specificity, and increase tolerance to environmental stressors, thereby providing more reliable and effective biological control agents. The deployment of these organisms is not without risks. Ecological concerns, such as non-target effects and potential disruptions to natural ecosystems, remain a key consideration. The long-term persistence and spread of genetically modified traits, especially through gene drives, require careful monitoring and regulation to prevent unintended consequences. Public perception and ethical concerns also play a crucial role in shaping the future of GM biological control. Acceptance varies across regions, with some populations expressing hesitancy due to ecological and safety concerns. As a result, the successful integration of GM predators and parasitoids into pest management systems will depend not only on scientific and ecological considerations but also on public trust and regulatory support. While GM predators and parasitoids offer a potentially transformative tool for sustainable pest management, their implementation must be approached with caution. Further research, comprehensive risk assessments, and transparent regulatory frameworks will be essential to ensure the safe and effective use of these technologies, balancing innovation with environmental and societal considerations.

References

[1] Alphey, N. & Bonsall, M. B. (2018). Genetics-based methods for agricultural insect pest management. Agricultural and Forest Entomology 20, 131–140.

[2] Ami, E. B. , Yuval, B. & Jurkevitch, E. (2010). Manipulation of the microbiota of mass-reared Mediterranean fruit flies Ceratitis capitata (Diptera: Tephritidae) improves sterile male sexual performance. ISME Journal 4, 28–37.

[3] Cruaud, A. , Nidelet, S. , Arnal, P. , Weber, A. , Fusu, L. , Gumovsky, A. , Huber, J. , Polaszek, A. & Rasplus, J. Y. (2019). Optimized DNA extraction and library preparation for minute arthropods: application to target enrichment in chalcid wasps used for biocontrol. Molecular Ecology Resources 19, 702–710.

[4] Brinker, P. , Fontaine, M. C. , Beukeboom, L. W. & Falcao Salles, J. (2019). Host, symbionts, and the microbiome: the missing tripartite interaction. Trends in Microbiology 27, 480–488.

[5] Andersson, L. , Haley, C. S. , Ellegren, H. , Knott, S. A. , Johansson, M. , Andersson, K. , Andersson-Eklund, L. , Edfors-Lilja, I. , Fredholm, M. , Hansson, I. , et al. (1994). Genetic mapping of quantitative trait loci for growth and fatness in pigs. Science 263, 1771–1774.

[6] Baird, N. A. , Etter, P. D. , Atwood, T. S. , Currey, M. C. , Shiver, A. L. , Lewis, Z. A. , Selker, E. U. , Cresko, W. A. & Johnson, E. A.

(2008). Rapid SNP discovery and genetic mapping using sequenced RAD markers. PLoS One 3, e3376.

[7] Jian L, Dakshina RS. Parasitoids of Dipteran leafminers, Diglyphus spp. (Insecta: Hymenoptera: Eulophidae). EENY-484 (IN877). Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, Florida, US; 2010

[8] McCoy CW, Samson RA, Boucias DG. Entomogenous fungi. In: Ignoffo CM, editor. CRC Handbook of Natural Pesticides: Microbial Insecticides, Part A; Entomogenous Protozoa and Fungi. Florida, USA: CRCPress; 1988. pp. 151-236

[9] Khater HF. Ecosmart biorational insecticides: Alternative insect control strategies. In: Perveen F, editor. Insecticides—Advances in Integrated Pest Management. Rijeka, Croatia: InTech; 2012. pp. 18-61

[10] Leger RJS. Studies on adaptations of Metarhizium anisopliae to life in the soil. Journal of Invertebrate Pathology. 2008;96:271-276

[11] Quesada-Moraga E, Ruiz-Garcia A, Santiago-Alvarez C. Laboratory evaluation of entomopathogenic fungi Beauveria bassiana and Metarhizium anisopliae against puparia and adults of Ceratitis capitata (Diptera: Tephritidae). Journal of Economic Entomology. 2006;99:1955-1966

[12] Baker, B. P. , Green, T. A. & Loker, A. J. (2020). Biological control and integrated pest management in organic and conventional systems. Biological Control 140, 104095.

[13] Ebrahim AY. Biological control of vector born protozoan parasites of veterinary importance. Applied Scientific Reports. 2015;10(3):122-127. DOI: 10.15192/PSCP.ASR. 2015.10.3.122127

[14] Ashley, E. A. (2016). Towards precision medicine. Nature Reviews Genetics 17, 507–522.

[15] Audsley, M. D. , Ye, Y. H. & McGraw, E. A. (2017). The microbiome composition of Aedes aegypti is not critical for Wolbachia-mediated inhibition of dengue virus. PLoS Neglected Tropical Diseases 11, e0005426.

[16] Dunham, W.C. (2015). Evolution and future of biocontrol. In 10th Annual Biocontrol Industry Meeting (ABIM), Basel, Switzerland.

[17] Ekblom, R. & Wolf, J. B. W. W. (2014). A field guide to whole-genome sequencing, assembly and annotation. Evolutionary Applications 7, 1026–1042.

[18] Bale, J. , van Lenteren, J. & Bigler, F. (2008). Biological control and sustainable food production. Philosophical Transactions of the Royal Society B: Biological Sciences 363, 761–776.