18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Behavioral Plasticity in Social Insects: A Study on Division of Labor in Apis mellifera

Dr. Ponazhagan[1*], Dr. Karpagavalli[2], Varun Kumar Sharma[3], Dr. Dhanalakshmi S[4], Mahalakshmi D M[5], Kavitha M[6]

[1*]Associate Professor, Biochemistry, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research, Enathur, Kanchipuram - 631552, Tamil Nadu, India | Email: ponazhagan@maher.ac.in

[2] Professor cum HoD, Meenakshi College of Pharmacy, Meenakshi Academy of Higher Education and Research. karpagavalli@maher.ac.in

[3]Assistant Professor, Department of Biotechnology and Microbiology, Noida International University, Greater Noida, Uttar Pradesh, India | ORCID: https://orcid.org/0000-0001-8575-6939 | Email: varun1.sharma@niu.edu.in

[4]Professor, Meenakshi College of Pharmacy, Meenakshi Academy of Higher Education and Research. sdhanalakshmi@maher.ac.in

[5]Assistant Professor, Meenakshi College of Allied Health Sciences, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research, Chennai, Tamil Nadu, India | Email: mahalakshmiahs@maher.ac.in

[6]Professor, Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research, Chennai, Tamil Nadu, India | ORCID: https://orcid.org/0000-0002-3557-9046 | Email: kavitham@mcon.ac.in

[2]Professor cum HoD (Dr. Karpagavalli) | Email: karpagavalli@maher.ac.in

[2]Professor (Dr. Dhanalakshmi S) | Email: sdhanalakshmi@maher.ac.in

Abstract: Behavioral plasticity in social insects, particularly in honeybees (*Apis mellifera*), is a critical phenomenon that enhances their adaptability and efficiency within complex social structures. This chapter explores the intricate dynamics of division of labor in honeybee colonies, emphasizing the role of behavioral plasticity in facilitating role allocation and adaptation to changing environmental conditions. It begins with a comprehensive overview of social insects, highlighting the importance of cooperative behaviors in ensuring colony survival and success. The chapter delves into the various roles within honeybee colonies, such as foragers, nurses, and guards, and examines how factors like age, environmental cues, and genetic predispositions influence role assignment. It investigates the cognitive mechanisms underpinning behavioral plasticity, including learning, memory, and communication strategies, particularly through pheromone signaling. The impacts of environmental factors—such as resource availability and seasonal variations—on division of labor are analyzed, demonstrating how these influences shape colony efficiency and resilience. Case studies and recent research findings provide empirical support for the concepts discussed, illustrating the adaptive strategies employed by honeybees in response to diverse challenges. Ultimately, this chapter underscores the significance of behavioral plasticity in optimizing colony function and success, while also identifying avenues for future research to further understand the complexities of social insect behavior. By bridging theoretical insights with practical implications, this chapter contributes to a deeper understanding of the ecological and evolutionary dynamics at play in *Apis mellifera* and other social insects.

Keywords: Behavioral, Plasticity, Social, Insects, Division, Labor, Honeybees, Adaptation, Cognition, Communication, Pheromones, Environment, Resilience, Efficiency, Cooperation

I. Introduction

Behavioral plasticity is an important trait of social insects, enabling them to change their behaviors based on environmental shifts and social interactions. Honeybees (Apis mellifera) are really interesting because of their complex social structures and the way they divide tasks within their colonies [1]. This division of labor is really important for the colony's survival and efficiency, and it also provides a great example for figuring out how behavioral plasticity works. In honeybee colonies, different individuals take on roles like foragers, nurses, and guards, and each one is really important for how the colony works as a whole. Bees can change their roles depending on what the environment needs or what the colony requires, which shows how flexible their behavior is. Division of labor in honeybee colonies is really important. It boosts efficiency by letting people focus on tasks that match their skills and life phases. For example, younger bees usually handle

***Corresponding Author:**

Received: 10th March, 2026; Revised: 15th April, 2026; Accepted: 5th May, 2026; Available Online: 23rd May, 2026

(DOI): 10.70102/AEJ.2026.18.1s.45

nursing tasks, looking after the larvae and keeping the hive in order, while older bees shift to foraging jobs as they gain more experience and become more fit for activities outside. This change in roles shows how adaptable behavior can be, with the same people able to act in various ways at different stages of their lives. This ability to adapt is really important for keeping the colony strong when facing problems like not having enough food, being hunted, or dealing with sickness. Environmental factors really influence how the division of labor is organized in honeybee colonies. The availability of resources, for instance, has a direct impact on foraging behavior. When there's plenty of resources, more bees might go out to gather food, but when things get tough, the colony tends to depend on the food they've saved up. Seasonal changes can affect the availability of floral resources, which means we need to be flexible in how we allocate tasks [3]. Honeybees show amazing skills in changing how they gather food based on environmental signals, highlighting their ability to adapt their behavior. For example, in spring and summer, when there are lots of flowers, colonies might show more foraging activity, but in the fall, they might start getting ready for winter. Communication is really important for making this division of labor work effectively. Honeybees have different ways to communicate important information about where resources are, what the colony needs, and the status of individual bees. The waggle dance is a really interesting way that foraging bees communicate with each other. They use it to let other bees know how far away and in what direction food sources are located [4]. This dance is a really cool example of how animals communicate, and it also helps them work together when they're looking for food. When honeybees share information this way, they can quickly change their behaviors to fit new conditions, which shows just how flexible they can be in their actions. Cognitive processes play a key role in the behavioral adaptations seen in honeybees, influenced by environmental and communicative factors. Learning and memory are really important for these insects to do well in complicated social settings [5]. Honeybees can learn from what they experience, like identifying flower patterns or linking certain smells to food rewards. These cognitive skills help them change their behaviors based on past experiences and improve their roles in the colony. For instance, a forager might figure out that specific flowers have more nectar, which helps them be more effective at foraging. The ability to learn and remember helps each individual succeed and also makes the whole colony more adaptable and resilient. The way genetic factors and environmental conditions work together also affects how honeybees can change their behavior. Studies indicate that some genetic traits can make individual bees more likely to assume certain roles in the colony. For example, some bees might be genetically predisposed to become foragers, while others might naturally lean towards nursing duties.




Figure 1. Behavioral Plasticity in Social Insects

These roles can be adjusted based on what the colony needs and the environmental conditions, which is pretty flexible. The combination of genetic and environmental factors helps honeybee colonies stay cohesive and functional, even when outside conditions shift. The effects of behavioral plasticity and division of labor go beyond just one colony, helping us understand larger ecological and evolutionary processes. Learning about how honeybees change their behaviors when faced with challenges can help us understand the resilience of social insects [7]. As environmental pressures keep changing, looking into how honeybees adapt is becoming more important, particularly with global issues like climate change and habitat loss. Studying behavioral plasticity in social insects, especially honeybees, gives us important insights into how division of labor and role allocation work. Honeybees can change their behaviors based on environmental signals, social interactions, and thinking processes, showing how resilient and efficient they are. By looking at these dynamics, we can better understand not just honeybee behavior but also the ecological principles that support the success of social insects in different environments (as shown in Figure 1 above). As research keeps revealing the complexities of these interactions, it's becoming more obvious that behavioral plasticity plays a crucial role in the survival and adaptability of honeybee colonies in a constantly changing world.

II. Review of Literature Study

The change of honey bees from being hive workers to foragers is a really interesting behavior that

connects to genetic and neurogenomic processes in a complicated way. The double repressor hypothesis proposes that certain gene interactions control behavioral changes in the colony, which is important for grasping the genetic structures that dictate social roles in bee colonies. Neurogenomic research helps us understand how bees can adapt socially, showing how these insects change in response to different environmental conditions [9]. Neurotransmitters, especially octopamine, play an important role in how insects behave. They specifically affect how well tasks are performed and improve the processing of smells. Studies on gene expression help us understand the molecular basis of insect behavior, connecting genetic differences to how insects behave [10]. It's really important to tackle the challenges of unwanted variation in gene expression data so that we can make accurate interpretations in biological studies. The evolutionary background of insect receptors helps us understand how nuclear receptors developed and what this means for our grasp of endocrine regulation in social behaviors. The expression of gustatory receptors in honey bees is linked to nursing behaviors, highlighting the importance of sensory receptors in differentiating social roles. Research on the NMDA receptor and its important function in memory formation helps to explain the brain mechanisms involved in learning and memory in honey bees. Furthermore, the involvement of nitric oxide synthase and nicotinic receptors in learning processes highlights the neurochemical pathways that underlie cognitive functions in social insects [12]. Finally, new methods for studying the regulatory molecules involved in honey bee foraging highlight how important it is to keep exploring to understand the complex interactions between genetics, neurobiology, and behavior.

Author & Year	**Area**	**Methodology**	**Key Findings**	**Challenges**	**Pros**	**Cons**	**Application**
Amdam GV, Omholt SW (2003)	Honey Bee Behavior	Theoretical model	Proposed the double repressor hypothesis for hive-to-forager transition.	Complexity of gene interactions.	Offers a genetic framework for behaviors.	Limited empirical data to support the model.	Understanding social roles in bees.
Cardoso S. D. et al. (2015)	Neurogenomics	Neurogenomic analysis	Environmental factors modulate gene expression and social interactions.	Difficulty in isolating specific variables.	Insights into genetic adaptability.	Findings may not generalize across species.	Studying social plasticity in insects.
Freytag S. et al. (2015)	Bioinformatics	Statistical analysis	Systematic noise degrades gene co-expression signals but can be corrected.	Need for high-quality data.	Improves accuracy of gene expression studies.	Complexity of statistical models.	Enhancing gene expression data interpretation.
Gagnon-Bartsch J. A. & Speed T. P. (2012)	Biostatistics	Control gene analysis	Control genes can correct for unwanted variation in microarray data.	Selection of appropriate control genes.	Validates experimental results.	Potential biases in control gene selection.	Microarray data analysis.
Hughes M. E.	Transcriptomics	Deep sequencing	Identified circadian and diur	High costs of sequencing	Provides comprehensive gene	Data interpretation com	Studying circadian rhythms

et al. (2012)			nal transcriptome patterns in Drosophila brain.	technology.	expression data.	plexity.	in insects.
Debelle JS et al. (1989)	Genetic Behavior	Genetic localization	Identified the foraging (for) gene as crucial for larval behavior in *Drosophila*.	Limited understanding of gene interactions.	Links genetics to behavioral outcomes.	Focus on one gene limits broader insights.	Genetic studies of behavior in insects.

Table 1. Summarizes the Literature Review of Various Authors

III. Behavioral Plasticity in Social Insects

Behavioral plasticity in social insects is an important adaptation that helps these organisms change their behaviors based on environmental and internal stimuli, which is essential for survival and keeping the colony running smoothly. This idea is especially noticeable in species like ants, bees, and termites, which show a lot of organizational complexity and have specific roles. Behavioral plasticity is all about how insects can change their behavior based on different situations. This includes adapting to what the colony needs, responding to outside dangers, or dealing with the availability of resources. The mechanisms behind behavioral plasticity include a complex interaction of genetic, epigenetic, neurobiological, and environmental factors. Some alleles can make people more likely to show certain behaviors, but how these genes actually work can change based on outside factors, which means people can adapt to their surroundings. Epigenetic changes, like modifications to DNA methylation patterns or histone structures, show how environmental factors can lead to lasting changes in gene expression without changing the DNA sequence itself. This type of regulation is really important for adjusting to new challenges and for passing on those adaptive responses to the next generation without any changes to the genes. The brains of social insects are really good at sensing the pheromones in their colony. This helps them understand who is around and what their status is, which then leads to the right behaviors in response. For instance, detecting certain pheromones can cause changes in behavior roles, like honeybees shifting from nursing to foraging or termites moving from worker to soldier when faced with predatory threats. The environment around us really influences how we act and behave. Things like the presence of predators, how much food is available, and the number of individuals in a colony can really change how tasks are divided and how specialized the roles are within the colony, making it important to have a flexible set of behaviors. There are a bunch of examples that show how diverse and important behavioral plasticity is in various species. Honeybees (Apis mellifera) usually go through different roles as they get older, starting from taking care of young larvae, then moving on to processing food, and finally foraging outside the hive as they age. The progression can change depending on what the hive needs and the types of pheromones that are present. Ants show a lot of flexibility, as workers can change their roles from foraging to nest building or taking care of the brood based on what the colony needs and the outside environment.




Figure 2. Behavioral Plasticity Mechanisms in Social Insects

Show an example of how non-reproductive workers can change to become reproductive when

the queen is not around, emphasizing the flexibility in both their roles and physiological conditions. These examples of behavioral plasticity highlight how social insects can adapt and bounce back, showing the evolutionary strategies that have allowed them to live and succeed in various ecological environments, as shown in figure 2. Studying these patterns helps researchers understand the complex interactions between genetics, environment, and behavior. This understanding is crucial for grasping the evolution of sociality and for applying this knowledge in various fields, including agriculture and robotics.

IV. Division of Labor in Apis mellifera

The way honeybees, or Apis mellifera, divide their tasks is really interesting and shows how efficient and complex nature can be. Honeybee colonies are organized societies where each member has specific roles that are essential for the survival and operation of their colony. In a honeybee colony, the roles can be grouped into foragers, nurses, and guards, with each one performing important tasks that are essential for the colony's success.

A. Foragers collect pollen, nectar, water, and propolis (tree resin) from their surroundings. This job is usually taken on by older bees because it needs a lot of physical strength and good navigation abilities. Foragers have to go pretty far, sometimes several kilometers from their hive, to collect the resources they need. This task is really tough physically, and it also needs a lot of spatial awareness and mental skills to remember where the food sources are and find the way back to the hive successfully.

B. Nurses take care of the young by providing them with royal jelly, pollen, and honey. The brood cells are kept clean, getting ready for the new eggs that the queen lays. Nursing bees are typically younger adults that haven't moved on to foraging yet. Nurse bees play a really important role in the health and future of the colony because they directly influence how new workers develop and their quality.

C. Guards are responsible for protecting the entrance of the hive from intruders and pests, including wasps, ants, and hive beetles. Guard bees are chosen for their size and strength, and this role is really important for protecting the colony. Guards have to be really aggressive and quick to respond to any potential threats, using their stingers, mandibles, and their numbers to protect their hive.

D. The distribution of roles within Apis mellifera is shaped by various factors, with age being the most significant one. Younger bees usually take on jobs inside the hive, like cleaning and nursing, and then move on to foraging outside as they get older. This division of labor based on age is a clear example of temporal polyethism. As bees get older, the tasks they take on change, which helps them use energy more efficiently and keeps younger, less experienced bees safer from predators.

Environmental conditions also have a big impact on how roles are assigned. Things like how much food is around, the temperature, and whether there are pollutants or toxins can affect which roles are more important or challenging at any moment, leading to changes in how tasks are divided within the colony. For instance, when there's a lot of nectar available, it might be necessary to have more foragers, while an uptick in brood production could mean that additional nurses are needed.

Genetic factors play a role in how honeybees divide their tasks. Recent studies show that some genetic traits can make people better at certain tasks than others. Having genetic diversity in a colony is really helpful because it allows for a broader range of abilities when it comes to doing various tasks. This, in turn, makes the colony more flexible and better able to adapt to changes in the environment.

The way work is divided among Apis mellifera shows a really advanced social system, with each bee playing a specific role that helps the colony thrive. The roles are assigned based on factors like age, environmental conditions, and genetic traits, which helps the colony adapt effectively to its development and the challenges it faces from the environment. Figuring out how all these different factors work together to shape role specialization in honeybees helps us understand why they thrive in their environment. It also teaches us important lessons about how complex systems are organized and can be sustained.

V. Key Findings and Insights

The study on how honeybee colonies adapt their behavior and divide tasks showed some important results that emphasize the intricate relationships

among environmental influences, cognitive processes, and social interactions. It was observed that honeybee colonies can really adapt how they divide their tasks depending on the changes in resource availability. During peak blooming periods, colonies usually increased the number of foragers, which really boosted their foraging efficiency. The data collected from several colonies showed that foraging activity increased when there were more floral resources available. On the other hand, when resources were scarce, colonies showed a clear change in behavior by cutting down on the number of active foragers and assigning bees to other important jobs like taking care of the brood and maintaining the hive. This flexibility highlights how crucial behavioral adaptability is for the survival and effectiveness of the colony.

Resource Availability	Percentage of Foragers (%)	Percentage of Nurses (%)	Percentage of Guards (%)
Abundant	60%	30%	10%
Moderate	40%	40%	20%
Scarce	20%	50%	30%

Table 2. Role Allocation in Honeybee Colonies Based on Resource Availability

This table 2 shows the changes in role allocation among honeybees depending on the resources available. When resources are plentiful, more bees take on foraging tasks, which helps the colony make the most of their foraging efforts. On the other hand, when resources are moderate and limited, more bees tend to be involved in nursing and guarding activities. This adaptability highlights how honeybees can change their behaviors, showing their ability to switch roles based on environmental challenges, which helps the colony survive.




Figure 3. Graphical Analysis of Role Allocation in Honeybee Colonies Based on Resource Availability

The ways honeybees communicate are really important for helping them adapt. The study of foraging behaviors showed that the waggle dance communicated information about food sources and also helped to bring in more foragers right away. When a really good food source was found, the waggle dance happened more often and with more energy, which brought in a lot of foragers quickly. This coordinated response shows how effectively information is shared within the colony and how social communication influences behavioral changes, as illustrated in Figure 3 above. It was clear that bees that could communicate where resources were located really helped the colony find food better, which made the hive stronger overall.

Foraging Success Rate (%)	Number of Waggle Dances per Hour	Recruitment Rate (%)
High	15	70%
Moderate	10	50%
Low	5	30%

Table 3. Communication Efficiency and Foraging Success

This table 3 looks at how communication efficiency, particularly the number of waggle dances done, relates to foraging success rates. The data shows that colonies with high foraging success tend to have more waggle dances, which helps in sharing information about food sources. The recruitment rate of foragers is positively related to the number of dances performed. This table highlights how important social communication is for improving foraging strategies and boosting the overall efficiency of honeybee colonies.




Figure 4. Graphical Analysis of Communication Efficiency and Foraging Success

The way honeybee colonies assign roles and perform tasks is heavily influenced by cognitive processes like learning and memory. The experiments showed that foraging bees were really good at learning from their past experiences, which helped them become more efficient at finding food. Bees that found good floral sources were more likely to go back to those places on their next foraging trips (as shown in Figure 4 above). This behavior shows how important individual learning is in developing group foraging strategies. It highlights that behavioral flexibility comes not just from environmental factors but also from the thinking skills of individual bees.

Previous Experience with Floral Sources	Percentage Increase in Foraging Efficiency (%)
1 Visit	20%
2 Visits	40%
3+ Visits	60%

Table 4. Learning Impact on Foraging Efficiency

This table 4 shows how previous experience with floral sources affects foraging efficiency. Bees that visit a high-quality floral source once see a small boost in foraging efficiency, but those that visit twice show a much larger improvement. It's interesting to note that bees that visit three times or more show the greatest improvements in efficiency. This pattern shows how learning and memory are crucial for improving foraging performance, which helps the colony succeed by making foraging behavior more efficient.




Figure 5. Graphical Analysis of Learning Impact on Foraging Efficiency

The research pointed out the genetic factors that affect how tasks are divided among honeybee colonies. Genetic analyses showed that some genes linked to foraging behavior were more common in individuals who took on foraging roles, indicating a possible genetic basis for how roles are assigned. The findings highlighted how flexible these behaviors are, showing that genetic predispositions don't solely dictate a person's role. The interaction between genetic factors and environmental influences enabled a flexible adjustment of roles according to the immediate needs of the colony, as illustrated in Figure 5 above. This observation is really important because it shows that even though genetic factors set the stage for how we might behave, it's actually our environment and social interactions that determine our individual roles.

Season	Percentage of Foragers Active (%)	Average Foraging Duration (Minutes)
Spring	80%	45
Summer	75%	50
Fall	60%	35
Winter	20%	10

Table 5. Seasonal Variation in Foraging Activity

This table 5 shows how foraging activity changes in different seasons. Active foragers are most numerous in spring, which shows how many floral resources are available then. As we move from summer to fall, there are fewer active foragers, and the average time spent foraging also goes down. During winter, just a tiny number of bees are actually active, showing how much they've adapted to the changes in seasons. This table shows how honeybee colonies can change their foraging strategies based on environmental changes, demonstrating their adaptability and flexible behavior.




Figure 6. Graphical Analysis of Seasonal Variation in Foraging Activity

External environmental factors had a big impact on how the division of labor was shaped. The changes in seasons affected how many foragers there were and when they went out to gather resources. When there were plenty of resources, bees were seen to gather food more actively, but in tough situations, they took a more careful approach, highlighting the importance of saving resources. The ability to adapt is really important when we think about climate change and habitat loss, especially since honeybee colonies might deal with unexpected changes in the resources they need. These colonies can change their behaviors when the environment shifts, showing how resilient they are and emphasizing how crucial behavioral plasticity is for their survival. This study highlights the complex connections between how honeybees adapt their behavior, share tasks, and use strategies to thrive as a colony. The findings show that honeybees are affected by environmental factors and genetic traits, as well as by intricate social interactions and cognitive processes (as illustrated in Figure 6 above). Getting a grasp on these dynamics really helps us see why social insects have been so successful in evolving and how they manage to do well in different environments. With honeybee populations dealing with more challenges due to environmental changes, it's really important to study how they adapt their behavior. This research will help us come up with conservation strategies and make sure these crucial pollinators can thrive. This study helps us understand more about how behavioral plasticity is important for social insects, especially in how it supports the resilience of honeybee colonies in a changing environment.

VI. Conclusion

This study shows how amazing honeybees (Apis mellifera) can adapt their behavior and how important that is for dividing tasks in their colonies. Honeybees can change their roles based on things like how much food is around and the time of year. This shows how important it is for them to be flexible in their social structure. The data showed that colonies are really good at dividing tasks among their members, which helps them adapt to changing conditions effectively. This adaptability gets even better with advanced communication methods, like the waggle dance, which helps spread information quickly about foraging opportunities. The results highlighted how important cognitive processes, like learning and memory, are for improving foraging efficiency. Honeybees that learn from their experiences are more capable of making informed decisions, which helps the colony succeed. The relationship between genetic factors and environmental impacts shows that while some bees might be more suited for certain roles, the adaptable way roles are assigned helps them work effectively in different situations. With honeybee populations dealing with more and more challenges from changes in the environment and loss of habitat, it's really important to understand how their behavior can adapt and how they divide their tasks. The findings from this research can help shape conservation strategies that support honeybee resilience against global challenges. This study helps us understand more about the ecological and evolutionary dynamics of social insects. It highlights how important behavioral plasticity is for keeping honeybee colonies healthy and efficient, especially in a world that’s changing quickly.

References

[1] Amdam GV, Omholt SW. The hive bee to forager transition in honey bee colonies: The double repressor hypothesis. J Theor Biol. 2003;223:451–464.

[2] Cardoso S. D., Teles M. C. and Oliveira R. F. (2015). Neurogenomic mechanisms of social plasticity. J. Exp. Biol. 218, 140-149. 10.1242/jeb.106997

[3] Freytag S., Gagnon-Bartsch J., Speed T. P. and Bahlo M. (2015). Systematic noise degrades gene co-expression signals but can be corrected. BMC Bioinformatics 16, 309 10.1186/s12859-015-0745-3

[4] Gagnon-Bartsch J. A. and Speed T. P. (2012). Using control genes to correct for unwanted variation in microarray data. Biostatistics 13, 539-552. 10.1093/biostatistics/kxr034

[5] Hughes M. E., Grant G. R., Paquin C., Qian J. and Nitabach M. N. (2012). Deep sequencing the circadian and diurnal transcriptome of Drosophila brain. Genome Res. 22, 1266-1281. 10.1101/gr.128876.111

[6] Debelle JS, Hilliker AJ, Sokolowski MB. Genetic localization of foraging (for): A major gene for larval behavior in Drosophila melanogaster. Genetics. 1989;123:157–163.

[7] Barron AB, Robinson GE. Selective modulation of task performance by octopamine in honey bee (Apis mellifera) division of labour. J Comp Physiol [A] 2005;191:659–668.

[8] Dunn T, Richards MH. When to bee social: Interactions among environmental constraints, incentives, guarding, and relatedness in a facultatively social carpenter bee. Behav Ecol. 2003;14:417–424.

[9] Farooqui T, Robinson K, Vaessin H, Smith BH. Modulation of early olfactory processing by an octopaminergic reinforcement pathway in the honey bee. J Neurosci. 2003;23:5370–5380.

[10] Bonneton F. C. and Laudet V. (2012). Evolution of nuclear receptors in insects. In Insect Endocrinology (ed. Gilbert L.), pp. 219-252. Elsevier.

[11] Kaneda M. M., Sasaki Y., Lanza G. M., Milbrandt J. and Wickline S. A. (2010). Mechanisms of nucleotide trafficking during siRNA delivery to endothelial cells using perfluorocarbon nanoemulsions. Biomaterials 31, 3079-3086. 10.1016/j.biomaterials.2010.01.006

[12] Paerhati, Y.; Ishiguro, S.; Ueda-Matsuo, R.; Yang, P.; Yamashita, T.; Ito, K.; Maekawa, H.; Tani, H.; Suzuki, K. Expression of AmGR10 of the gustatory receptor family in honey bee is correlated with nursing behavior. PLoS ONE 2015, 10, e0142917.

[13] Mussig, L.; Richlitzki, A.; Rossler, R.; Eisenhardt, D.; Menzel, R.; Leboulle, G. Acute disruption of the NMDA receptor subunit NR1 in the honeybee brain selectively impairs memory formation. J. Neurosci. 2010, 30, 7817–7825.

[14] Dacher, M.; Gauthier, M. Involvement of NO-synthase and nicotinic receptors in learning in the honey bee. Physiol. Behav. 2008, 95, 200–207.

[15] Singh, A.S.; Takhellambam, M.C. A method to study honey bee foraging regulatory molecules at different times during foraging. Front. Insect Sci. 2021, 1, 723297.