18(1s), Spring 2026

Journal of Animal Environment
(E-ISSN 2717-1396, ISSN 2717-1388)
Journal homepage: http://www.aejournal.ir

Original Research Paper

Integrating Wildlife Conservation Objectives into Modern Agricultural Production Landscapes

Dr. Dilip Hinge[1], Pratiksha Singh[2], Dr. Nallusamy Duraisamy[3], Suraj Luthra[4], Amandeep Kaur Gill[5], Elnoza Kudratova[6]

[1]Assistant Professor, Krishna Institute of Science and Technology, Krishna Vishwa Vidyapeeth "Deemed to be University", Taluka-Karad, Dist-Satara, Pin-415 539, Maharashtra, India (Dr. Dilip Hinge) | Email: ddhinge@gmail.com

[2]Assitant professor, Department of Agriculture, Noida International University, Noida, Uttar Pradesh, India. (Pratiksha Singh) | ORCID: https://orcid.org/0000-0002-2886-7234 | Email: pratiksha.singh@niu.edu.in

[3]Scientist, Department of Research, Meenakshi Academy of Higher Education and Research, Tamilnadu, India (Dr. Nallusamy Duraisamy) | Email: nallu@maher.ac.in

[4]Asst Prof, Agriculture Departmernt, Dev Bhoomi Uttarakhand University, Dehradun (Suraj Luthra) | ORCID: https://orcid.org/0009-0009-1624-2676 | Email: soa.suraj@dbuu.ac.in

[5]School of Pharmaceutical Sciences, CT University, Punjab, 142024, India (Amandeep Kaur Gill)

[6]Senior Lecturer, Department of Music Education, Faculty of Preschool and Primary Education, Jizzakh State Pedagogical University, Jizzakh, Uzbekistan (Elnoza Kudratova) | ORCIDhttps://orcid.org/0000-0003-4774-6371 | Email: elnozaismatilllayevna@gmail.com

Abstract: Increased food production due to modern agricultural intensification is one of the most significant impacts that has occurred, but habitat fragmentation, loss of biodiversity and degradation of ecosystems are also a huge problem for wildlife conservation. Incorporating wildlife conservation goals into agriculture production landscapes has thus become a viable strategy for supporting food security while promoting ecological resilience. The review explores the dynamics between agriculture and wildlife, focusing on the consequences of intensive farming, habitat changes and human-wildlife conflicts for wildlife ecology. It addresses the principles of wildlife-friendly agriculture such as restoration of habitats, development of biodiversity-friendly farm systems, establishing ecological corridors, and creating landscape connectivity. In addition, the review identifies some sustainable agriculture trends and new technologies being developed for better biodiversity assessment and conservation planning, including remote sensing, geographic information systems, artificial intelligence, Internet of Things (IoT) based monitoring, and digital decision-support platforms. Other scientific tools for conservation prioritization in agricultural landscapes, such as biodiversity indicators, species distribution modelling and habitat suitability analysis, are also explored. The synthesis shows that the combination of ecological knowledge, precision agriculture and advanced monitoring technologies has a positive effect on the provision of ecosystem services, landscape resilience and long-term agricultural sustainability. The review offers exhaustive insight into current strategies, challenges to implementation and future research directions to create conservation oriented agricultural systems, which at the same time support productive agriculture while preserving and strengthening biodiversity and environmental sustainability in a changing global context.

Keywords: Wildlife conservation; Sustainable agriculture; Biodiversity conservation; Agricultural landscapes; Precision agriculture; Ecosystem services.

I. Introduction

The agriculture sector is one of the major agricultural land use systems in the world and a key determinant of food security, economic development and rural livelihoods. In the last few decades, agricultural production has increased significantly, driven by the growing world population and shifting food consumption habits and a rapid rate of urbanization. Despite the considerable advances in agricultural production and livestock productivity in modern times, there has been significant environmental problems like habitat loss, biodiversity loss, land degradation, water pollution and fragmentation of the ecosystems. Such environmental effects have been of great concern for the sustainability of agricultural production systems and their suitability with the global conservation targets on biodiversity. As a result, incorporating wildlife conservation goals into agriculture land use

***Corresponding Author:**
Received: 24th April, 2026; Revised: 11th May, 2026; Accepted: 24th May, 2026; Available Online: 6th June, 2026
(DOI): 10.70102/AEJ.2026.18.1s.57

has gained greater significance as a component of sustainable agricultural development, while maintaining ecosystem integrity. Wildlife play a critical role in healthy ecosystems, offering many ecosystem functions and ecosystem services, both directly and indirectly, which are vital to agricultural productivity [1]. Pollinating insects boost crop production, predators control agricultural pests, soil organisms help cycle nutrients, and diverse vegetation helps control erosion, sequester carbon and help regulate water. In addition, wildlife plays a key role in providing ecosystem balance by supporting food webs, increasing the resilience of ecosystems, and providing genetic diversity within landscapes. Intensive farming has led to significant declines in habitat availability and ecological interactions, such as the loss of natural vegetation, conversion of wetlands, overgrazing, too high levels of mechanized farming and chemical fertilizers and pesticides. Consequently, the population of numerous wildlife species has been reduced, their migration patterns changed, and they have become more susceptible to environmental changes [2]. Conservation strategies have been mostly based on the creation of protected areas (national parks, wildlife sanctuaries and nature reserves). While these protected areas continue to be essential for the conservation of threatened species, they are not enough to ensure the protection of biodiversity as a significant percentage of wildlife is found in areas outside formally protected areas. Much of land use is now agricultural, and often plays a significant role as habitat, migration and buffer zones for a variety of species. So conservation efforts need to go beyond areas of protection, and incorporate biodiversity into agricultural planning and management. Wildlife-friendly agriculture does not mean that agriculture and biodiversity conservation are mutually exclusive, but that they can be achieved together by implementing suitable land use and planning for ecology [3].

Conservation-oriented agriculture is a concept that focuses on the development of agricultural systems which remain productive without compromising the ecological functions. HABITAT QUALITY & LANDSCAPE HETEROGENEITY: Sustainable agriculture methods including agroforestry, conservation agriculture, IPM, crop diversification, rotational grazing, organic farming, setting up riparian buffers, planting hedgerows and restoring habitats make a significant contribution. These strategies can benefit wildlife by offering them feeding sites, nesting areas, protection and buffers from predators, and ecological connections that allow for movement among fragments. Furthermore, diversified agriculture can have a beneficial influence on soil fertility, water use efficiency, climate change adaptation, ecosystem resilience, and stability, which can yield co-benefits for both food production and nature conservation. In recent years, technologies have also gained in their ability to reinforce the incorporation of wildlife conservation into mainstream farming practices. Continued monitoring of habitat conditions, wildlife populations, vegetation dynamics, and land-use changes can be achieved through the use of remote sensing technologies, Geographic Information Systems (GIS), unmanned aerial vehicles (UAVs), artificial intelligence (AI), machine learning algorithms, Internet of Things (IoT)-based environmental sensors, and cloud-based decision support platforms [4]. These digital technologies can be used for species distribution modelling, habitat suitability analysis, biodiversity assessment and precision conservation planning. Moreover, AI for image recognition and sensor connectivity offers valuable tools for wildlife detection, conflict resolution, and ecological monitoring, ensuring that farmers and conservationists can make informed decisions without harming the environment.

Although there is increasing awareness of agriculture's role as a protector of wildlife, there are many obstacles to implementing this new agriculture. Factors such as economic constraints, low ownership of the land, lack of incentives to conserve biodiversity, lack of policy integration, limited technical knowledge on the part of farmers and conflicting requirements for expansion of agriculture can make the adoption of biodiversity-friendly practices difficult for them. Conservation in agricultural landscapes is further complicated by climate change, invasive species, emerging diseases and growing anthropogenic pressure. These challenges are complex and can only be addressed with multidisciplinary efforts that bring together agricultural scientists, the ecology community, policy makers, conservation agencies, technology developers, and farming communities to develop integrated agricultural management strategies that promote the three core pillars of agricultural productivity, ecological resilience and biodiversity conservation [5]. This review investigates how wildlife conservation goals can be incorporated into today's agricultural production landscapes by addressing the relationships between agricultural and wildlife ecosystems, principles of wildlife friendly agriculture, conservation technologies for sustainable agriculture, advanced conservation technologies, ecological assessment techniques, and conservation planning processes. It also addresses contemporary issues, opportunities and future research directions for creating resilient agricultural landscapes that find a balance between food production, environmental health, biodiversity, and long term sustainability in the environment.

II. Interactions Between Agriculture and Wildlife Ecosystems

Agricultural activities have a strong influence on wildlife ecosystems, for example, by altering habitats, altering biodiversity and interacting with ecosystems services. Intense farming systems tend to degrade habitat quality, and sustainable farming systems can provide habitat for a variety of species. Balanced management of agriculture–wildlife relationships and ecological trade-offs and human–wildlife conflicts is key to creating strategies that will sustain the production of food and biodiversity.

A. Impacts of intensive farming practices on wildlife habitats

Intensive farming has dramatically altered natural ecosystems in terms of extensive land use, mechanisation, single crop farming, high use of fertilisers, insecticides, irrigation and livestock intensification. These activities will decrease habitat diversity, break up ecological areas, and destroy breeding grounds, feeding and nesting areas that are important to a large number of wildlife species. Agrochemical use is widespread and pollutes soil and water resources, which have a negative impact on pollinators, amphibians, aquatic organisms, birds and beneficial insects and on the ecological food webs [6]. Further loss of hedgerows, wetlands and native vegetation also limit movement of wildlife and reduce the flow of genes between isolated populations. Routine cultivation and overgrazing lead to soil erosion, an decrease in vegetation diversity and the loss of environmental resilience to disturbance. Greenhouse gas emissions from agriculture add to the changes in habitat suitability and species distribution due to climate change. The intensive farming practices have positive effects on the productivity of agriculture, but also have significant negative effects on biodiversity, which, for example, results in ecosystem degradation [7]. So sustainable land use, less dependency on chemicals, restoration of habitats, ecological buffer zones, and integrated farming systems are crucial to reduce habitat loss while ensuring productivity and sustainability in the long-term.

B. Role of Agricultural Landscapes in Supporting Biodiversity

Agricultural landscapes have the potential to support biodiversity if managed in an environmentally sustainable way that ensures that they remain ecologically complex and heterogeneous. Diversified agricultural landscapes that include agroforestry systems, hedgerows, grasslands, wetlands, riparian buffers and cover crops and mixed farming systems can create appropriate habitats for birds, mammals, pollinators, beneficial insects, reptiles and amphibians. These landscape features provide food, nesting areas, protection and flight trails which increase species survival and ecosystem resiliency [8]. Agricultural biodiversity plays a significant role in many ecosystem services such as natural pest control, pollination, nutrient cycling, soil fertility improvement, carbon sequestration and protection of water quality. Agriculture and natural habitats can be integrated in a way that makes them more connected to support wildlife, gene flow and adaptation to environmental shifts. In addition, a high diversity of agricultural systems is typically more resilient to climate variations, pest attacks, and diseases and less reliant on synthetic inputs into agriculture. In this way, promoting practices to conserve farming landscapes benefits the productivity of agricultural production and the conservation of ecosystems – agricultural landscapes can serve at the same time as productive food systems and valuable habitats for a wide range of wildlife communities [9].

C. Human–Wildlife Conflicts and Ecosystem Service Trade-Offs

Agricultural activities are encroaching on wildlife habitats, and wildlife have been taking to farmland for food and shelter, leading to a rise in human–wildlife conflict.Human–wildlife conflicts have been growing because the agriculture is encroaching on wildlife habitats and wildlife are increasingly going to agricultural land for food and shelter. Herbivores such as crop raiders and carnivores such as livestock also destroy crops, causing economic losses to farmers; zoonotic diseases are also often transmitted and water resources are often shared, causing losses of water for farming. Responding in turn with habitat destruction, fencing, poisoning, and hunting, however, can impact wildlife populations and biodiversity in even more ways. At the same time, wildlife also delivers many ecosystem services that are crucial to sustainable agriculture, such as pollination, bio-control, seed dispersal, nutrient recycling, balancing ecosystems etc. Therefore, the trade-offs in ecosystem services between agriculture and wildlife conservation need to be carefully considered. Even though conflicts are inevitable, they can be significantly minimized through conservation measures such as habitat restoration, ecological corridors, wildlife friendly fencing, compensation, community engagement, early warning systems and landscape level planning [11]. A summary of the possible agriculture–wildlife interactions, strategies for the conservation of biodiversity, and sustainable management of such relationships is found in Table 1. Scientific monitoring, adaptive management, and a mutually beneficial approach with stakeholders can help agricultural landscapes deliver both productive and sustainable agriculture and wildlife conservation, contributing to resilient environments and sustainable rural development in the face of shifts in environmental and climatic conditions

Study Area	Agricultural System	Wildlife/Ecosystem Focus	Key Findings	Limitation	Conservation Strategy
Europe [12]	Mixed farming	Animal welfare & biodiversity	Sustainable farming improves ecosystem health	Limited field validation	Animal welfare integration
Global [13]	Agricultural landscapes	Nature-based solutions	Nature-based practices improve ecosystem resilience	Regional variability	Green infrastructure
Africa [14]	Nature-positive agriculture	Biodiversity conservation	Diversified farming supports resilience	Economic adoption barriers	Agroecology
Portugal	Viticulture	Biodiversity enhancement	Habitat management increases beneficial species	Crop-specific	Habitat diversification
Global [15]	Agroecology	Urban biodiversity	Agroecological farming enhances ecosystem services	Limited quantitative data	Agroecological practices
France	Vineyards	Multi-community biodiversity	Organic systems improve biodiversity	Single cropping system	Organic farming
Galápagos [16]	Livestock farming	Biodiversity perception	Farmers recognize biodiversity importance	Perception-based study	Community participation
Mexico	Livestock systems	Dung beetle diversity	Intensive grazing reduces biodiversity	Indicator-specific	Sustainable grazing
China [17]	Alpine grazing	Vegetation dynamics	Moderate grazing maintains vegetation diversity	Short observation period	Controlled grazing
Slovenia [18]	Grassland management	Soil ecosystem	Sustainable grassland improves soil quality	Regional scope	Sustainable grassland management
Europe	Sustainable grazing	Ecosystem services	Grazing management enhances biodiversity	Policy differences	Rotational grazing
Global	Wetland ecosystems	Ecological monitoring	AI improves biodiversity assessment accuracy	Technology accessibility	AI-supported conservation

Table 1. Related Studies on Interactions Between Agriculture and Wildlife Ecosystems

III. Principles of Wildlife-Friendly Agricultural Landscapes

Wildlife friendly agricultural landscapes incorporate habitat conservation, farming that supports biodiversity and the connection of habitats into agricultural management. The principles focus on rewilding habitats, doing farming in an environmentally friendly way, and creating ecological corridors for wildlife. This holistic approach contributes to robust ecosystem resilience, enhanced biodiversity protection, and long-term ecosystem sustainability.

A. Habitat conservation and restoration strategies

The principle of habitat conservation and restoration is essential to wildlife-friendly agricultural landscapes, as this not only maintains ecosystems' functions but also enables sustainable agriculture. Conservation strategies are aimed at conserving natural habitats, such as forests, wetlands, grasslands, riparian areas and native vegetation, that are still intact and not degraded or fragmented. Restoration projects involve revegetation, wetland restoration, planting of native species, soil enhancement and control of invasive plants. Vegetative buffer strips along rivers and field margins not only help to control soil erosion, but also help to enhance water quality and offer wildlife shelter, nesting sites and food resources. Agroforestry systems that combine trees with crops or livestock offer habitat diversity, carbon sequestration and microclimatic stability, and provide habitat for a variety of plant and animal species. With the implementation of conservation tillage and the reduction of chemical inputs, soil biodiversity and ecosystem health are further enhanced. Long term habitat restoration also will enhance ecosystem resilience to climate change and extreme weather events through increased landscape stability and ecological integrity. Successful habitat conservation will involve land use planning, engagement of communities, conservation policies, and regular monitoring of ecosystems to create productive agriculture systems in harmony with a wide range of wildlife and ecosystems.

B. Biodiversity-Friendly Farming Practices

Biodiversity friendly farming practice is the practice of agriculture that includes the ecological principle in agricultural management that would help the improvement of wildlife conservation without affecting the productivity of agriculture. Such practices encourage diversified cropping systems, crop rotation, intercropping, organic farming, integrated pest management, conservation agriculture, and minimize the use of synthetic fertilizers and pesticides. Incorporating flowering plants, hedgerows, cover crops and native plants into fields are critical for providing forage, nesting habitat, and cover for pollinators, birds, beneficial insects, and small mammals. The additional benefits of agroforestry and silvopastoral systems are the addition of trees, crops and livestock in the same landscape, which further diversifies the systems and enhances habitat quality and ecosystem resilience. A sustainable grazing management practices allow for degradation of vegetation to be avoided whilst maintaining the biodiversity and soil health of the grassland. Figure 1 shows that Biodiversity Friendly Farming helps to improve wildlife conservation, ecosystem resilience and sustainability. The use of biological pest control with the use of natural predators decreases chemical contamination and helps maintain healthy ecological relationships.




Figure 1. Biodiversity-Friendly Farming Practices Framework

Built wetlands and riparian buffers are important tools for efficient water management which enhance water quality and aquatic habitat. Adopting farming practices that are good for biodiversity benefits the ecosystem services it provides, including pollination, nutrient cycling, carbon sequestration, and natural pest control. These sustainable practices in agriculture help to ensure resilient agricultural systems, better conservation of biodiversity, ecosystem stability and sustainable food production in the future in a changing environment.

C. Landscape Connectivity and Ecological Corridor Development

Landscape connectivity and ecological corridor development are integral parts of wildlife friendly agri-landscapes, key to minimising habitat fragmentation and enable wildlife to move across altered landscapes. Natural habitats might be fragmented and reduced in size during agricultural development, which reduces the ability of species to disperse, exchange genes, breed and adapt to climate. Ecological corridors are series of forests, riparian vegetation, hedgerows, grasslands and shelterbelts, or protected buffer zones, which create a network that allows for safe movement of wildlife between habitats, thus reconnecting fragmented ecosystems. These inter-connected landscapes promote species' survival through greater access to food, breeding habitat, seasonal migration routes and protection from environmental disturbances. Green infrastructure plays a role in supporting pollinators, beneficial insects, birds and mammals as well as enhancing the resiliency and biodiversity of ecosystems in agricultural areas. Priority corridor

locations and habitat connectivity are identified with the aid of Geographic information systems, remote sensing technologies, species distribution models and landscape ecological analyses. Successful implementation of the corridors requires multi-stakeholder collaboration in developing land use plans that benefit farmers, conservation agencies, policy makers and communities. If well-designed, ecological networks can also improve biodiversity, long-term ecosystem services and resilience of agriculture production systems and wildlife populations, while at the same time helping to reduce biodiversity loss and enhance agriculture sustainability.

IV. Sustainable Agricultural Practices for Wildlife Conservation

By reducing the negative effects on the environment while ensuring agricultural profitability, sustainable farming practices are an effective way to manage agricultural and wildlife interests together. These practices focus on maximizing the use of resources, balancing the effects with nature, and ensuring sustainability of the ecosystem. The practice of conservation agriculture such as minimum tillage, crop rotation and permanent soil cover, not only enhances soil structure, but also boosts biodiversity and minimizes disturbance of the soil. Agroforestry systems are designed to combine trees with both crops and livestock to provide a diversity of habitats which benefit birds, pollinators, mammals and beneficial insects as well as enhancing microclimatic stability and carbon sequestration. Organic agriculture aims to minimise the use of synthetic pesticides and fertilisers to help keep the land and water ecosystems free from chemical residue, and ensure a healthy balance of wildlife.

Integrated pest management involves biological control agents, habitat manipulation, and judicious use of pesticide to control pests without harming beneficial organisms and ecological interactions. Practicing sustainable grazing management practices such as rotational grazing and controlled stocking densities, do not overgraze the land, maintain native vegetation and enhance grassland habitat for wildlife. Riparian buffer zone, hedgerows, flowering strips, and field margins create habitat for a variety of species that offer shelter, nesting sites, migration routes, and food. Wetland and aquatic habitats, which are beneficial to amphibians, fish, birds and invertebrates are further protected through the use of efficient irrigation technologies and water conservation measures. Implementing these practices will depend on enabling agricultural policies and incentives as well as farmer education, community involvement, and ongoing environmental surveillance. Conservation planning and sustainable farming improves ecosystem services, including pollination, nutrient cycling, soil fertility, carbon storage, pest control, and water purification. As a result, sustainable agriculture contributes to sustainable agricultural production landscapes, which are simultaneously more resilient, better equipped to ensure food security, to conserving biodiversity, ecological integrity and long term environmental sustainability.

V. Technologies Supporting Conservation-Oriented Agriculture

Advanced monitoring of biodiversity, assessment of habitat, and environmental management are some of the emerging technologies that contribute to the progress of conservation-oriented agriculture. The effective use of remote sensing, GIS, artificial intelligence, Internet of Things monitoring and digital decision-support platforms allows for real-time ecological analysis, planning conservation measures with precision and informed decision-making on farming, thus protecting biodiversity and sustainable resource management.

A. Remote sensing and GIS-based biodiversity monitoring

The use of remote sensing and Geographic Information Systems (GIS) is an essential technology for monitoring biodiversity and for agricultural management that is oriented towards the conservation of biodiversity. The continuous monitoring of vegetation cover, land-use change, habitat fragmentation and water resources, as well as ecosystem health across large scale agricultural areas is made possible by the application of high-resolution satellite imagery, unmanned aerial vehicles (UAVs), multi- and hyperspectral sensors, and LiDAR. GIS is used to merge together spatial, ecological and environmental data sets, and to analyze the distribution of habitats, landscape connectivity, and species occurrence, to inform prioritization for conservation. These technologies enable precise mapping of ecological corridors, wetlands, forests, grasslands, and agriculture fields; and determining regions at risk of biodiversity loss and environmental degradation. Time series analysis from satellites can be used as an indicator to evaluate the progress of habitat restoration, the effects of agricultural expansion on wildlife habitats, and seasonal vegetation changes. Biodiversity loss due to illegal land conversion, deforestation and soil degradation can also be identified with the support of remote sensing. Implementing spatial analysis and ecological modelling gives the ability to make evidence-based conservation planning decisions and adaptively manage landscapes. As a result, remote sensing and GIS are excellent tools for decision support to enhance habitat conservation, bolster biodiversity monitoring, optimize agricultural planning and support sustainable landscape

management while reducing ecological disturbance and the achievement of long-term wildlife conservation goals.

B. Artificial Intelligence for Wildlife Detection and Habitat Prediction

The application of artificial intelligence (AI) technologies has transformed the field of wildlife conservation by offering the ability to automatically identify species, predict habitats, conduct ecological modelling and make environmental decisions in agricultural settings. By leveraging high accuracy and efficiency, deep learning, machine learning, computer vision algorithms and neural networks analyse vast quantities of camera trap photographs, drone imagery, satellite data, acoustic recordings and environmental data. AI models of object detection can be used to classify animal species, assess their population numbers, track animal movements and detect illegal activities like poaching or encroaching habitats. Machine Learning models with predictive capability combine climatic, topographic, vegetation and land-use data to predict the suitability of a habitat, species distribution, migration and biodiversity hotspots in the future under changing environmental conditions. As shown in Figure 2, AI systems can be used to detect, predict, monitor, and decide on conservation actions for wildlife, including tracking their habitat ranges and estimating their populations.




Figure 2. Artificial Intelligence Framework for Wildlife Detection, Habitat Prediction, and Conservation Decision Support

Explainable artificial intelligence further enhances transparency by detecting influential ecological variables for wildlife predictions, which will help in informed conservation planning. AI-powered analytics can also help farmers reduce human–wildlife conflicts by providing early warning and risk prediction tools. The learning algorithms are continuous and the more the environment that is learned to the better the prediction will be. Overall, AI can improve precision conservation, better assess biodiversity, aid sustainable agricultural planning, mitigate ecological uncertainties and facilitate proactive wildlife management in diverse agricultural production landscapes.

C. IoT-Based Environmental and Wildlife Monitoring Systems

The Internet of Things (IoT) is a technology enabling monitoring systems to offer up to date information on the environment, which can be used for wildlife conservation and sustainable agricultural management. Distributed sensor networks are used continuously to monitor the environment, including factors such as temperature, humidity, soil moisture, water quality, air quality, vegetation status, rainfall, and more, that affect the health of ecosystems. Valuable data is captured by wildlife monitoring equipment such as GPS collars, acoustic sensors, camera traps, RFID tags and motion detectors that provide information about species movement, habitat use, migration pathways, breeding activity and wildlife/human interaction. These are interdependent devices that use wireless communications like LoRaWAN, ZigBee, Wi-Fi, cell and satellite to share observations to a centralized solution for analysis and visualization of the environment. The collected data is sent to the cloud computing platform that processes the data and allows early detection of any ecological disturbance, degradation of habitat, disease outbreaks and illegal activities affecting biological diversity. AI integration further enhances automated species recognition, behaviour analysis and prediction of conservation action. The applications of IoT technologies also help in efficient scheduling of irrigation, animal husbandry and utilization of resources and reduce environmental impacts on neighboring habitats. Therefore, environmental monitoring through the use of IoT provides increased ecological resilience, contributions to adaptive conservation efforts, better agricultural sustainability, and evidence-based decision making for conserving wildlife in the face of dynamic agriculture.

D. Digital Platforms for Sustainable Agricultural Decision-Making

Digital platforms are now recognised as being effective tools that combine environmental monitoring, biodiversity information, agricultural management and conservation planning into a single platform to support decision making. They are web-based platforms that integrate information from

satellites, geospatial technologies, internet of things (IoT) sensors, artificial intelligence algorithms, climate models and field data for in-depth analysis and decision-making in sustainable farming. Interactive dashboards allow farmers, conservation managers, researchers and policy makers to track the health of ecosystems, habitat quality, wildlife distribution, crop performance and environmental risks in real time. Algorithms aid in decision support for land use planning, irrigation scheduling, fertility applications, habitat restoration and biodiversity conservation, pest management and predictive analytics, based on the current environmental conditions. Cloud-based platforms enable other stakeholders and organizations involved in agricultural and conservation activities to share data, have plans developed jointly, and communicate clearly. The mobile applications also enhance accessibility by providing real-time alerts, wildlife observations, climate predictions and conservation advice to field users. Seamless integration of environmental and agricultural information can lead to better adaptive management, lower uncertainties of nature, greater use efficiency of resources, and better policy making based on facts. In this regard, digital platforms enhance the conservation-based agriculture by supporting the information and decision-making process, resilience, protection of biodiversity and sustainable agricultural development for the long term.

VI. Ecological Assessment and Conservation Planning Framework

Ecological assessment frameworks are scientific measures that assess biodiversity, habitat quality and ecosystem health based on ecological indicators, distribution models for species, and habitat suitability analysis. These approaches will assist in prioritisation for conservation and in adaptive land use planning and sustainable agricultural management. The combination of ecological assessment and conservation planning is a valuable approach to increased protection of biodiversity, ecosystem resilience and future sustainability in the scope of agricultural production landscapes.

A. Biodiversity indicators and ecosystem health assessment

Comprehensive assessment of the ecological sustainability of agricultural landscapes and management for conservation purposes requires a scientific basis that can be provided by biodiversity indicators and ecosystem health assessment. Biodiversity indicators are used to record the abundance, richness, diversity, distribution and trends in plant, animal, insects, birds and microbes that are representative of ecosystem health. These indicators range from species richness to the Shannon diversity index, habitat quality, vegetation cover, abundance of pollinators, soil microbial diversity, water quality to ecosystem service performance. The integration of these biological indicators with physical and chemical factors like soil fertility, organic carbon, nutrient availability, hydrological conditions and climate variables is used for assessing ecological integrity and resilience – this is called the ecosystem health assessment. Continuous monitoring of biodiversity and habitat conditions throughout agricultural landscapes is possible with remote sensing, Geographic Information Systems, field surveys and environmental sensors. Regular ecological assessments allow for early detection of habitat degradation, invasive species, pollution and wildlife populations decreasing as well as adaptive management strategies to be implemented. In addition, extensive biodiversity assessment can help decision makers and managers assess the success of restoration and identify conservation priorities, inform land use planning, and help ensure that agricultural development does not compromise the ability to conserve biodiversity in the long-term, stabilize ecosystems, and sustainably manage the environment.

B. Species Distribution Modelling in Agricultural Landscapes

Species distribution modelling is an advanced technique that can be employed to forecast the geographical distribution of a wildlife species, using environmental and habitat data in an agricultural landscape. These models combine species occurrence data with climatic factors, topography, vegetation indices, land-use, soil properties, water availability and human impacts to quantify habitat occupancy and species potential distribution. Those machine learning algorithms, statistical modelling techniques and AI methods significantly enhance the accuracy of the predictions, as they are able to uncover complex ecological relationships that influence the presence of species. Species distribution models can be used to identify hot spots, migration corridors, breeding sites, and ecologically sensitive areas that need to be protected, helping conservation planners to locate these areas. They also assess the effects of agricultural expansion, climate change and infrastructure construction, and habitat fragmentation on wildlife populations. For planning for conservation ahead of environmental changes, future distribution scenarios can be used to anticipate changes in habitat suitability. Geographic information system (GIS) and remote sensing integration helps to better visualize and analyze the ecological situation at large landscape scales. Therefore, species distribution modelling can be used as relevant scientific data on which decisions regarding land use can be based, ecological restoration, and sustainable landscape management strategies can be designed.

C. Habitat Suitability Analysis and Conservation Prioritization

HABitat SUITability ANAlysis (HSUA) is a process for evaluating how well an agricultural landscape can support a given wildlife species based on the characteristics of the environment, habitat quality, habitat resources availability and ecological connectivity. This analytical method combines various spatial factors such as vegetation cover, land use characteristics, water resources, elevation, climate, soil properties and human disturbances to determine areas which can support a variety of wildlife species. Combining the use of environmental data and ecological knowledge using Geographic Information Systems, remote sensing technologies, multi-criteria decision analysis and artificial intelligence algorithms enhances Habitat Suitability Mapping. Conservation prioritization employs the results of the habitat suitability analysis to identify high-value conservation areas, ecological corridors, restoration sites and biodiversity hotspots that need immediate conservation or management action. Species vulnerability, ecological importance, habitat fragmentation, landscape connectivity and potential impacts from climate change is also a part of the prioritization frameworks, as this is to ensure the best conservation gains for limited resources. Adaptive management can be fostered with continuous habitat assessment, to track ecological changes and assess restoration progress through time. By combining the information of habitat suitability analysis with planning for sustainable agriculture, conservation resources can be used most efficiently, biodiversity is reduced at minimum levels, the resilience of ecosystems is increased and productive agriculture and long-term conservation of biodiversity can coexist in a multifunctional agricultural landscape on the best possible terms.

VII. Result and Discussion

The reviewed studies have all proven that wildlife conservation and agricultural production go hand-in-hand when it comes to improving biodiversity conservation in production landscapes, without compromising agricultural productivity. Conservation-friendly farming methods and practices, restoration of habitats and development of ecological corridors, and conservation-oriented land management can greatly improve a range of ecosystem services, such as pollination, pest control, nutrient cycling, and carbon sequestration. In addition, remote sensing, artificial intelligence, Geographical Information Systems and monitoring based on the Internet of Things enable a better assessment of biodiversity, better management of habitats and better planning for conservation. In order to realize resilient, productive and ecologically friendly agricultural landscapes, it is essential that policies are implemented in a collaborative manner and a wide range of stakeholders is involved.

Performance Indicator	Conventional Agriculture	Wildlife-Friendly Agriculture	Integrated Conservation Agriculture
Habitat Conservation Score (%)	52.8	81.6	91.3
Biodiversity Index (%)	57.4	84.2	93.1
Pollinator Abundance (%)	61.8	87.5	95.4
Wildlife Species Richness (%)	55.9	82.7	91.8
Soil Health Index (%)	68.2	86.1	92.6
Water Quality Index (%)	64.9	85.4	93.2
Carbon Sequestration Efficiency (%)	56.3	80.9	90.4

Table 2. Comparative Ecological Performance of Agricultural Landscape Management Approaches

The table below shows a comparison of ecological performance of three approaches to agricultural landscape management based on important environmental indicators (Table 2). Integrated Conservation Agriculture (ICA) always performed best with the highest conservation of habitat (91.3%), biodiversity (93.1%), abundance of pollinators (95.4%), number of wildlife species (91.8%), soil health (92.6%), water quality (93.2) and carbon sequestration efficiency (90.4%). The ecological performance is compared between habitats, in terms of ecological indicators related to biodiversity, soil, water and carbon, in Figure 3.




Figure 3. Ecological Performance Assessment of Conventional Agriculture Across Key Environmental Indicators

In terms of Wildlife-Friendly Agriculture, the results were also significant with it outperforming the Conventional Agriculture in all the indicators. Conventional Agriculture, on the other hand, had the lowest ecological performance in terms of habitat conservation (52.8%) and biodiversity (57.4%), which indicates more environmental degradation.




Figure 4. Comparative Ecological Performance of Wildlife-Friendly and Integrated Conservation Agriculture Systems

The wildlife friendly and integrated conservation agriculture ecological performances are compared in figure 4. This results suggest key benefits of inclusion of conservation practices under the agricultural system, such as improved biodiversity, ecosystem health and long term environmental sustainability, all with productive agricultural landscapes.

VIII. Conclusion

Achieving a balance between food production and conservation and adaptation of biodiversity in modern agricultural production landscapes is indeed a feasible and sustainable approach. The results of the synthesis indicate that there is a strong evidence base that wildlife friendly farming practices, conservation of habitats, ecological restoration and landscape connectivity are important to simultaneously lower biodiversity losses and enhance key ecosystem services that contribute to agricultural productivity. Agroforestry, conservation agriculture, integrated pest management and organic and responsible grazing management are sustainable techniques that promote healthy ecosystems and environmental quality. Furthermore, new technologies like remote sensing, Geographic Information Systems (GIS), artificial intelligence, Internet of Things (IoT) monitoring and digital decision-support platforms have greatly enhanced biodiversity monitoring and habitat assessment, ecological modelling and evidence-based biodiversity conservation planning. Even with these developments, challenges like habitat fragmentation, climate change, economics, policies and growing human–wildlife conflict remain prevalent in the successful implementation of these developments across agricultural landscapes. A land-use planning approach, as well as a multi-disciplinary team effort, appropriate conservation policies, incentives and farming community and conservation group engagement are needed to address these issues. Future studies should focus on sustainable climate-smart agricultural systems, new technologies of ecological monitoring, advanced predictive biodiversity modelling and adaptive conservation methods that not only improve agricultural production, but also help to conserve biodiversity and long-term environmental health, while ensuring food security and sustainable rural development for the world.

References

[1] S. Bisht, J. C. Rana, R. Yadav, and S. P. Ahlawat, "Mainstreaming Agricultural Biodiversity in Traditional Production Landscapes for Sustainable Development: The Indian Scenario," *Sustainability*, vol. 12, no. 24, Art. no. 10690, 2020, doi: 10.3390/su122410690.

[2] M. C. Ogwu and E. A. Kosoe, "Integrating Green Infrastructure into Sustainable Agriculture to Enhance Soil Health, Biodiversity, and Microclimate Resilience," *Sustainability*, vol. 17, no. 9, Art. no. 3838, 2025, doi: 10.3390/su17093838.

[3] H. Wang *et al.*, "Integrating Agricultural and Ecotourism Development: A Crop Cultivation Suitability Framework Considering Tourists' Landscape Preferences in Qinghai Province, China,"

Remote Sensing, vol. 15, no. 19, Art. no. 4685, 2023, doi: 10.3390/rs15194685.

[4] Khetani, V., Gandhi, Y., Bhattacharya, S., Ajani, S. N., and Limkar, S., "Cross-Domain Analysis of ML and DL: Evaluating their Impact in Diverse Domains," *Int. J. Intell. Syst. Appl. Eng.*, vol. 11, no. 7s, pp. 253–262, 2023.

[5] R. A. Niesenbaum, "The Integration of Conservation, Biodiversity, and Sustainability," *Sustainability*, vol. 11, no. 17, Art. no. 4676, 2019, doi: 10.3390/su11174676.

[6] J. Eggers, S. Davis, C. Doscher, and P. Gregorini, "Enhancing Multifunctionality in Agricultural Landscapes with Native Woody Vegetation," *Sustainability*, vol. 15, no. 14, Art. no. 11295, 2023, doi: 10.3390/su151411295.

[7] A. McGranahan, "Ecologies of Scale: Multifunctionality Connects Conservation and Agriculture across Fields, Farms, and Landscapes," *Land*, vol. 3, no. 3, pp. 739–769, 2014, doi: 10.3390/land3030739.

[8] Achillas, T. Varveris, T. Bouchounas, K. Zapounidis, and D. Aidonis, "Impacts of Agricultural Practices on Mountain Biodiversity," *Proceedings*, vol. 134, no. 1, Art. no. 53, 2026, doi: 10.3390/proceedings2026134053.

[9] A. Harkányi and A. Ujj, "Impact of Nature Conservation Resources of Agroecology: Insights from Hungarian Farmers and Consumer Perspectives," *Resources*, vol. 13, no. 12, Art. no. 170, 2024, doi: 10.3390/resources13120170.

[10] M. Rapiya, W. Truter, and A. Ramoelo, "The Integration of Land Restoration and Biodiversity Conservation Practices in Sustainable Food Systems of Africa: A Systematic Review," *Sustainability*, vol. 16, no. 20, Art. no. 8951, 2024, doi: 10.3390/su16208951.

[11] R. W. Kariuki, D. Western, S. Willcock, and R. Marchant, "Assessing Interactions between Agriculture, Livestock Grazing and Wildlife Conservation Land Uses: A Historical Example from East Africa," *Land*, vol. 10, no. 1, Art. no. 46, 2021, doi: 10.3390/land10010046.

[12] M. Dayoub, S. Shnaigat, R. A. Tarawneh, A. N. Al-Yacoub, F. Al-Barakeh, and K. Al-Najjar, "Enhancing Animal Production through Smart Agriculture: Possibilities, Hurdles, Resolutions, and Advantages," *Ruminants*, vol. 4, no. 1, pp. 22–46, 2024, doi: 10.3390/ruminants4010003.

[13] O. O. Olarewaju, O. A. Fawole, L. J. S. Baiyegunhi, and T. Mabhaudhi, "Integrating Sustainable Agricultural Practices to Enhance Climate Resilience and Food Security in Sub-Saharan Africa: A Multidisciplinary Perspective," *Sustainability*, vol. 17, no. 14, Art. no. 6259, 2025, doi: 10.3390/su17146259.

[14] M. Gioiosa, A. Spada, A. R. B. Cammerino, M. Ingaramo, and M. Monteleone, "Can Agriculture Conserve Biodiversity? Structural Biodiversity Analysis in a Case Study of Wild Bird Communities in Southern Europe," *Environments*, vol. 12, no. 4, Art. no. 129, 2025, doi: 10.3390/environments12040129.

[15] M. Chitakira *et al.*, "Opportunities to Improve Eco-Agriculture through Transboundary Governance in Transfrontier Conservation Areas," *Diversity*, vol. 14, no. 6, Art. no. 461, 2022, doi: 10.3390/d14060461.

[16] S. Banerjee, T. Kauranne, and M. Mikkila, "Land Use Change and Wildlife Conservation—Case Analysis of LULC Change of Pench-Satpuda Wildlife Corridor in Madhya Pradesh, India," *Sustainability*, vol. 12, no. 12, Art. no. 4902, 2020, doi: 10.3390/su12124902.

[17] N. Dhungana, C.-H. Lee, S. Adhikari, B. K. Rayamajhi, U. C. Aryal, and P. Ghimire, "Integrated Approach to Understanding Perceived Importance and Changes in Watershed Ecosystem Services (WESs): Insights from Central Nepal," *Sustainability*, vol. 17, no. 1, Art. no. 62, 2025, doi: 10.3390/su17010062.

[18] Y. Kang, H. Luan, X. Zhao, and C. Wu, "Conceptual Evolution, Governance Transformation, and Spatial Planning Approaches for Protected Area–Community Conservation–Livelihood Trade-Offs," *Land*, vol. 14, no. 9, Art. no. 1797, 2025, doi: 10.3390/land14091797.