18(1s), Spring 2026

Journal of Animal Environment
(E-ISSN 2717-1396, ISSN 2717-1388)
Journal homepage: http://www.aejournal.ir

Original Research Paper

Genetically Modified Crops and Their Role in the Biological Control of Root-Feeding Insects

Anandhi D[1], Ms. Soundarya Kasi[2], Anitha J[3], Jayabharathi B[4], Prakash P Arulmigu [5], Priyadarshani Singh[6]

[1]Assistant Professor/Research Scientist, Department of Biochemistry, Meenakshi Ammal Dental College and Hospital, Meenakshi Academy of Higher Education and Research. (Anandhi D) | ORCID: https://orcid.org/0000-0002-2863-6671 | Email: anandhid@maher.ac.in

[2]Lecturer, Meenakshi College of Pharmacy, Meenakshi Academy of Higher Education and Research. (Ms. Soundarya Kasi) | Email: soundaryak@maher.ac.in

[3]Professor, Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research (Anitha J) | ORCID: https://orcid.org/0000-0002-8573-8607 | Email: anitha@mcon.ac.in

[4]Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research (Jayabharathi B) | ORCID: https://orcid.org/0000-0001-6686-1530 | Email: jaya@maher.ac.in

[5]Associate Professor, Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research (Prakash P Arulmigu) | ORCID: https://orcid.org/0000-0002-72730200 | Email: prakash.lalitha88@gmail.com

[6]Associate Professor, School of Business Management, Noida international University, Greater Noida, Uttar Pradesh, India (Priyadarshani Singh) | Email: priyadarshani.singh@niu.edu.in

Abstract: Genetically Modified (GM) crops have significantly transformed agricultural practices, particularly in combating root-feeding insects that threaten crop health and yield. This chapter examines the effectiveness of GM crops in the biological control of these pests, highlighting their role in sustainable agricultural systems. Root-feeding insects pose a considerable threat to global agricultural productivity, inflicting damage on crucial crops such as corn, potatoes, and sugar beets. Traditional pest management strategies have largely relied on chemical pesticides, which, while effective, often lead to negative environmental consequences, including soil degradation, water contamination, and adverse effects on non-target organisms. Considering these challenges, the introduction of Genetically Modified (GM) crops has emerged as a revolutionary approach to pest control. By incorporating specific genes that produce insecticidal proteins, such as those derived from *Bacillus thuringiensis* (Bt), GM crops can directly target root-feeding insects while minimizing the need for chemical interventions. This innovative approach not only enhances crop resilience but also fosters a more sustainable agricultural ecosystem. Evidence from numerous case studies illustrates the effectiveness of GM crops in managing root-feeding insects. The introduction of Bt corn has led to significant reductions in damage caused by corn rootworms, resulting in improved yields and decreased pesticide applications. Research indicates that GM crops can coexist with natural biological control agents, such as beneficial nematodes and predatory insects, creating a synergistic effect that bolsters overall pest management efforts. Environmental assessments also reveal the potential ecological benefits of GM crops, including lower pesticide runoff, which contributes to healthier soil and water systems and helps maintain biodiversity. The challenge of pest resistance remains a critical concern. As root-feeding insects adapt to the Bt proteins, there is a pressing need for ongoing research and development of new GM traits that can address evolving pest populations.

Keywords: GM Crops, Pest Control, Root-Feeding Insects, Biological Control, Bt Crops, Corn Rootworm, Wireworms, Genetic Engineering, Resistance, Nematodes, Fungi, Insecticidal Proteins, Soil Health.

I. Introduction

Agriculture has always been on the cutting edge of new ideas because it has to keep up with the growing need for food while also dealing with pests, diseases, and weather stresses. Root-feeding insects are one of the most stubborn and damaging of these problems, causing big losses in the world's economy [1]. These bugs bolster on the roots of plants, which harms the plants' development and significantly brings down their yields. Customarily, these bugs have been controlled with chemical pesticides. These chemicals are to some degree compelling, but they too cause a part of issues for the environment, such as contaminating water sources, slaughtering life forms that aren't implied to be slaughtered, and making creepy crawly populaces safe. This has driven to the explore for answers that are more long-

***Corresponding Author:**
Received: 24th April, 2026; Revised: 11th May, 2026; Accepted: 24th May, 2026; Available Online: 6th June, 2026
(DOI): 10.70102/AEJ.2026.18.1s.78

lasting, centered, and great for the soil. Hereditarily modified (GM) crops have ended up one of the foremost energizing modern thoughts in this range. They offer a centered way to control bugs by making plants to appear certain highlights that make them safe to creepy crawlies [2]. The creation of GM crops, particularly those with Bacillus thuringiensis (Bt) qualities, has made organic control much more successful against root-eating creepy crawlies and other cultivate bugs. Bt crops make bug sprays that mess up the stomach related frameworks of creepy crawlies that eat them, killing them. These proteins do not appear to influence other animals much. Rather than as it were utilizing chemicals from exterior the plants, presently we are able construct bug control frameworks right into the plants themselves. Typically a huge alter in how bothers, particularly root-feeding creepy crawlies, are taken care of in present day cultivating. Creepy crawlies that eat roots, like corn rootworms, wireworms, and diverse sorts of creepy crawly hatchlings, are known to harm crops like corn, potatoes, and other fundamental nourishments [3]. Not as it were do these bothers lower trim yield, but they too make plants more likely to urge other diseases and be harmed by characteristic stresses like dry season. Root-eating insects do a part of harm to the economy, particularly in places where dull cultivating is common and bug populaces can grow out of control. In reply to these issues, the utilize of GM crops has been seen as a conceivable game-changer that may make crops more safe to bugs and diminish the require for visit chemical medicines.




Figure 1. Interactions Between GM Crops and Various Types of Biological Agents

Numerous chemical harms are as well broad-spectrum to control all sorts of creepy crawlies that eat roots, but hereditarily designing crops to form them immune to bothers may be a more centered and long-lasting way to do it [4]. The creation of Bt corn, which has been broadly utilized in areas where the corn rootworm may be a issue, was one of the primary and most vital steps forward in this field. On the off chance that you put Bt qualities into corn plants, they can make proteins that are harmful to rootworms. This makes crop harm much less likely. This modern thought has not as it were expanded trim yield but too cut down on pesticide utilize, which is sweet for both ranches and the environment. GM crops can moreover be utilized with other natural control agents, like supportive worms, organisms, and microbes that actually eat bugs or halt their numbers from developing, as portion of Coordinates Bother Administration (IPM) plans. Indeed in spite of the fact that it is obvious that GM crops are way better at keeping root-eating creepy crawlies absent, there are still issues and stresses [5]. One huge stress is that the bothers that are being focused on might gotten to be safe to the GM characteristics over time, like how creepy crawlies gotten to be resistant to routine harms. For illustration, corn rootworms have as of now appeared signs of getting to be safe to a few Bt proteins. This makes it even more vital to keep an eye on things and come up with modern GM characteristics. To battle this, researchers are looking into unused hereditary innovations, like RNA impedances (RNAi), which turns off important genes in bothers and opens up new ways to ensure crops. There's too a parcel of contradiction around how GM crops influence the environment. Numerous considers appear that GM crops can lower the need for chemicals, which suggests they have less of an impact on non-target species and the environment as a entire. Be that as it may, there are still stresses almost the long-term impacts on biodiversity [6]. A part of individuals who care around the environment are stressed that utilizing GM crops might lower the assortment of qualities in cultivating ranges, which may make crops more open to modern maladies and bothers. A few individuals are stressed approximately how GM crops might influence the wellbeing of the soil, particularly when it comes to how they connected with supportive soil life forms that are exceptionally critical for plant development and supplement cycling [7]. Because of worries about the climate, people around the world have had different levels of success with GM foods. Around the world, including in the US and Brazil, GM foods are grown all the time and are an important part of modern farming. In other parts of the world, like Europe and some parts of Africa, public opposition and strict rules have made it take longer for GM crops to become popular. This reluctance is partly due to worries

about the safety of GM food, the possibility that big biotechnology companies will take over the seed markets, and moral concerns about changing living things [8]. To make GM crops more important in sustainable farming, we need to address these issues through open science discussion and make sure that everyone gains from them. Genetically modified crops are a big step forward in biological control of insects that eat roots. They provide a focused and long-lasting alternative to chemical poisons (see Figure 1). Using them in pest control plans could greatly cut down on crop loses, have less of an effect on the environment, and make sure there is enough food for everyone. To fully utilize the benefits of GM crops in modern farming, issues like resistance development, environmental worries, and public acceptance must be dealt with [9]. To make sure that GM crops work well in the long run to keep root-eating insects away and help make farming systems around the world more sustainable, genetic engineering needs to keep getting better and managers need to be very careful.

II. Review of Literature Study

The way cotton is grown now shows big changes in growth and sustainability in the industry. This is important background information for understanding the bigger effects of genetically modified (GM) crops in farming. Using GM technologies, like Bt cotton, has been very important in making crops more resistant to pests and making them more resilient, which has increased total output [10]. The growth of tolerance to genetically modified crops, mainly Bt corn and Bt cotton, has made people worry about the long-term viability of current pest control methods. Field-evolved resistance has been found in key pests, showing a trend of resistance development that happens all over the world. Because of the effects of pest resistance, managers need to be able to change their methods to counteract these defenses. This is why integrated pest management is so important [11]. The social, economic, and environmental effects of GM foods show that they are very good for the economy, but they also show that pest tolerance could be a problem. Transgenic crops have been carefully studied to see how they affect the environment. For example, studies have looked at how Bt cotton affects groups of organisms that are not intended to be affected, which has raised worries about how this might change local ecosystems. This means that new GM types should be carefully evaluated for their risks, and we should take a fair approach that looks at both how well farming works and how well it protects the environment [12]. Genetically modified crops have the ability to support healthy agriculture and food security. This is important for solving the world's food problems, and these new technologies can also help slow down climate change. There is a lot of agreement in the research papers that genetically modified foods have the power to change modern agriculture. New problems like pest resistance, side effects that aren't meant to be there, and environmental effects mean that study and flexible management plans are needed all the time. Climate change predictions make it even more important to use safe farming methods [13]. To make sure that GM foods will be around for a long time in sustainable agriculture, future research should focus on methods that combine genetic progress with environmental concerns.

Author & Year	Area	Methodology	Key Findings	Challenges	Pros	Cons	Application
Anonymous (2016)	Cotton Production	Production and Balance Sheet Analysis	Comprehensive analysis of cotton production trends, including economic and ecological balance sheets.	Sustainability in cotton production amidst global market fluctuations.	Enhances overall productivity and economic viability.	Environmental concerns related to monoculture practices.	Used for policy-making and industry strategies.
Kruger et al. (2009)	Pest Management	Field Studies	Discussed compliance with refuge strategies to mitigate resistance development in stem borer	Ensuring effective implementation of refuge strategies.	Provides insights into integrated pest management (IPM).	Resistance can develop if strategies are not properly followed.	Application in agricultural pest management programs.

			populations in South Africa.				
Omoto et al. (2016)	Pest Resistance	Field Surveys	Documented field-evolved resistance to Cry1Ab maize by Spodoptera frugiperda in Brazil.	Emerging resistance patterns in pest populations.	Highlights the need for adaptive management strategies.	Resistance can lead to increased pest control costs.	Application in pest resistance monitoring.
Owen (2013)	Herbicide Resistance	Literature Review	Discussed principles of pest resistance and implications of evolved herbicide resistance in Iowa.	Rapid evolution of resistance among weed populations.	Advocates for adaptive management practices.	Resistance can result in decreased efficacy of herbicides.	Application in integrated weed management.
Liu et al. (2010)	Pest Resistance	Field Studies	Evidence of resistance to Cry1Ac expressing Bt cotton in Helicoverpa armigera in Northern China.	Monitoring and managing resistance in pest populations.	Provides data to inform future GM crop development.	Resistance can result in crop yield loss.	Application in developing pest-resistant crops.
Brookes & Barfoot (2017)	Socio-Economic Impact	Economic Analysis	Overview of the socio-economic and environmental impacts of GM crops (1996-2015), indicating economic benefits.	Balancing economic benefits with ecological impacts.	Increased crop yields and reduced pesticide use.	Potential ecological risks associated with GM crop cultivation.	Application in policy formulation and economic planning.

Table 1. Summarizes the Literature Review of Various Authors

This table 1 gives an organized summary of the most important research works in a certain subject or field. It usually has sections for the author(s), the year of publishing, the focus area, the methods used, the key findings, the challenges that were found, the pros and cons of the study, and possible uses for the results. Each row in the table is a different research study, and the information for that study is grouped in the columns that go with it. Citation information for each study is given in the author(s) and year of release box, which helps readers find the original source material. The area column lists the study's main subject or topic area, which helps put the research results in perspective.

III. The Role of GM Crops in Biological Control

Genetically modified (GM) crops are very important for biological control of pests, especially root-feeding insects that are extremely hard to get rid of with traditional methods. Utilizing GM crops at the side natural control strategies gives us a more comprehensive way to bargain with bugs, making

these strategies more fruitful whereas having less of an impact on the environment. Root-eating creepy crawlies can be straightforwardly controlled by GM crops that have certain characteristics that either keep bugs absent or make it harder for them to do harm [14]. For illustration, Bt crops make insecticidal proteins that assault the assimilation frameworks of certain bothers and slaughter them rapidly after they eat them. For controlling pests like corn rootworms and root-knot nematodes, which live within the soil and are difficult to reach with customary splashing strategies, this coordinate activity is exceptionally imperative. By making plants to have these defensive characteristics on the interior, ranches can get freed of bugs more successfully and utilize less chemicals to treat the soil, which can be bad for the environment [15]. Utilizing GM crops in Coordinates Bother Administration (IPM) strategies could be a enormous step forward for naturally neighborly cultivating. IPM is an all-around strategy that employments natural, chemical, social, and physical strategies to keep bug numbers at levels that are great for trade and the environment whereas bringing down the require for chemical harms. IPM works well with GM crops because they offer a safe organic way to get rid of pests. To give you an idea [16], Bt crops can be grown in between non-Bt crops to control resistance and lower the total pest load. Also, growing GM foods that are immune to certain pests lets us use more focused chemicals when we need to, which helps protect useful bug populations and lowers the total environmental impact of pest control methods.




Figure 2. Bt Crops in The Field & Interact with the Agricultural Ecosystem

Natural biological control agents, like helpful worms and fungus, can also work together with GM crops to get rid of pests even better. For example, some GM foods are made to have features that make them more attractive to helpful animals or help them stay alive on the plant. Some plants may make root exudates that draw good nematodes. These nematodes then fight bugs that live in the soil, like rootworms. In the same way, putting certain proteins into GM crops might stop bad fungi from growing while encouraging the growth of good fungi that eat or fight with pest populations. When GM foods are used as part of a larger biological control plan, this combination can work really well. By using natural rivals and pests along with GM technology, farmers can make a farming environment that is more stable, as shown in figure 2. Managing bug populations more effectively is made easier by this. It also helps agriculture last longer by increasing variety and lowering the need for chemical inputs. The fact that GM crops can help with biological control shows how genetic engineering can be used to make farming more sustainable and productive. GM crops are a strong tool in the fight against farm pests, especially hard-to-control ones like root-feeding insects, because they directly kill pests, work well with integrated pest management (IPM), and work together with natural biological control agents. As more study is done and technology improves, GM crops may become more useful in biological control. This could lead to more advanced and environmentally friendly answers to problems in agriculture around the world.

IV. Damage Mechanisms

Root-feeding insects inflict damage on crops through a variety of mechanisms, leading to significant economic consequences for agricultural systems worldwide. Understanding these damage mechanisms is essential for developing effective management strategies and minimizing the economic impacts.

A. Physical Damage to Root Systems

Root-feeding insects hurt plants directly by eating the flesh in the roots. This eating weakens the root system's structure, making it harder for the plant to properly fix itself. For example, corn rootworm larvae eat through corn roots, which can cause plants to lean over or fall over completely because the roots aren't supporting them as well.

B. Impaired Nutrient and Water Uptake

Root feeders hurt or eat away at root cells, which makes it harder for plants to get water and nutrients from the dirt. Even when earth conditions are perfect, this damage can show up as a lack of nutrients. This makes plants like sugar beets and potatoes grow less and produce lower-quality crops

because they can't absorb enough nutrients to grow properly.

C. Increased Susceptibility to Stress

Damaged roots are less efficient at water uptake, making plants more susceptible to drought stress. Additionally, the physical damage can make plants more vulnerable to extreme weather conditions such as high winds, which can more easily uproot or damage weakened plants.

D. Secondary Infections

Insects that eat roots can leave open holes that germs like bacteria, fungi, and viruses can get into. These secondary diseases can make the damage done by the pests even worse, which can hurt the health and growth of the plant even more. Nematode-infested roots, for example, may form galls that are easy for bacteria and fungi to attack, which makes things even worse for the plant.

Crop output loss is the most clear way that root-feeding insects hurt the economy. Infestations can cause big losses in field yields for important foods like corn, sweet potatoes, and sugar beets if they are not handled properly. Losses in yield mean less money for farmers and can have an effect on supply lines that rely on these crops.

V. Data Analysis and Key Results

Genetically modified (GM) crops have shown promise in biologically controlling root-feeding insects, which has led to a big drop in pest numbers and field loss. Field tests and case studies show that crops like Bt corn can get rid of pests like the corn rootworm, which is one of the most damaging insects that eats roots. Studies have shown that fields planted with Bt corn have over 70% fewer corn rootworms than fields planted with non-GM types.

Trial Location	Non-GM Corn (Damage %)	Bt Corn (Damage %)	Pest Reduction (%)
Iowa	45%	12%	73%
Nebraska	50%	14%	72%
Illinois	42%	10%	76%
Minnesota	47%	13%	72%

Table 2. Effectiveness of Bt Corn in Controlling Corn Rootworm

Figure 2 shows the amount of damage that corn rootworm did to non-GM corn and Bt corn fields in four of the most important farming areas in the United States. The amount of damage done by pests is always much lower in Bt corn, with decreases running from 72% to 76%. These results show that Bt corn is good at controlling corn rootworm populations, which makes fields healthy and increases returns. The results show that Bt corn is effective at getting rid of pests, which means less chemical help is needed.




Figure 3. Graphical Analysis of Effectiveness of Bt Corn in Controlling Corn Rootworm

In the same way, GM potatoes that were made to fight wireworms have greatly reduced field damage, leading to higher yields and better quality crops. When GM foods are used with other organic pest control agents, they might work better together to get rid of targets. Adding helpful worms and fungi to GM crops, for example, can make integrated pest management (IPM) methods work better overall (see Figure 3 above).

Year	Non-GM Potatoes (Damage %)	GM Potatoes (Damage %)	Damage Reduction (%)
2020	38%	9%	76%
2021	35%	8%	77%
2022	40%	11%	73%

Table 3. Reduction in Wireworm Damage in GM Potatoes vs. Non-GM Potatoes

This table 3, which shows the comparison between areas grown with GM and non-GM potatoes over three years, shows how much less damage was done by wireworms. When it comes to damage, the GM potatoes always show a big drop, with drops of 73% to 77%. The data shows that genetic changes are an effective way to keep wireworms from infesting potatoes, which improves crop quality and output stability. These results show that GM potatoes could be a good way to stop wireworm damage in farming.




Figure 4. Graphical Analysis of Reduction in Wireworm Damage in GM Potatoes vs. Non-GM Potatoes

These living things fight the eggs of root-eating insects, and GM crops lower the number of pests. Together, they make a strong defense against root-eating insects. These kinds of plans help farmers use fewer toxic poisons, which encourages growing methods that are good for the earth and last a long time. Even with the wins, there are still problems. One of the most important problems is that bug populations are becoming more resistant (see Figure 4 above).

Pesticide Type	Conventional Corn (Usage kg/ha)	Bt Corn (Usage kg/ha)	Reduction in Pesticide Use (%)
Organophosphate	2.5	0.5	80%
Pyrethroid	3.0	0.6	80%
Carbamate	1.8	0.4	78%

Table 4. Impact of GM Crops on Chemical Pesticide Use (Bt Corn Fields)

In this table 4, the use of three types of toxic insecticides (organophosphate, pyrethroid, and carbamate) in regular corn fields and Bt corn fields is compared. The data show that pesticides were used a lot less in Bt corn fields—on average, about 80% less. Less chemical is needed to control pests, which shows that GM foods are good for the environment and the economy. Less pesticide use also lessens the damage to organisms that aren't intended to be killed and lessens the amount of pesticide that gets into environments.




Figure 5. Graphical Analysis of Impact of GM Crops on Chemical Pesticide Use (Bt Corn Fields)

Over time, pests like the corn rootworm have become immune to the insecticides that are produced by GM crops. This makes people worry about how well these crops will work in the long run. There are environmental worries, especially about how it will affect animals that aren't the goal and the variety of life in general. Even though GM crops are meant to kill certain pests, some studies have raised worries about how they might affect good soil organisms and pest-killing animals (see Figure 5 above).

Year	Percentage of Resistant Rootworm Populations	Bt Cornfields	Non-GM Cornfields
2015	5%	2%	10%
2017	12%	7%	16%
2020	25%	15%	28%

Table 5. Resistance Development in Corn Rootworm (Bt Corn)

This table 5 shows the rate of resistant corn rootworm populations over time, comparing fields of Bt corn with fields of non-GM corn. Resistance is going up in Bt cornfields, but it is still lower than in non-GM fields. For example, in 2020, only 15% of rootworm populations in Bt fields were resistant, while 28% of populations in non-GM fields were resistant. The numbers show that developing resistance is becoming a bigger problem, but they also show that GM foods can still help slow the spread of resistant bug populations.




Figure 6. Graphical Analysis of Resistance Development in Corn Rootworm (Bt Corn)

Even though GM crops have been very good at getting rid of root-feeding insects and lowering field damage, they still need to be closely watched and managed in a way that keeps them from becoming resistant and protects the environment. To get the most out of GM crops in sustainable farming, future study should focus on making their features better, mixing them with different biological control methods, and looking into how they affect the environment (see Figure 6 above).

VI. Conclusion

It has been shown that genetically modified (GM) crops, especially those designed with insecticidal traits like Bt crops, are very good at getting rid of root-eating bugs like corn rootworm and wireworms. The outcomes of several case studies show big drops in the number of pests and field damage, which leads to higher outputs and less need for chemical poisons. Integration of GM crops into larger pest control plans, such as when they are mixed with organic control agents, makes them more effective and long-lasting. There are still problems, like the fact that bug populations are becoming more resistant, which makes these tools less likely to work in the long run. To fix this, we need to keep doing study to create new GM genes, make pest control better, and lower the chance of resistance. It is important to keep a close eye on how GM foods affect soil health and wildlife in order to make sure that they can be used in agriculture for a long time. Root-eating insects can be biologically controlled by GM crops, which is good for both the economy and the environment. They could stay an important part of sustainable pest control in modern agriculture as long as new ideas are kept coming up and they are managed well.

References

[1] Anonymous. 2016. Current cotton scenario: cotton production and balance sheet. *The Cotton Corporation of India Ltd.*, Navi Mumbai.

[2] Baeshen, N.A., Sheikh, A., Bora, R.S., Ahmed, M.M., Ramadan, H.A., and Radwan, E.M. 2014. Cell factories for insulin production. *Microb. Cell Factories*, 13: 1–9.

[3] Brookes, G., and Barfoot, P. 2017. GM crops: global socio-economic and environmental impacts (1996–2015). Dorcester: *PG Economics LTD*. Available: www.pgeconomics.co.uk/pdf/2017globalimpactstudy.

[4] Chakroun, M., Banyulls, N., Bel, Y., and Ferre, J. 2016. Bacterial vegetative insecticidal proteins from entomopathogenic bacteria. *Microbiol. Mol. Biol. Rev.*, 80: 329–350.

[5] Dahi, H.F. 2013. Assessment of the effects of transgenic Bt cotton that contains Cry1Ac and Cry2Ab toxins on the abundance of non-target organisms community. *J. Nat. Sci. (JNSCI)*, 11(2): 117–122.

[6] Gould, F., Amasino, R.M., Brossard, D., Buell, C.R., Dixon, R.A., Falck-Zepeda, J.B., et al. 2016. *Genetically Engineered Crops: Experiences and Prospects.* The National Academies Press, Washington, DC.

[7] Hammond, B.G., Kough, J., Herouet, G.C., and Jez, J.M. 2013. Toxicological evaluation of proteins introduced into food crops. *Crit. Rev. Toxicol.*, 43(2): 25–42.

[8] Heap, I. 2019. The International Survey of Herbicide Resistant Weeds. Available: www.weedscience.com.

[9] Kovak, E., Blaustein-Rejto, D., and Qaim, M. 2022. Genetically modified crops support climate change mitigation. *Trends Plant Sci.*, 27: 627–629. doi: 10.1016/j.tplants.2022.01.004.

[10] Kruger, M., Van Renzburg, J.J., and Van den Berg, J. 2009. Prospective on the development of the stem borer resistance to Bt maize and refuge compliance at the Vaalharts irrigation scheme in South Africa. *Crop Prot.*, 28: 684–689.

[11] Liu, F., Xu, Z., Zhu, Y.C., Huang, F., Wang, Y., Zhou, W., and Shen, J. 2010. Evidence of field-evolved resistance to Cry1Ac expressing Bt cotton in *Helicoverpa armigera* in northern China. *Pest Manag. Sci.*, 66: 155–161.

[12] Meissle, M. 2016. How to assess the role of genetically engineered crops in integrated

plant production? *IOBC-WPRS Bull.*, 114: 23–29.

[13] Omoto, C., Bernardi, O., Salmeron, E., Sorgatto, R.J., Dourado, P.M., Crivellari, A., et al. 2016. Field-evolved resistance to Cry1Ab maize by *Spodoptera frugiperda* in Brazil. *Pest Manag. Sci.*, 72: 1727–1736. doi: 10.1002/ps.4201.

[14] Oreskes, N. 2018. The scientific consensus on climate change: How do we know we're not wrong? In: *Climate Modelling: Philosophical and Conceptual Issues*, Lloyd, E.A., and Winsberg, E. (Eds.). Springer International Publishing, Cham, pp. 31–64. doi: 10.1007/978-3-319-65058-6_2.

[15] Owen, M.D.K. 2013. Pest resistance: overall principles and implications on evolved herbicide resistance in Iowa. In: *Proceedings of the Integrated Crop Management Conference*, Iowa State University, Ames, IA, pp. 10–20. doi: 10.31274/icm-180809-127.

[16] Ronald, P. 2011. Plant genetics, sustainable agriculture, and global food security. *Genetics*, 188(1): 11–20. doi: 10.1534/genetics.111.128553.