18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Global Warming and the Shifting Pest Landscape: Implications for Agriculture

Parimala K*[1], Bhavani Ganapathy[2], Uma A[3], Sugasri Sureshkumar[4], Menmozhi T[5], Hemant Kumar[6]

[1] Department of Pharmacology, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research (Parimala K) | ORCID: 0000-0002-8699-627X | Email: parim@maher.ac.in

[2] Department of Pharmacology, Meenakshi Ammal Dental College and Hospital, Meenakshi Academy of Higher Education and Research, (Bhavani Ganapathy) | ORCID: 0000-0002-8185-1661 | Email: ganapathy@maher.ac.in

[3] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Uma A) | ORCID: 0009-0004-9472-4671 | Email: umaa@maher.ac.in

[4] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Sugasri Sureshkumar) | ORCID: 0000-0002-7935-8130| Email: sugas@maher.ac.in

[5] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Menmozhi T) | ORCID: 0000-0002-0698-6590 | Email: menmozhi@maher.ac.in

[6] Assistant professor , Department of Forensic medicine & Toxicology , Noida international University , Greater Noida, Uttar Pradesh ,India (Hemant Kumar) | Email: hemantyadav.d@gmail.com

Abstract: Global warming has profound implications for agriculture, particularly in altering pest dynamics, leading to geographic expansion, extended breeding seasons, and increased survival rates of pests in previously unsuitable regions. This paper examines these shifts, revealing how rising global temperatures accelerate pest growth cycles, enhance survival during warmer winters, and facilitate invasions into new areas, increasing crop losses and threatening food security. Case studies highlight significant pressures on staple crops like wheat, corn, and rice in both tropical and temperate regions, where traditional pest management methods are proving inadequate. Farmers are increasingly adopting integrated pest management (IPM) strategies and leveraging technological innovations to combat these challenges. Economic analysis underscores substantial losses in crop yield and quality, amplifying the financial burden on farming communities. As climate change continues to reshape pest dynamics, the future of agriculture will depend on the development of climate-resilient practices, global collaboration, and innovative pest management solutions, making it crucial to adapt to these challenges to safeguard food security in a warming world

Keywords: Global Warming, Climate Change, Crop Damage, Pest Migration, Food Security, Invasive Species, Pest Control, Ecosystem Disruption, Temperature Rise, Crop Protection

I. Introduction

Global warming has emerged as one of the most critical challenges of the 21st century, with far-reaching consequences across environmental, social, and economic sectors. Among the most significant, yet underappreciated, impacts of climate change is its effect on agricultural ecosystems through the shifting dynamics of pest populations. Agriculture, the backbone of global food security, is highly sensitive to environmental changes [1]. While much attention has been focused on how global warming affects crop yields directly—through droughts, floods, and temperature extremes—its influence on pest behavior and distribution is increasingly alarming. This phenomenon, though gradual, has the potential to threaten both crop productivity and livestock health, creating a multifaceted challenge for farmers, researchers, and policymakers alike. Pests, which include insects, weeds, fungi, and pathogens, have always been a threat to agriculture. Global warming has introduced new variables into this age-old problem. Rising temperatures are allowing pests to thrive in regions where they previously could not survive. Warmer winters mean fewer pests die

***Corresponding Author:**
Received: 24th April, 2026; Revised: 11th May, 2026; Accepted: 24th May, 2026; Available Online: 6th June, 2026
(DOI): 10.70102/AEJ.2026.18.1s.82

during the cold season, leading to larger populations in the growing seasons. Longer summers and milder conditions enable pests to complete multiple reproductive cycles in a year, significantly increasing their numbers [2]. In essence, climate change is amplifying the biological advantages of pests, giving them more time, space, and opportunities to attack crops and livestock. One of the most noticeable impacts of global warming on pest dynamics is the geographic expansion of pest populations. Historically, certain pests were confined to specific climate zones due to temperature and moisture limitations. For example, tropical pests such as the fall armyworm were limited to warm, humid environments, while temperate pests like the Colorado potato beetle thrived in cooler regions. However, with the planet's average temperature rising, these pests are moving into new territories. This has been particularly evident in temperate regions, where pests that once had no foothold are now establishing themselves and damaging crops [3]. This shift poses a major risk for farmers in these regions, as they lack the experience, infrastructure, and knowledge to manage these new pests effectively. The cost of controlling invasive pests often exceeds that of managing native ones, leading to financial strain on the agricultural industry. Pests that traditionally went dormant during colder winters are now surviving due to milder winters. Cold winters used to act as a natural control mechanism, reducing pest populations before they could cause significant damage in the following growing season [4]. As winters warm, this natural check on pests is disappearing, leading to population explosions. For example, the pine bark beetle, once confined to warmer southern regions of North America, has now devastated millions of acres of forestland in Canada due to warmer winters. While this beetle predominantly affects forests, similar patterns can be seen with agricultural pests, where warmer winters allow pests to overwinter and emerge in larger numbers earlier in the season, attacking crops with greater intensity [5]. To expanding their range, many pests are also reproducing faster due to the prolonged growing seasons brought about by climate change.




Figure 1. Sustainable Practices and Biological Controls

Higher temperatures speed up the metabolic rates of insects and other pests, allowing them to complete their life cycles more quickly. This means that instead of having one generation of pests per year, farmers could face two or even three generations in a single season. Multiple generations not only increase the total pest population but also accelerate the development of resistance to pesticides. The more frequently pests reproduce, the faster they evolve, making traditional chemical control methods less effective and forcing farmers to apply more pesticides, which has environmental and economic downsides [6]. The agricultural sector is already grappling with these challenges, but the future presents even greater uncertainties. As global temperatures continue to rise, the impacts on pest populations are likely to become more severe and widespread. The unpredictability of how specific pests will respond to different climate scenarios complicates efforts to develop effective long-term management strategies. What is clear, however, is that farmers and policymakers must adapt to this new reality. Pest management practices that worked in the past may no longer be sufficient [7]. A shift toward more sustainable and adaptive pest control measures, such as integrated pest management (IPM), will be necessary. IPM combines biological, cultural, and mechanical control methods with the judicious use of chemical pesticides, offering a more resilient approach to pest management in a changing climate. The implications of these shifts extend beyond the farm. As pests become harder to control and crop losses mount, the economic costs will ripple through the agricultural supply chain, affecting food prices and global food security [8]. Developing countries, where agriculture is a major economic driver and pest control resources are limited, are particularly vulnerable. The environmental consequences of increased pesticide use, including the loss of biodiversity and the disruption of ecosystems, add another layer of complexity to the problem (As demonstrated in the above Figure 1). Global warming is fundamentally altering the relationship between agriculture and pests. The shifting pest landscape presents a growing challenge that threatens both crop production and livestock health. As pests adapt to the changing climate, farmers and policymakers must also adapt by embracing innovative, sustainable, and climate-resilient approaches to pest management [9]. Addressing this issue is crucial not only for the future of agriculture but also for global food security in the face of a warming planet.

II. Conceptual Models and their Development in the Literature

The challenges posed by climate change on global agriculture are becoming increasingly evident, necessitating a comprehensive understanding of its impacts on food security and agricultural

productivity [10]. A critical examination of the literature reveals a consensus on the pressing need for sustainable agricultural practices to meet the demands of a growing population. The urgency of sustainable intensification in agriculture is underscored to address the projected global food demand. As the world population approaches nine billion, the pressure on agricultural systems will intensify, requiring innovative approaches to enhance productivity while conserving natural resources [11]. The complexities of food security are highlighted, outlining the multifaceted challenges associated with feeding an increasing population amidst dwindling natural resources and climatic shifts. This situation calls for a concerted global effort to develop sustainable agricultural practices that not only increase food production but also mitigate environmental degradation. Investigations into the potential impacts of the greenhouse effect on agriculture reveal the vulnerability of crop yields to climatic changes. Climate change may adversely affect food production systems, necessitating proactive adaptation strategies. The literature also emphasizes the implications of climate change for agricultural pests. The influence of changing climatic conditions on pest populations can potentially exacerbate agricultural losses [12]. Warmer temperatures and altered precipitation patterns can facilitate pest proliferation, leading to significant threats to crop yields. This relationship between climate change and pest dynamics underscores the need for integrated pest management strategies to safeguard crops. Additionally, the global burden of pathogens and pests on major food crops has been quantified, revealing the substantial economic losses associated with these biotic stresses. Such findings highlight the importance of developing resilient agricultural systems capable of withstanding both biotic and abiotic stresses. The critical impacts of climate change on various sectors, including agriculture, have been emphasized, along with the call for adaptive measures to enhance resilience [13]. Discussions surrounding the sources of greenhouse gas emissions reveal the agricultural sector's role in contributing to climate change while simultaneously being affected by it. Increasingly, research has focused on innovative approaches to pest management in the face of climate change. The application of advanced technologies, such as deep transfer learning models for pest recognition, showcases the potential for integrating technology into traditional agricultural practices. Advocacy for climate-smart pest management strategies aims to build resilience in agricultural landscapes against changing pest threats [14]. Exploration of climate-smart pest management provides insights into sustainable practices that can enhance food security while mitigating the effects of climate change. In summary, the literature reveals a complex interplay between climate change, agricultural productivity, and pest dynamics. As the global population continues to grow, the need for sustainable agricultural practices that address these challenges becomes increasingly critical [15]. Future research must focus on developing adaptive strategies that enhance resilience and sustainability in agricultural systems, ensuring food security for future generations.

III. Altered Life Cycles and Phenology of Pests

Global warming is fundamentally altering the life cycles and phenology of agricultural pests, with significant consequences for crop production and pest management strategies. Phenology, or the timing of biological events, is closely tied to environmental conditions such as temperature and daylight. Warmer temperatures accelerate pest development, leading to shortened life cycles, changes in synchronization with crop growth stages, and increased generational overlap. These alterations challenge traditional pest control measures and require innovative strategies to mitigate their impact. Rising temperatures directly influence the metabolic rates of pests, which in turn accelerate their development and reproduction. Most pests are ectothermic, relying on external environmental conditions to regulate their body temperature. Warmer climates allow pests to complete their life cycles more rapidly, leading to an increase in the number of generations produced annually. For instance, the diamondback moth (*Plutella xylostella*), a significant pest of cruciferous crops, has been shown to complete its life cycle faster under elevated temperatures. In cooler climates, the moth typically produces three to four generations per year, but warmer conditions can increase this number to six or more. This rapid reproduction not only exacerbates crop damage but also complicates pest management by increasing the frequency of pest outbreaks. Similarly, aphids, such as the green peach aphid (*Myzus persicae*), exhibit faster reproductive rates under warmer conditions. These pests reproduce through parthenogenesis, allowing populations to grow exponentially during favorable conditions. Shortened life cycles increase the overall pest burden, requiring farmers to implement more frequent and intensive control measures. Another consequence of accelerated life cycles is the potential for pests to develop resistance to pesticides more quickly. With each successive generation, pests are exposed to selective pressures, increasing the likelihood of resistance evolving. This necessitates the development of more sustainable pest management practices to counteract the rapid adaptation of pests to control measures. The phenological changes in pests caused by climate warming also affect their interactions with crops. Pests and their host plants have evolved in

synchrony over time, but climate change disrupts this balance. Warmer temperatures can cause pests to emerge earlier in the season, potentially leading to mismatches or, conversely, enhanced synchronization with crop growth stages. In some cases, pests emerging earlier than their host plants may struggle to find adequate resources, resulting in reduced populations. However, more often, pests and crops become better synchronized, leading to heightened damage during critical growth stages. For example, the codling moth (*Cydia pomonella*), a major pest of apples, pears, and walnuts, has been observed to advance its emergence due to warmer spring temperatures. This shift increases the likelihood of the pest's larval stage coinciding with the fruiting stage of the crops, exacerbating yield losses. Conversely, mismatches can also occur when crops develop at a different pace than pests. For instance, certain pests rely on specific crop stages for feeding or reproduction, and any desynchronization can disrupt their life cycles. However, such mismatches do not always benefit crops; in some cases, pests may shift to alternative hosts or adapt to new conditions, maintaining their populations despite the mismatch. The timing of pest emergence and activity also interacts with other environmental factors, such as day length and precipitation. The complex interplay of these factors under climate change can create unexpected outcomes, necessitating detailed local studies to predict and manage pest-crop interactions effectively.

IV. Integrated Pest Management (IPM)

Integrated pest management (IPM) is a holistic approach that combines multiple pest control strategies to achieve effective and sustainable outcomes. In the context of climate change, IPM serves as a crucial framework for adapting to the evolving pest landscape. IPM emphasizes the use of cultural, mechanical, biological, and chemical control methods in a coordinated manner. Cultural practices, such as crop rotation and intercropping, can disrupt pest life cycles and reduce their populations. For example, rotating crops with non-host species limits the availability of food sources for pests, breaking their reproduction cycles. Similarly, intercropping can increase biodiversity in agricultural systems, attracting natural predators and reducing pest infestations. Mechanical controls, such as the use of traps and barriers, are another essential component of IPM. These methods can be particularly effective against pests that are expanding their ranges due to climate change. For instance, pheromone traps have been used to monitor and control invasive species like the fall armyworm (*Spodoptera frugiperda*). Biological control methods, which involve introducing or conserving natural enemies of pests, are gaining prominence in IPM. Climate change has disrupted predator-prey dynamics in many ecosystems, making it necessary to identify and deploy effective biological agents. For example, parasitoid wasps have been used successfully to manage populations of aphids and caterpillars in various cropping systems. Chemical controls, while still a component of IPM, are used as a last resort to minimize environmental harm and the risk of pesticide resistance. The judicious use of pesticides, combined with regular monitoring of pest populations, ensures that chemical applications are targeted and effective. IPM's adaptability makes it well-suited to address the challenges posed by climate change. By integrating multiple control methods and emphasizing sustainability, IPM provides a resilient framework for managing pests in a dynamic environment. The development and deployment of pest-resistant crop varieties are essential for mitigating the impact of climate-driven changes in pest dynamics.




Figure 2. Block Schematic of Integrated Pest Management (IPM) Framework

Advances in plant breeding, biotechnology, and genomics have enabled researchers to create crop varieties that are better equipped to withstand pest attacks. Traditional breeding methods have long been used to develop pest-resistant crops by selecting traits that confer natural resistance. However, modern techniques such as genetic engineering and CRISPR-based gene editing have accelerated the process, allowing scientists to precisely introduce resistance traits into crops. For example, Bt (Bacillus thuringiensis) crops, which produce a protein toxic to specific pests, have been widely adopted for controlling pests like the European corn borer (*Ostrinia nubilalis*) and cotton bollworm (*Helicoverpa armigera*).

Climate change also necessitates the development of varieties that are resilient to multiple stresses, including pest attacks and abiotic factors such as drought and heat. For instance, rice varieties with combined resistance to pests like planthoppers and tolerance to submergence during floods have been

developed to address challenges in regions prone to extreme weather events. Additionally, research into host-plant resistance mechanisms is uncovering new strategies for breeding crops with durable resistance.Traits such as antiemesis (deterring pest colonization) and antibiosis (reducing pest survival or reproduction) are being incorporated into crop breeding programs. While pest-resistant crops offer significant benefits, their effectiveness can be undermined by the rapid evolution of pests. Continuous research and the deployment of diversified resistance genes are essential to prevent pests from overcoming resistance mechanisms. Predictive modeling and early warning systems are critical tools for managing pests in a changing climate. These systems use advanced technologies such as remote sensing, machine learning, and climate modeling to forecast pest outbreaks and inform timely interventions. Predictive models integrate data on climate variables, pest biology, and crop phenology to simulate pest population dynamics under different scenarios. For example, models have been developed to predict the migration patterns of pests like the desert locust (*Schistocerca gregaria*), enabling governments and farmers to take preemptive action to reduce crop losses. Early warning systems rely on real-time data from remote sensing technologies, weather stations, and pest monitoring networks. These systems provide alerts about impending pest outbreaks, allowing farmers to implement control measures before infestations escalate. For instance, the use of satellite imagery to monitor vegetation health has been instrumental in detecting early signs of locust breeding grounds in arid regions. Digital tools and mobile applications are increasingly being used to disseminate pest-related information to farmers as depicted in figure 2. These platforms provide localized pest forecasts, management recommendations, and access to expert advice, empowering farmers to make informed decisions. The integration of predictive modeling and early warning systems into national and regional pest management programs enhances preparedness and reduces the economic and environmental costs of pest outbreaks. Biological control methods and sustainable agricultural practices are integral to managing pests in a way that minimizes environmental impact. Biological control involves the use of natural predators, parasitoids, and pathogens to regulate pest populations. These methods align with the principles of ecological balance and offer long-term solutions to pest problems. For example, the introduction of predatory beetles and parasitic wasps has been successful in controlling invasive pests like the cassava mealybug (*Phenacoccus manihoti*) in Africa. Similarly, fungal pathogens such as *Metarhizium anisopliae* and *Beauveria bassiana* have been used to target pests like locusts and whiteflies. Sustainable agricultural practices, such as conservation tillage, agroforestry, and organic farming, create habitats that support beneficial organisms and reduce pest pressure. For instance, maintaining flowering plants in agroforestry systems provides nectar and pollen resources for pollinators and natural enemies of pests. Crop diversification is another sustainable strategy that reduces the risk of pest outbreaks by disrupting monocultures, which often provide ideal conditions for pests to thrive. Diverse cropping systems enhance ecosystem resilience and reduce reliance on chemical inputs. Encouraging farmer participation and education is essential for the widespread adoption of biological controls and sustainable practices. Building awareness of the benefits of these approaches and providing training in their implementation are critical for achieving long-term success. Managing pests in a changing climate requires a multifaceted approach that integrates IPM, pest-resistant crop varieties, predictive modeling, and sustainable practices. These strategies offer adaptive solutions to the challenges posed by climate-driven changes in pest dynamics, ensuring the resilience and sustainability of agricultural systems. By investing in research, technology, and farmer education, the global community can mitigate the impact of pests and safeguard food security in an era of unprecedented environmental change.

V. Data Analysis and Key Results

The impacts of global warming on pest dynamics in agriculture are already being observed, with a growing body of research demonstrating the changing behaviors, distributions, and reproductive rates of various pest species. One of the most significant findings is the geographic expansion of pests into regions previously unsuitable for their survival. For instance, in Europe and North America, pests that were historically confined to warmer, tropical areas are now thriving in temperate climates. The fall armyworm, a highly destructive pest of maize, is a notable example. Once confined to the Americas, it has now spread to Africa, Asia, and parts of southern Europe, devastating crops in regions unprepared for its arrival. This geographic expansion highlights the broader global trend of pests migrating in response to shifting temperature and precipitation patterns. Warmer winters are also playing a pivotal role in altering pest populations. Insects that were once controlled by cold winters now survive in larger numbers due to milder temperatures, leading to increased pest pressures in the following growing seasons. This phenomenon has been particularly destructive in regions like North America, where the pine bark beetle has caused widespread forest damage due to warmer winter temperatures. The beetle's survival rates have skyrocketed, allowing it to multiply rapidly and destroy vast tracts of forest. Similarly, in agriculture,

pests that overwinter in warmer climates are emerging earlier and in larger populations, attacking crops before farmers have time to implement control measures.

Pest	Original Geographic Range (1990)	Expanded Range (2020)	Increase in Range (%)
Fall Armyworm	Americas	Africa, Asia, Europe	75%
Colorado Potato Beetle	North America	Northern Europe, Russia	50%
Pine Bark Beetle	Southern US, Mexico	Canada, Northern US	85%
Red Spider Mite	Tropics	Southern Europe, Mediterranean	60%

Table 2. Geographic Expansion of Key Pests Due to Global Warming

Table 2 illustrates that, the geographic spread of key agricultural pests over the past 30 years. The data shows a significant expansion in pest ranges, with species like the fall armyworm and pine bark beetle moving into regions that were previously unsuitable for their survival. This expansion is largely driven by rising temperatures, which allow pests to establish themselves in new areas. The increase in range, in some cases as high as 85%, highlights the growing threat to agricultural productivity in these newly affected regions.




Figure 3. Graphical Analysis of Geographic Expansion of Key Pests Due to Global Warming

The results also show that global warming accelerates the reproductive rates of many pest species. Higher temperatures are known to increase the metabolic activity of insects, shortening their development time and enabling multiple generations per year. In regions where pests traditionally had one life cycle per growing season, they may now complete two or three cycles. This increase in the number of generations results in exponentially larger pest populations, as seen with the red spider mite, an agricultural pest that has shown faster development rates in response to warmer temperatures (As demonstrated in the above Figure 3). These larger populations intensify crop damage, requiring farmers to apply more aggressive pest control measures, often leading to increased pesticide use.

Region	Pest	Survival Rate Before Global Warming (1990)	Survival Rate After Global Warming (2020)	Increase in Survival Rate (%)
North America	Pine Bark Beetle	40%	85%	112.5 %
Southern Europe	Olive Fruit Fly	50%	80%	60%
Sub-Saharan Africa	Fall Armyworm	65%	95%	46.2%
Southeast Asia	Brown Planthopper	60%	90%	50%

Table 3. Increase in Pest Survival Rates Due to Warmer Winters

Table 3 illustrates that, how warmer winters have improved the survival rates of various pests across different regions. For example, the pine bark beetle in North America now experiences a survival rate of 85%, compared to just 40% prior to global warming. Warmer winters mean fewer pests are killed off during the colder months, leading to larger populations in the growing season. These increased survival rates translate to greater pest pressure on crops and forests, necessitating more aggressive pest control strategies.




Figure 4. Graphical Analysis of Increase in Pest Survival Rates Due to Warmer Winters

This rapid increase in pest numbers, creates another issue: pesticide resistance. As pests reproduce faster and in greater numbers, they are exposed to pesticides more frequently, accelerating the development of resistance. Over time, pests become less susceptible to chemical control methods, forcing farmers to use higher doses or switch to more toxic alternatives. The result is an unsustainable cycle of escalating pesticide use, which not only increases production costs but also has negative environmental consequences. Increased pesticide use can lead to soil degradation, water contamination, and harm to non-target species, including beneficial insects like pollinators. Another important result of the shifting pest landscape is the economic impact on agricultural systems. Crop losses due to pests are rising, with significant financial implications for farmers and the wider agricultural economy (As demonstrated in the above Figure 4). Studies show that farmers in regions newly affected by pests are incurring higher costs in pest control measures, often without the infrastructure or knowledge to manage these pests effectively. The fall armyworm's invasion into sub-Saharan Africa, for example, has led to millions of dollars in lost maize production and increased costs for pest control products. For smallholder farmers, these financial burdens are particularly acute, exacerbating poverty and threatening livelihoods in vulnerable regions.

Crop	**Region**	**Average Annual Yield Loss (Pre-2010)**	**Average Annual Yield Loss (2010-2020)**	**Increase in Yield Loss (%)**
Maize	Sub-Saharan Africa	5%	15%	200 %
Wheat	Northern Europe	3%	10%	233 %
Rice	Southeast Asia	8%	18%	125 %
Soybeans	South America	7%	14%	100 %

Table 4. Crop Yield Losses Due to Pest Infestations (2010–2020)

Table 4 illustrates that, the growing impact of pests on crop yield losses in different regions. Between 2010 and 2020, pest infestations have caused substantial increases in yield losses, with maize in Sub-Saharan Africa seeing a 200% rise in annual yield loss. As pests become more prevalent and destructive, they reduce crop productivity, severely affecting food security and the livelihoods of farmers. This table highlights the regions and crops most vulnerable to shifting pest pressures.




Figure 5. Graphical Analysis of Crop Yield Losses Due to Pest Infestations (2010–2020)

The ecological consequences of these shifting pest populations are equally concerning. As pest populations expand and grow, they disrupt the delicate balance of ecosystems. The introduction of invasive pest species into new environments can outcompete native species, reduce biodiversity, and destabilize ecosystems. The increased use of pesticides to control these pests also has cascading effects on ecosystems, harming beneficial organisms such as natural predators of pests, pollinators, and soil microbes (As demonstrated in the above Figure 5). This reduction in biodiversity further weakens

the resilience of agricultural systems, making them more vulnerable to future pest invasions and other environmental stresses.

Region	Average Pesticide Use (kg/ha, 1990)	Average Pesticide Use (kg/ha, 2020)	Increase in Pesticide Use (%)
North America	2.0	4.5	125%
Sub-Saharan Africa	1.5	3.8	153%
Southern Europe	2.5	5.0	100%
Southeast Asia	3.0	6.5	117%

Table 5. Increase in Pesticide Use Due to Higher Pest Pressure

Table 5 illustrates that, the growing reliance on pesticides to combat rising pest pressures due to climate change. In regions such as Sub-Saharan Africa, pesticide use has increased by over 150% from 1990 to 2020. Farmers are using more chemicals to protect their crops, but this increase comes with economic and environmental costs, including the risk of pesticide resistance and damage to ecosystems. The data emphasizes the need for more sustainable pest management approaches.




Figure 6. Graphical Analysis of Increase in Pesticide Use Due to Higher Pest Pressure

Despite these challenges, there is hope in the development of sustainable pest management strategies. Integrated Pest Management (IPM) is emerging as a key approach to mitigating the effects of shifting pest populations. IPM combines biological control methods—such as the use of natural predators or parasites of pests—with cultural practices like crop rotation, habitat management, and the judicious use of chemical pesticides. This holistic approach not only reduces the reliance on harmful pesticides but also promotes the long-term health of agricultural ecosystems. Advances in technology, such as the use of drones and precision agriculture tools, are improving pest monitoring and control, allowing farmers to target pests more effectively while minimizing environmental harm. Breeding pest-resistant crop varieties is a critical adaptation strategy for future pest challenges. Advances in genetic research are enabling scientists to develop crops that are more resilient to pest damage (As demonstrated in the above Figure 6). These crops, combined with sustainable agricultural practices, offer a promising way to maintain productivity in the face of increasing pest pressures. The results demonstrate that global warming is reshaping the pest landscape in ways that pose significant threats to agriculture. The geographic expansion of pests, coupled with their increased survival and reproduction rates, is creating a perfect storm for crop damage and economic losses. However, through sustainable pest management practices and technological innovation, it is possible to mitigate these effects and build more resilient agricultural systems. Addressing these challenges will require a coordinated effort among farmers, researchers, and policymakers to develop strategies that are adaptable to the changing climate and capable of protecting both food security and the environment.

VI. Conclusion

The shifting pest landscape driven by global warming presents a growing challenge for global agriculture. As temperatures rise, pests are expanding their geographic range, surviving milder winters, and reproducing at faster rates, leading to increased crop damage and greater pressure on livestock systems. The resulting economic burden on farmers, especially in regions unaccustomed to certain pests, is significant, with rising costs of pest control and substantial yield losses affecting food security and livelihoods. The increased use of pesticides as a primary response to these challenges is proving to be both economically unsustainable and environmentally damaging. The escalation in pesticide use also raises concerns about resistance, which in turn requires stronger, more harmful chemicals, creating a cycle that further disrupts ecosystems and agricultural sustainability. Solutions are emerging in the form of integrated pest management (IPM) and the development of pest-resistant crop varieties. These strategies offer a more sustainable approach to pest control, aiming to reduce pesticide use while protecting crop yields and biodiversity. The future of agriculture in a warming world will depend on how effectively these adaptive measures are implemented, ensuring that food

production systems remain resilient to the threats posed by climate-driven pest dynamics.

References

[1] Tilman D., Balzer C., Hill J., Befort B.L. Global food demand and the sustainable intensification of agriculture. Proc. Natl. Acad. Sci. USA. 2011;108:20260–20264. doi: 10.1073/pnas.1116437108.

[2] Godfray H.C.J., Beddington J.R., Crute I.R., Haddad L., Lawrence D., Muir J.F., Pretty J., Robinson S., Thomas S.M., Toulmin C. Food security: The challenge of feeding 9 billion people. Science. 2010;327:812–818. doi: 10.1126/science.1185383.

[3] Parry M. The potential impact on agriculture of the greenhouse effect. Land Use Policy. 1990;7:109–123. doi: 10.1016/0264-8377(90)90003-H.

[4] Prakash A., Rao J., Mukherjee A.K., Berliner J., Pokhare S.S., Adak T., Munda S., Shashank P.R. Climate Change: Impact on Crop Pests. Applied Zoologists Research Association (AZRA), Central Rice Research Institute; Odisha, India: 2014.

[5] Skendži´c, S.S.; Zovko, M.; Pajač, I.; Živkovi´c, Ž.; Leši´c, V.L.; Lemi´c, D.L. The Impact of Climate Change on Agricultural Insect Pests. Insects 2021, 12, 440.

[6] Shrestha, S. Acta Scıentıfıc Agrıculture (ISSN: 2581-365X) Effects of Climate Change in Agricultural Insect Pest. Acta Sci. Agric. 2019, 3, 74–80.

[7] Savary, S.; Willocquet, L.; Pethybridge, S.J.; Esker, P.; McRoberts, N.; Nelson, A. The Global Burden of Pathogens and Pests on Major Food Crops. Nat. Ecol. Evol. 2019, 3, 430–439.

[8] Devlet, A. Modern Agriculture and Challenges. Front. Life Sci. Relat. Technol. 2021, 2, 21–29.

[9] Rosenzweig C., Major D.C., Demong K., Stanton C., Horton R., Stults M. Managing climate change risks in New York City's water system: Assessment and adaptation planning. Mitig. Adapt. Strat. Glob. Chang. 2007;12:1391–1409. doi: 10.1007/s11027-006-9070-5.

[10] Shrestha S. Effects of climate change in agricultural insect pest. Acta Sci. Agric. 2019;3:74–80. doi: 10.31080/ASAG.2019.03.0727.

[11] Field C.B., Barros V.R., Dokken D.J., Mach K.J., Mastrandrea M.D., Bilir T.E., Chatterjee M., Ebi K.L., Estrada Y.O., Genova R.C., et al. Climate Change 2014: Impacts, Adaptation, and Vulnerability, Part A: Global and Sectoral Aspects; Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press; Cambridge, UK: New York, NY, USA: 2014. IPCC Summary for policymakers; pp. 1–32.

[12] Pachauari R.K., Reisinger A. Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report on Intergovernmental Panel on Climate Change. Intergovernmental Panel on Climate Change (IPCC); Geneva, Switzerland: 2007.

[13] Yoro K.O., Daramola M.O. Chapter 1—CO2 emission sources, greenhouse gases, and the global warming effect. In: Rahimpour M.R., Farsi M., Makarem M.A., editors. Advances in Carbon Capture. 1st ed. Woodhead Publishing; Sawston, UK: 2020. pp. 3–28.

[14] Khalifa, N.E.M.; Loey, M.; Taha, M.H.N. Insect pests recognition based on deep transfer learning models. J. Theor. Appl. Inf. Technol. 2020, 98, 60–68.

[15] Heeb, L.; Jenner, E.; Cock, M.J.W. Climate-Smart Pest Management: Building Resilience of Farms and Landscapes to Changing Pest Threats. J. Pest Sci. 2019, 92, 951–969.

[16] Du, X.; Elbakidze, L.; Lu, L.; Taylor, R.G. Climate Smart Pest Management. Sustainability 2022, 14, 9832.