18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Bacillus thuringiensis: Mechanisms of Action and Synergism with Biotechnological Tools

Parimala K*[1], Prathiba S[2], Uma A[3], Menmozhi T[4], Balamurugan N[5], Karan Srivastava[6]

[1] Department of Pharmacology, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research (Parimala K) | ORCID: 0000-0002-8699-627X | Email: parim@maher.ac.in

[2] Lecturer, Department of Pharmacology, Meenakshi Ammal Dental College and Hospital, Meenakshi Academy of Higher Education and Research (Prathiba S) | ORCID: 0000-0002-5090-5903 | Email: prathibhas@maher.ac.in

[3] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Uma A) | ORCID: 0009-0004-9472-4671 | Email: umaa@maher.ac.in

[4] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Menmozhi T) | ORCID: 0000-0002-0698-6590 | Email: menmozhi@maher.ac.in

[5] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Balamurugan N) | ORCID: 0000-0002-0778-902X |Email: bala@maher.ac.in

[6] Assistant Professor, Department of Pharmacology , Noida international University , Greater Noida, Uttar Pradesh ,India (Karan Srivastava) | Email: kman180386@gmail.com

Abstract: Bacillus thuringiensis (Bt) has emerged as a pivotal biotechnological tool in sustainable agriculture, renowned for its insecticidal properties. This paper explores the mechanisms of action of Bt, its efficacy in pest control, the development of resistance among pests, and its synergistic potential with modern biotechnological tools. Bacillus thuringiensis, a soil-dwelling bacterium, produces proteins toxic to specific insect larvae, making it a cornerstone in pest management strategies. Its historical application dates back to the 1930s, with significant advancements in biotechnology leading to the development of genetically modified crops expressing Bt proteins. The importance of Bt in modern agriculture lies in its ability to reduce reliance on chemical pesticides, promote environmental sustainability, and enhance crop yields. The analysis reveals that Bt crops significantly reduce pest infestation rates, with reductions averaging over 80% compared to non-Bt counterparts. The data also indicate a gradual development of resistance in key pest species over time, underscoring the necessity for effective resistance management strategies. Environmental assessments demonstrate that Bt crops mitigate pesticide use and decrease non-target organism mortality, enhancing overall ecological health. Bacillus thuringiensis remains an essential component of integrated pest management, balancing efficacy with environmental sustainability. Continued innovation, including the combination of Bt with other biotechnological advancements, is vital for enhancing pest control effectiveness and mitigating resistance. Future research and responsible application of Bt technologies will be crucial in addressing the growing challenges of agricultural sustainability and food security.

Keywords: Cry Toxins, Vip Proteins, Insecticide, Transgenic Crops, Resistance, Bioinsecticide, Pest Control, Gene Pyramiding, Biopesticide, Toxin Synergy.

I. Introduction

Bacillus thuringiensis (Bt) is a Gram-positive, soil-dwelling bacterium that has become an indispensable tool in pest control and agricultural biotechnology. First discovered in 1901 by the Japanese biologist Ishiwata Shigetane and later isolated in 1911 by Ernst Berliner, Bt has garnered global attention due to its ability to produce proteins that are toxic to certain insect species [1]. These proteins, primarily Cry and Vip toxins, are highly specific to insect pests and have little to no effect on humans, animals, or beneficial insects, making Bt a popular choice for integrated pest management (IPM) systems. The bacterium forms endospores and produces crystal proteins during sporulation, which act as potent insecticides when ingested by insect larvae. Bt operates through a highly specific mode of action, targeting the digestive system of insect larvae [2]. Upon ingestion, Bt toxins bind to receptors in the insect's gut, leading to pore

***Corresponding Author:**

Received: 24th April, 2026; Revised: 11th May, 2026; Accepted: 24th May, 2026; Available Online: 6th June, 2026

(DOI): 10.70102/AEJ.2026.18.1s.84

formation and subsequent disruption of the gut cells. This ultimately causes paralysis, starvation, and death of the insect. Because of its specificity, Bt has been extensively used in various formulations, such as sprays and dusts, and has become the genetic basis for numerous transgenic crops engineered to resist pests.

A. Historical Perspective of Bt Use in Pest Control

The potential of *Bacillus thuringiensis* in pest control was recognized as early as the 1920s, but it wasn't until the 1950s and 1960s that Bt-based products became commercially available. Early formulations consisted of Bt spores and crystalline toxins applied as biopesticides in agriculture [3]. These formulations were considered groundbreaking at the time, as they provided a natural and targeted means of controlling agricultural pests, reducing reliance on synthetic chemical insecticides. Bt was particularly effective against Lepidopteran pests (caterpillars), such as the European corn borer, which were major threats to crops like corn, cotton, and vegetables. The adoption of Bt as a bioinsecticide rapidly increased during the 1970s and 1980s due to growing concerns about the environmental and health impacts of chemical pesticides. Bt offered a solution that was both environmentally sustainable and effective, contributing to improved crop yields and reduced pesticide residues in the food supply. Governments and regulatory agencies supported the use of Bt-based products because of their safety profile, leading to widespread adoption by organic and conventional farmers alike [4]. It was the advent of genetic engineering in the 1990s that truly revolutionized the use of Bt in pest control. Scientists successfully inserted genes encoding Bt toxins into crops, creating transgenic plants that could produce their own insecticidal proteins. This innovation eliminated the need for external pesticide applications, providing continuous protection from insect pests throughout the growing season.




Figure 1. Bt in Modern Agriculture and Biotechnology

The first Bt crops, including Bt corn and Bt cotton, were commercialized in the mid-1990s and were quickly adopted by farmers around the world [5]. These crops were designed to combat some of the most damaging pests, such as the corn borer and the cotton bollworm, leading to significant reductions in pesticide use and associated environmental impacts. The widespread success of Bt crops has made them a cornerstone of modern pest management strategies. Today, Bt remains a critical component of integrated pest management (IPM) programs and plays a vital role in reducing the use of synthetic insecticides, preserving beneficial insect populations, and mitigating the environmental

footprint of agricultural practices [6]. In modern agriculture, *Bacillus thuringiensis* plays an essential role in promoting sustainable and environmentally friendly farming practices. Bt-based bioinsecticides are valued for their specificity to insect pests, minimal impact on non-target species, and reduced likelihood of bioaccumulation in ecosystems compared to synthetic chemical pesticides [7]. This makes Bt an attractive option for both organic and conventional farming systems, particularly in regions where pest outbreaks are a significant threat to crop productivity. One of the most significant advancements in agricultural biotechnology has been the development of Bt transgenic crops. These genetically modified crops contain one or more Bt toxin genes, enabling them to produce insecticidal proteins internally. As a result, Bt crops provide consistent, season-long protection against target pests, reducing the need for repeated pesticide applications [8]. The adoption of Bt crops has led to substantial environmental and economic benefits, including reduced chemical pesticide use, decreased labor and operational costs, and improved crop yields. Notably, Bt crops have been shown to have a positive impact on biodiversity by helping to preserve beneficial insect populations, such as pollinators and natural predators of pests, which might otherwise be harmed by broad-spectrum insecticides. The success of Bt in agriculture extends beyond transgenic crops [9]. Advances in biotechnology have enabled the development of new Bt-based formulations and delivery systems, improving the stability, effectiveness, and application of Bt bioinsecticides. These innovations include encapsulation techniques that protect Bt toxins from environmental degradation, as well as the use of nanotechnology to enhance the precision of Bt applications in the field. Bt has been integrated into IPM strategies, where it is combined with other biological, chemical, and cultural pest control methods to achieve optimal results with minimal environmental impact. Beyond pest control, Bt is also an important tool in scientific research. The bacterium's unique ability to produce insecticidal proteins has made it a model organism for studying protein structure, gene expression, and microbial ecology. Researchers are continuously exploring new ways to harness Bt's capabilities, including developing novel Bt toxins that target resistant insect species and expanding Bt's use to control pests in forestry, aquaculture, and urban environments. Despite its many benefits, the use of Bt in agriculture faces challenges, particularly the development of resistance in insect populations [10]. As with any pest control method, overreliance on Bt can lead to the selection of resistant insect strains, reducing the effectiveness of both Bt bioinsecticides and Bt crops. To address this, scientists are employing strategies such as gene pyramiding (stacking multiple Bt toxin genes in a single crop) and implementing refuge strategies, where non-Bt crops are planted alongside Bt crops to slow the evolution of resistance. Biotechnological tools like RNA interference (RNAi) and CRISPR-Cas gene editing offer promising avenues for enhancing the sustainability of Bt-based pest control (As presented in Figure 1 above). *Bacillus thuringiensis* has emerged as a powerful ally in the fight against agricultural pests, providing an eco-friendly alternative to conventional insecticides [11]. Its integration into biotechnology has expanded its utility, offering innovative solutions to pest management challenges. As research continues, Bt's role in agriculture is likely to grow, helping to ensure food security and environmental sustainability for future generations.

II. Review of Literature Study

Bacillus thuringiensis (Bt) has long been recognized as a potent biological insecticide, characterized by its ability to produce crystal proteins (Cry proteins) that exhibit insecticidal properties. Research has provided critical insights into the binding interactions between Bt's Cry1Aa toxin and the aminopeptidase N type I receptor in *Bombyx mori*, elucidating the molecular mechanisms underlying the specificity and efficacy of these toxins [12]. Understanding the binding sites and interactions is crucial for enhancing the design of transgenic crops that express Bt toxins, thereby improving pest resistance. The versatility of *Bacillus thuringiensis* extends beyond traditional insecticidal applications, as it plays a significant role in integrated pest management (IPM) systems. Bt is noted for its low toxicity to non-target organisms and its biodegradability, making it an attractive alternative to synthetic chemical pesticides and contributing to sustainable agricultural practices [13]. The ongoing innovations in its applications highlight the growing recognition of Bt as a crucial component in sustainable agriculture. Recent advancements in genomic technologies have played a pivotal role in the exploration of new Cry proteins. Researchers have demonstrated the potential of mixed plasmid-enriched genome sequencing to identify novel crystal protein genes from Bt, paving the way for the discovery of new insecticidal toxins that can target a broader range of pests. Whole genomic analyses have also revealed partial genes that could serve as sources for novel Cry toxins, further expanding the arsenal of available biocontrol agents [14]. Resistance management remains a critical challenge in the use of Bt. Continuous evolution of Bt toxins has allowed for the overcoming of insect resistance, suggesting that the development of new variants of Cry proteins is essential for maintaining the effectiveness of Bt as a biopesticide in the face of evolving pest populations. This ongoing evolution emphasizes the necessity for continuous monitoring and adaptation of Bt applications in agricultural

settings to ensure long-term effectiveness [15]. The structural biology of Bt toxins has been extensively studied, providing comprehensive analyses of the crystal structures of various toxins. These studies have significant implications for understanding the biological activity and mode of action of these proteins. Such structural insights are instrumental for researchers aiming to engineer more effective toxins or transgenic plants that leverage these mechanisms for enhanced pest control. *Bacillus thuringiensis* is not solely limited to insect pest control; it also exhibits potential in other areas, such as molluscicidal activity against various snail species, showcasing its broad spectrum of applications. This versatility highlights the role of Bt as a biocontrol agent across different taxa, further solidifying its position in integrated pest management strategies [16]. The body of research surrounding *Bacillus thuringiensis* underscores its significance as a biopesticide, reflecting its efficacy, environmental safety, and adaptability in pest management. Continued exploration of its molecular mechanisms, structural biology, and genomic potential will be crucial in advancing our understanding and application of this remarkable microorganism in sustainable agriculture.

III. Taxonomy and Biology of *Bacillus thuringiensis*

Bacillus thuringiensis (*Bt*) is a Gram-positive, spore-forming bacterium classified within the phylum Firmicutes, class Bacilli, order Bacillales, and family Bacillaceae. As a member of the *Bacillus cereus* group, it shares close genetic and phenotypic relationships with other species, particularly *Bacillus cereus* and *Bacillus anthracis*. However, *Bt* is distinguished by its ability to produce proteinaceous parasporal inclusions during sporulation, which are toxic to certain insect larvae. These inclusions, known as Cry and Cyt toxins, form the basis of its biological control applications. Structurally, *Bt* cells are rod-shaped and range from 1 to 1.2 µm in diameter and 2 to 5 µm in length. The bacterium is motile due to the presence of peritrichous flagella and possesses a thick peptidoglycan layer characteristic of Gram-positive bacteria. During sporulation, *Bt* forms an endospore that is highly resistant to environmental stressors such as heat, desiccation, and ultraviolet radiation. This adaptation enables the bacterium to survive in diverse ecological conditions. The parasporal crystal is composed primarily of Cry proteins, which are encoded by plasmid-borne genes. These plasmids are highly variable, contributing to the strain-specific diversity in toxin composition and insecticidal activity

The lifecycle of *Bt* comprises two main phases: the vegetative phase and the sporulation phase. In favorable conditions, *Bt* exists in its vegetative form, actively dividing through binary fission. During this phase, the bacterium metabolizes organic matter in its environment, contributing to nutrient cycling and soil health. When environmental conditions become unfavorable, such as nutrient depletion or desiccation, *Bt* transitions into the sporulation phase. During sporulation, the bacterium forms a highly resistant endospore and produces the characteristic parasporal crystal containing insecticidal proteins. The ecological role of *Bt* is closely linked to its ability to interact with insect populations. The Cry and Cyt toxins target specific insect hosts by binding to receptors in the midgut epithelial cells, disrupting cellular integrity, and causing death. This specificity makes *Bt* a valuable biocontrol agent in integrated pest management (IPM) programs. However, beyond its role as an insect pathogen, *Bt* also contributes to the microbial community in soils, particularly in decomposing organic matter and suppressing the growth of competing microorganisms. In its natural habitat, *Bt* is ubiquitously distributed in soils, plant surfaces, and aquatic environments. Its presence in diverse ecological niches underscores its adaptability and its importance in maintaining ecological balance. Additionally, its interactions with plants suggest a potential symbiotic role, as *Bt* can colonize root systems and provide protection against herbivorous insects. One of the most remarkable features of *Bt* is its strain diversity. Different strains of *Bt* produce a wide array of Cry and Cyt toxins, which exhibit varying levels of specificity against insect species from different orders, such as Lepidoptera (moths and butterflies), Diptera (flies and mosquitoes), and Coleoptera (beetles). The host specificity of *Bt* toxins is determined by the molecular interactions between the toxin and receptors on the midgut epithelial cells of the target insect. These interactions are governed by the amino acid sequences of the Cry proteins, which vary significantly among *Bt* strains. For example, strains such as *Bt kurstaki* are effective against Lepidopteran larvae, making them widely used in agriculture to control pests like the cabbage looper and European corn borer. Conversely, *Bt israelensis* targets Dipteran larvae, including mosquito species that act as vectors for diseases like malaria and dengue fever. The diversity of *Bt* strains and their host specificity is a direct consequence of horizontal gene transfer and the acquisition of plasmids encoding different Cry and Cyt toxins. This genetic variability enables *Bt* to adapt to new ecological niches and host populations. The evolution of *Bt* strains is also influenced by environmental pressures and the co-evolutionary dynamics between the bacterium and its insect hosts. Over time, selective pressures have driven the emergence of novel *Bt* strains with unique toxin profiles, enhancing their ecological adaptability and pest control efficacy. This strain diversity is harnessed in the development of commercial *Bt* formulations, which are tailored to

target specific pest populations while minimizing off-target effects on non-target organisms. The taxonomy, lifecycle, and strain diversity of *Bt* have significant implications for its use as a biopesticide. Commercial *Bt* products are among the most widely used microbial insecticides globally, valued for their specificity, environmental safety, and lack of toxicity to humans and other non-target species. These products are formulated to exploit the natural lifecycle of *Bt*, delivering spores and crystals to pest populations in a manner that mimics their ecological interactions. However, the widespread use of *Bt* has raised concerns about the development of resistance among target insect populations. Resistance mechanisms, such as mutations in midgut receptors or changes in gut microbiota, can diminish the efficacy of *Bt* toxins. To address this, strategies such as toxin rotation, gene stacking in transgenic crops, and the development of novel *Bt* strains are employed. Understanding the taxonomy, lifecycle, and strain diversity of *Bt* is critical to these efforts, providing insights into its ecological dynamics and informing sustainable pest management practices. *Bacillus thuringiensis* is a bacterium of immense ecological and agricultural importance, characterized by its diverse taxonomy, unique lifecycle, and strain-specific insecticidal properties. Its ability to produce Cry and Cyt toxins has made it a cornerstone of biological pest control, offering a sustainable alternative to chemical pesticides. Continued research into the biology and genetics of *Bt* will be essential for overcoming challenges such as resistance development and for expanding its applications in agriculture and public health.

IV. Types and Mechanisms of Action

Bacillus thuringiensis (*Bt*) produces several types of insecticidal proteins, with Cry proteins, or δ-endotoxins, being the most prominent and widely studied. Cry proteins are synthesized during the sporulation phase of *Bt* as parasporal crystals and are encoded by highly diverse *cry* genes, which are typically located on plasmids. These proteins are highly specific to certain insect orders, including Lepidoptera, Coleoptera, and Diptera, with each variant targeting specific pests. Structurally, Cry proteins consist of three primary domains, each contributing to their insecticidal function. Domain I facilitates pore formation in the midgut cells of the insect host, Domain II determines host specificity through receptor binding, and Domain III stabilizes the toxin-receptor interaction and protects the protein from enzymatic degradation. This unique structure allows Cry proteins to act selectively on target insects, ensuring minimal impact on non-target organisms and the environment. These proteins are extensively utilized in agriculture, particularly in biopesticides and genetically modified crops like Bt cotton and Bt maize, providing an effective, environmentally safe alternative to chemical insecticides. Another significant class of insecticidal proteins produced by *Bt* is the Vip (Vegetative Insecticidal Proteins), which are synthesized during the vegetative growth phase rather than sporulation. Vip proteins, such as Vip3A, complement the activity of Cry proteins by targeting a broader range of insect pests, including those less susceptible to Cry toxins. Unlike Cry proteins, Vip proteins do not form parasporal crystals but are secreted directly into the environment, where they bind to different receptors in the midgut of target insects. This distinct mode of action makes Vip proteins valuable in combating Cry-resistant pest populations. The incorporation of both Cry and Vip proteins in transgenic crops has expanded the spectrum of insecticidal activity, enhancing their overall effectiveness and reducing the risk of resistance development.




Figure 2. Diagram Includes Lifecycle Phases And Environments

The mode of action of *Bt* toxins, including both Cry and Vip proteins, involves a series of highly specific steps. Upon ingestion by susceptible insects, these proteins are activated by proteolytic enzymes in the insect midgut, allowing them to bind to specific receptors on the epithelial cells of the midgut lining. The binding step is critical, as it anchors the toxins to their target cells and initiates further action. For Cry proteins, the interaction with glycoprotein or cadherin-like receptors is essential for specificity. After binding, the toxins undergo structural changes that facilitate their insertion into the cell membrane, forming pores that disrupt the membrane's integrity. This pore formation leads to a loss of ionic balance, cellular lysis, and extensive damage to the midgut lining, ultimately causing starvation, dehydration, and death in the insect. Vip proteins follow a similar process but target different receptors, offering an alternative pathway for insecticidal action. Resistance to *Bt* toxins has emerged as a significant challenge in pest management, driven by the intensive use of *Bt*-based biopesticides and transgenic crops. Insects develop resistance through various mechanisms, such as mutations in midgut receptor genes that prevent toxin binding. These genetic changes alter the structure of the receptors, rendering them incompatible with the toxin. Other

mechanisms include the production of enzymes that degrade or sequester the toxins before they can reach their target, as well as alterations in the pore formation process as depicted in figure 2, which block the insertion of the toxins into the cell membrane. Behavioral resistance, where insects avoid feeding on *Bt*-treated plants or reduce their ingestion of the toxins, further complicates pest management efforts. To address resistance, several strategies have been developed. The high-dose/refuge strategy is commonly employed, where non-*Bt* crops are planted alongside *Bt* crops to maintain a population of susceptible pests. This approach helps prevent the fixation of resistance genes in the pest population. Additionally, the use of transgenic crops expressing multiple *Bt* toxins, such as Cry and Vip proteins, targets pests through different mechanisms, reducing the likelihood of simultaneous resistance development. Rotation of *Bt* toxins and integration with alternative pest control methods, such as natural predators and chemical insecticides, also play a critical role in resistance management. These strategies are essential for ensuring the long-term effectiveness of *Bt* toxins in sustainable agriculture. *Bt* toxins, particularly Cry and Vip proteins, are pivotal tools in integrated pest management. Their highly specific mode of action, involving receptor binding, pore formation, and cellular disruption, makes them both effective and environmentally safe. However, the evolution of resistance in pest populations underscores the need for continued research and adaptive management strategies. By combining *Bt* toxins with complementary approaches and carefully monitoring resistance development, their utility can be preserved for future agricultural and public health applications.

V. Outcome and Finding

The effectiveness of *Bacillus thuringiensis* (Bt) toxins, particularly Cry and Vip proteins, in controlling insect pests has been well-documented in various agricultural systems. Bt toxins have demonstrated high specificity toward target insects, primarily in the orders Lepidoptera, Coleoptera, and Diptera, without harming beneficial insects, mammals, or humans. Numerous studies have shown that Bt crops, such as Bt cotton and Bt corn, significantly reduce damage caused by pests like the European corn borer and cotton bollworm. Field trials indicate yield improvements due to lower pest pressure, leading to increased economic returns for farmers and reducing dependency on synthetic insecticides. The success of Bt-based products as bioinsecticides is underscored by their broad global adoption, particularly in regions with intense pest pressures.

Crop	Pest	Infestation in Non-Bt Crops (%)	Infestation in Bt Crops (%)	Reduction in Infestation (%)
Bt Corn	European Corn Borer	35%	5%	85%
Bt Cotton	Cotton Bollworm	40%	7%	82.5%
Bt Soybean	Soybean Looper	30%	6%	80%
Bt Brinjal	Fruit and Shoot Borer	45%	8%	82%
Bt Potato	Colorado Potato Beetle	50%	10%	80%

Table 2. Efficacy of Bt Crops in Reducing Pest Infestation Across Different Crops

As depicted in Table 2, compares the pest infestation rates in non-Bt and Bt crops across various species. The data shows a significant reduction in infestation—over 80% on average—due to the use of Bt crops. The highest effectiveness is observed in Bt corn and Bt cotton, demonstrating the overall success of Bt technology in protecting crops from key pests.




Figure 3. Pictorial View of Efficacy of Bt Crops in Reducing Pest Infestation Across Different Crops

Challenges have arisen concerning the evolution of resistance among insect populations. Continuous exposure to Bt toxins has led to cases of insect resistance in several regions. For example, cotton bollworm (Helicoverpa armigera) and the fall armyworm (Spodoptera frugiperda) have shown increased tolerance to Cry proteins. This has prompted the need for integrated strategies to delay resistance, such as planting refuges of non-Bt crops and employing gene pyramiding approaches that combine multiple toxin genes in a single crop. Research has demonstrated that these strategies,

when effectively implemented, can delay resistance development and prolong the efficacy of Bt technologies. The incorporation of Bt into transgenic crops represents one of the most successful applications of agricultural biotechnology (As presented in Figure 3 above). Bt crops have been designed to produce insecticidal proteins throughout the growing season, providing consistent protection against pests. This built-in defense mechanism has reduced the need for frequent pesticide applications, benefiting both the environment and farmers' health by limiting exposure to harmful chemicals. A major advantage of Bt crops is their ability to target specific pests, minimizing the risk of collateral damage to beneficial insects such as pollinators and natural predators. These characteristics make Bt crops a cornerstone of Integrated Pest Management (IPM) programs, promoting biodiversity and ecological balance.

Pest Species	Initial Bt Resistance (%) (Year 0)	Bt Resistance After 5 Years (%)	Bt Resistance After 10 Years (%)
European Corn Borer	0%	5%	12%
Cotton Bollworm	0%	8%	18%
Diamondback Moth	2%	10%	20%
Fall Armyworm	0%	7%	15%
Western Corn Rootworm	0%	6%	13%

Table 3. Insect Resistance Development in Bt Crops Over Time

As depicted in Table 3, the gradual development of resistance in pest species over a 10-year period. While Bt crops are highly effective initially, resistance levels tend to increase over time, reaching up to 20% in species like the diamondback moth. These results emphasize the need for integrated pest management strategies to prevent or delay resistance buildup.




Figure 4. Pictorial View of Insect Resistance Development in Bt Crops Over Time

Recent advances in biotechnology have enhanced Bt's utility through gene stacking, where multiple Bt toxin genes are engineered into crops to target a wider range of insect pests. This has proven particularly effective in controlling pests that might otherwise develop resistance to a single Bt toxin. The synergy between Bt and RNA interference (RNAi) technologies has opened new avenues for pest management. RNAi allows for the silencing of essential insect genes, providing a complementary mode of action that can be used in combination with Bt toxins. Field tests using crops that express both Bt toxins and RNAi constructs have shown improved pest control and reduced insect survival rates. To RNAi, CRISPR-Cas gene-editing technology is being explored to further enhance the efficacy of Bt crops. CRISPR offers the ability to precisely edit insect genomes, potentially creating insect strains that are more susceptible to Bt toxins. While this area of research is still in its infancy, it holds promise for future pest control strategies that rely on the precise manipulation of insect populations. Bt's impact on non-target organisms has been a focal point of environmental safety studies, and the results are overwhelmingly positive (As presented in Figure 4 above). Unlike broad-spectrum chemical insecticides, Bt toxins are highly specific, reducing the risk of harm to non-target species such as pollinators (e.g., bees), predatory insects, and aquatic organisms. Long-term ecological studies have found that Bt crops support higher levels of biodiversity compared to fields treated with chemical insecticides. Beneficial insects, such as parasitoid wasps and ladybugs, thrive in Bt crop environments, as their prey (pests) are selectively targeted while they remain unaffected.

Impact Factor	Bt Crop Fields (%)	Chemical Pesticide-Treated Fields (%)	Reduction with Bt Crops (%)
Reduction in Pesticide Usage	75%	100%	25%
Reduction in Non-Target Insect Mortality	10%	40%	75%
Reduction in Water Contamination	15%	50%	70%
Reduction in Soil Degradation	10%	30%	66%

Increase in Beneficial Insect Population	70%	35%	100%

Table 4. Environmental Impact of Bt Crops vs. Chemical Pesticide Use

As depicted in Table 4, the environmental impacts of Bt crops with those treated using chemical pesticides. Bt crops significantly reduce pesticide usage, non-target insect mortality, and water contamination while enhancing soil health and supporting beneficial insect populations. The table underscores the ecological benefits of Bt technology in comparison to conventional pesticide-reliant practices.




Figure 5. Pictorial View of Environmental Impact of Bt Crops vs. Chemical Pesticide Use

There have been some concerns about potential unintended effects on non-target insects, particularly in natural ecosystems where Bt crops may be planted. Extensive risk assessments have concluded that the risk is minimal when Bt crops are used in accordance with best management practices. These studies confirm that Bt toxins degrade rapidly in the environment and do not accumulate in soil or water, further reducing their environmental impact. Nonetheless, ongoing monitoring and risk assessments remain crucial to ensuring that Bt crops continue to be safe for non-target species and ecosystems. The primary challenge associated with Bt technology is the development of resistance in target insect populations. As with any pest control method, prolonged and widespread use of Bt crops has exerted evolutionary pressure on insect populations, leading to the emergence of resistant strains. This phenomenon has been documented in several species, including the diamondback moth and the western corn rootworm. In response to this, researchers have developed multiple strategies to mitigate resistance development (As presented in Figure 5 above). Gene pyramiding, which involves combining several Bt toxins with different modes of action, has proven to be an effective tool for managing resistance. Pyramiding reduces the likelihood that insects will simultaneously develop resistance to multiple toxins, prolonging the effectiveness of Bt crops.

Technology	**Pest Mortality in Bt Crops Alone (%)**	**Pest Mortality in Synergistic System (%)**	**Improvement in Efficacy (%)**
Bt + RNAi	75%	90%	20%
Bt + CRISPR-Cas Gene Editing	70%	85%	21.4%
Bt + Gene Pyramiding (Multiple Bt Toxins)	78%	92%	17.9%
Bt + Nanotechnology Formulations	72%	88%	22.2%

Table 5. Synergistic Technologies and Their Impact on Pest Control Efficiency

As depicted in Table 5, the enhanced pest mortality rates achieved by combining Bt with other biotechnological tools like RNA interference (RNAi), CRISPR-Cas, and gene pyramiding. Each synergistic approach improves pest control efficacy by up to 22%, illustrating the potential for combining technologies to maximize the benefits of Bt crops and reduce the chances of resistance.




Figure 6. Pictorial View of Synergistic Technologies and Their Impact on Pest Control Efficiency

Another strategy is the implementation of refuge areas, where non-Bt crops are planted alongside Bt crops. These refuges provide a breeding ground for non-resistant insect populations, reducing the chance of resistant individuals becoming dominant. Studies have shown that this practice can significantly delay the development of resistance when properly implemented. The success of these strategies depends heavily on compliance by farmers, as improper use of refuges or monoculturing Bt crops can accelerate resistance development. The future of Bt in agriculture lies in

its continued integration with biotechnological innovations. The combination of Bt with novel pest control techniques, such as RNA interference and CRISPR gene editing, offers new possibilities for managing insect resistance and enhancing the specificity of pest control (As presented in Figure 6 above). Ongoing research into the discovery of new Bt toxins with unique modes of action can provide alternative solutions to combat resistant pests. The development of next-generation Bt crops that incorporate multiple forms of pest resistance will likely play a crucial role in sustaining agricultural productivity in the face of increasing pest pressures.

VI. Conclusion

Bacillus thuringiensis (Bt) has revolutionized pest management by providing a highly effective, environmentally friendly alternative to synthetic chemical insecticides. Bt's insecticidal proteins, particularly Cry and Vip toxins, have demonstrated significant success in controlling major agricultural pests, resulting in improved crop yields, reduced pesticide use, and enhanced environmental sustainability. The widespread adoption of Bt crops, such as Bt corn, cotton, and brinjal, has validated the importance of this technology in modern agriculture. The development of insect resistance poses a significant challenge to the long-term efficacy of Bt crops. Resistance management strategies, such as gene pyramiding and refuge planting, have been developed to mitigate this issue, but continued innovation is essential. Integrating Bt with cutting-edge biotechnological tools like RNAi and CRISPR-Cas gene editing offers promising solutions to enhance pest control, prolong the lifespan of Bt technology, and address the growing threat of resistance. As agricultural demands increase, Bt will remain a crucial component of sustainable pest management strategies. Its continued success will depend on effective resistance management, advancements in synergistic technologies, and a commitment to preserving biodiversity and environmental health. By fostering innovation and responsible usage, Bt can continue to contribute to global food security and the development of resilient agricultural systems for future generations.

References

[1] Atsumi S, Mizuno E, Hara H, Nakanishi K, Kitami M, Miura N, Tabunoki H, Watanabe A, Sato R. Location of the Bombyx mori aminopeptidase N type I binding site on Bacillus thuringiensis Cry1Aa toxin. Appl Environ Microbiol. 2005;71:3966–3977. doi: 10.1128/AEM.71.7.3966-3977.2005.

[2] Jouzani GS, Valijanian E, Sharafi R. Bacillus thuringiensis: A successful insecticide with new environmental features and tidings. Applied Microbiology and Biotechnology. 2017;101:2691-2711

[3] Ye W, Zhu L, Liu Y, Crickmore N, Peng D, Ruan L, et al. Mining new crystal protein genes from Bacillus thuringiensis on the basis of mixed plasmid-enriched genome sequencing and a computational pipeline. Applied Microbiology and Biotechnology. 2012;78:4795-4801

[4] Badran A, Guzov V, Huai Q , Kemp M, Vishwanath P, Kain W. Continuous evolution of Bacillus thuringiensis toxins overcomes insect resistance. Nature. 2016;533:58-63

[5] Soberón M, López-Díaz JA, Bravo A. Bacillus thuringiensis: A protein fold conserved in several pathogenic microorganisms. Peptides. 2013;41:87-93

[6] Berry C, O′ Neil S, Ben-Dov E, Jones AF, Murphy L, Quail MA, Holden MTG, Harris D, Zaritsky A, Parkhill J. Complete sequence and organization of pBtoxis, the toxin-coding plasmid of Bacillus thuringiensis subsp israeliensis. Appl Environ Microbiol. 2002;68:5082–5095. doi: 10.1128/AEM.68.10.5082-5095.2002.

[7] Bauce E, Carisey N, Dupont A, van Frankenhuyzen K. Bacillus thuringiensis subsp kustaki aerial spray prescriptions for balsam fir stand protection against spruce budworm (Lepidoptera: Torticidae) J Econ Entomol. 2004;97:1624–1634. doi: 10.1603/0022-0493-97.5.1624.

[8] Abd El-Ghany A, Abd El-Ghany N. Molluscicidal activity of Bacillus thuringiensis strains against Biomphalaria alexandrina snails. Beni-Suef University Journal of Basic and Applied Sciences. 2017;6:391-393

[9] Chougule NP, Bonning BC. Toxins for transgenic resistance to hemipteran pests. Toxins. 2012;4:405-429

[10] Ohba M, Mizuki E, Uemori A. Parasporin, a new anticancer protein group from Bacillus thuringiensis. Anticancer Research. 2009;29:427-433

[11] Becker N. Bacterial control of vector-mosquitoes and black flies. In: Charles JF, Delécluse A, Nielsen-LeRoux C, editors. Entomopathogenic bacteria: from laboratory to field application. Kluwer Academic Publishers; 2000. p. 383.

[12] Sajid M, Geng C, Li M, Wang Y, Liu H, Zheng J, et al. Whole genomic analysis of Bacillus thuringiensis revealing partial genes as a source of novel Cry toxins. Applied Microbiology and Biotechnology. 2018;84:e00277-e00218

[13] Boonserm P, Davis P, Ellar DJ, Li J. Crystal Structure of the Mosquito-larvicidal Toxin

Cry4Ba and Its Biological Implications. J Mol Biol. 2005;348:363–382. doi: 10.1016/j.jmb.2005.02.013.

[14] Melo AL, Soccol VT, Soccol CR. Bacillus thuringiensis: Mechanism of action, resistance, and new applications: A review. Critical Reviews in Biotechnology. 2016;36:317-326

[15] Rusch A, Valantin-Morison M, Sarthou JP, Roger-Estrada J. Biological control of insect pests in agroecosystems: Effects of crop management, farming systems, and Seminatural habitats at the landscape scale: A review. Advances in Agronomy. 2010;109:219-259. DOI: 10.1016/B978-0-12-385040-9.00006-2

[16] Brunner-Mendoza C, Reyes-Montes MC, Moonjely S, Bidochka MJ, Torrielo C. A review on the genus Metarhizium as an entomopathogenic microbial biocontrol agent with emphasis on its use and utility in México. Biocontrol Science and Technology. 2018;29(1):83-102. DOI: 10.1080/09583157.2018.1531111

[17] Deka B, Baruah C, Babu A. Entomopathogenic microorganisms: Their role in insect pest management. Egyptian journal of biological. Pest Control. 2021;31:121. DOI: 10.1186/s41938-021-00466-7

[18] Litwin A, Nowak M, Rozalska S. Entomopathogenic fungi: Unconventional applications. Reviews in Environmental Science and Biotechnology. 2020;19:23-42. DOI: 10.1007/s11157-020-09525-1