18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Biological Control of Agricultural Pests: Innovations in RNA Interference and Transgenic Approaches for Pest Suppression

Varun kumar Sharma*[1], Saravanan M[2], Preethi Murali[3], Sridevi Sangeetha KS[4], Divya N[5], Uma Bhardwaj[6]

[1] Associate Professor , Biotechnology and Microbiology , Noida international University , Greater Noida, Uttar Pradesh ,India (Varun kumar Sharma) | Email: varun1.sharma@niu.edu.in

[2] Central Research Laboratory, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research (Saravanan M) | ORCID: 0000-0002-1696-9532| Email: saravanan@maher.ac.in

[3] Department of Oral Pathology, Meenakshi Ammal Dental College and Hospital, Meenakshi Academy of Higher Education and Research (Preethi Murali) | ORCID: 0000-0002-7483-1873 | Email: pmurali@maher.ac.in

[4] Meenakshi College of Allied Health Sciences, Meenakshi Academy of Higher Education and Research (Sridevi Sangeetha KS) | ORCID: 000-0002-8175-1484 | Email: ssks@maher.ac.in

[5] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Divya N) | ORCID: 0000-0002-5620-6820 | Email: divyan@maher.ac.in

[6] Professor, Biotechnology and Microbiology , Noida international University , Greater Noida, Uttar Pradesh ,India (Uma Bhardwaj) | Email: vc@niu.edu.in

Abstract: The growing use of chemical insecticides to control pests in agriculture has caused a lot of worry about the long-term health of people and the environment, as well as the ability of pests to become resistant to these chemicals. As a result, new biological control methods are being developed. RNA interference (RNAi) and transgenic techniques are at the head of these methods for getting rid of pests. RNAi is a natural process in cells that controls gene translation. It is an exact and effective way to get rid of pest populations. Scientists can effectively get rid of pests while having little to no effect on other living things by creating small interfering RNAs (siRNAs) that target specific genes in pest species. This stops important biological processes like growth, reproduction, and survival. New biopesticides have been made possible by recent progress in RNAi technologies. These can be used directly on crops or by changing plants. The use of RNAi to control pests not only cuts down on the need for traditional poisons, but it also reduces the damage that farming does to the environment. Using RNAi along with modified crops also has positive benefits that work together to make crops more resistant to pests and lower yield losses. Pest-resistant genes are added to crop genomes using transgenic methods. This gives crops features like producing Bt (Bacillus thuringiensis) poison or being resistant to insects that eat plants. Genetically modified organisms (GMOs) like these can greatly lower the number of pests while keeping crops productive. People often don't want modified crops to be used, though, because they're worried about biodiversity, environmental balance, and possible side effects that weren't meant to happen. This paper talks about the newest RNAi and transgenic technologies for controlling pests, focused on how they work, how well they work, and what effects they have on the environment.

Keywords: RNA Interference, Transgenic Crops, Pest Suppression, Biopesticides, Sustainable Agriculture, Genetic Engineering

I. Introduction

To meet the growing global food demand, which is expected to hit over 9 billion people by 2050, the farming industry is facing problems that have never been seen before. The spread of farm pests is one of the biggest threats to food security because they can seriously lower crop output and make it hard for people to make a living. Chemical insecticides have been the mainstay of traditional pest control methods. These chemicals work in the short term, but they have caused many health and environmental problems. Some of these are damage to the environment, the development of bug tolerance, and bad effects on species that aren't the goal, like humans and helpful insects [1]. To deal

***Corresponding Author:**
Received: 24th April, 2026; Revised: 11th May, 2026; Accepted: 24th May, 2026; Available Online: 6th June, 2026
(DOI): 10.70102/AEJ.2026.18.1s.85

with these problems, new biological control methods are being created. RNA interference (RNAi) and transgenic approaches are looking like good options for long-term pest control. RNA interference is a normal biological process that controls how genes are expressed in living cells. This process can be used to create focused pest control plans that hurt non-target species as little as possible. Scientists can successfully mess up important biological processes by creating small interfering RNAs (siRNAs) that target genes that are needed for pests to survive. RNAi has a lot of ability to get rid of many types of farming pests, such as insects and worms. Because it is so precise, it can kill specific pests that regular poisons can't. This lowers the risks that come with using chemicals. New advances in RNAi technology have made it easier to create new biopesticides that can be used in a variety of farming situations [2].

These biopesticides can be put on crops directly or given to plants through plant transformation, which makes plants make their own siRNAs that kill pests. These kinds of innovations can keep the ecosystem balanced while reducing the number of pests and food losses by a large amount. Also, the fact that RNAi can work against more than one species by changing the pattern of the siRNA opens up new ways to get rid of pests that are successful and good for the environment. Along with RNAi, genetic methods are becoming more popular in the fight against pests. For these ways to work, foods must be genetically modified to make them resistant to certain pests [3]. Putting Bacillus thuringiensis (Bt) genes into crops is one of the most well-known cases. Some bug pests are killed by toxins that Bt crops make. This successfully controls their numbers while greatly reducing the need for chemical poisons. The economic success of Bt crops like corn and cotton has led to big increases in output and decreases in the use of pesticides, showing that genetic methods work. But there has been some disagreement about the use of modified crops [4]. People are worried about species, the environment, and the moral issues that come up with genetic editing. These worries have led to discussions that often stop these technologies from being widely used.

Moreover, the improvement of bother resistance to Bt crops has made individuals ponder on the off chance that hereditary bother administration strategies will work within the long run. Since of this, combining RNAi and hereditary methods for bug control ought to be looked at with a reasonable see, taking under consideration both the stars and cons. Utilizing both RNAi and hereditary strategies together makes a interesting chance to progress strategies for controlling bugs. Researchers can make crops that are both hereditarily changed to be safe to bothers conjointly able to utilize RNAi to way better target and control bug populaces by combining these innovations [5]. This method that employments more than one way to induce freed of bothers can offer assistance ease a few of the stresses that come with depending on fair one, making the farming community more grounded. In expansion, the rules that administer hereditary propels in cultivating ought to alter to keep up with these advancements. To create beyond any doubt that the inquire about and utilize of RNAi and altered nourishments are secure, it is imperative to set up clear and successful administering frameworks [6]. This incorporates both exhaustive security tests and open contact with the open to induce them to believe and acknowledge these innovations. Counting agriculturists, clients, and legislators within the discussion almost biotechnology advancements is imperative for dealing with concerns and making beyond any doubt that these arrangements are utilized in a capable way.

II. Related Work

RNA obstructions (RNAi) has been examined increasingly as a conceivable way to urge freed of bugs in later a long time, with numerous ponders appearing that it might work. For case, researchers have been able to utilize RNAi to target particular qualities in imperative cultivating bothers, like the cotton bollworm (Helicoverpa armigera), which is known for harming cotton areas. Considers have appeared that giving these pests siRNA that targets imperative qualities caused tall passing rates and issues with growth, appearing that RNAi is an compelling way to urge freed of bugs. Moreover, advance in transport frameworks, like capsule strategies and nanoparticle shapes, have made it less demanding to make RNAi-based biopesticides [7]. These frameworks make siRNAs more steady and successful when connected to crops. At the same time, mechanical biotechnology has been centering on how to utilize adjusted strategies together. The improvement of Bt crops has changed the way bothers are controlled, giving a long-term choice to chemical harms. Many field tests and industrial uses of Bt cotton and Bt corn have shown that they lower bug numbers, boost output, and have minimal negative effects on the environment. According to research, using Bt crops can also have other benefits, like lowering the need for pesticides and helping farms make more money. But because some bug populations have become resistant, scientists are still looking into ways to stop resistance from spreading. Some ideas being looked into are shelter methods and using RNAi along with genetic traits [8]. Recently, researchers have also looked into how RNAi and genetic methods might work together to make next-generation foods that are immune to pests. For instance, scientists have created transgenic plants that carry RNAi constructs that target multiple genes in pests. This makes the plants

much more resistant to pests. This two-pronged method not only lowers the chance of developing resistance, but it also increases the types of pests that can be managed, making the problem of pest management more complete. On top of that, the rules that govern RNAi and genetic technologies are changing. Many countries are reviewing their rules to make room for these new technologies [9].

Table 1: Summary of Related Work

Aspect	Related Work	Future Trend	Challenges	Benefits
RNAi Applications	Successful use in cotton bollworm and fall armyworm	Expanded use in diverse crops	Target specificity in diverse pest species	High specificity reduces harm to non-target organisms
Transgenic Crops	Development of Bt crops for pest resistance	Increased stacking of resistance traits	Public acceptance and regulatory hurdles	Effective pest control with reduced chemical use
Integration [10]	Combining RNAi with traditional pest management	Development of multi-faceted IPM strategies	Resistance development in pests	Enhanced pest suppression through synergistic effects
Delivery Methods	Advancements in nanoparticle delivery systems	Innovations in plant-based RNAi production	Ensuring efficient and stable delivery to pests	Potential for reduced pesticide applications
Environmental Impact	Studies on non-target effects of RNAi	Improved assessments of ecological impacts	Long-term ecological monitoring	Improved ecosystem health and biodiversity
Economic Analysis [11]	Cost-benefit analyses of Bt crops	Economic evaluations of integrated approaches	Initial costs of technology adoption	Increased profitability for farmers
Crop Diversity	Research on diverse pest-resistant crops	Breeding programs for resilience	Limited genetic diversity in commercial crops	Enhanced resilience to pests and environmental stress
Education & Outreach	Awareness campaigns on biotechnology benefits	Greater focus on farmer education and training	Misinformation about GMOs and RNAi	Informed decision-making and acceptance
Regulatory Framework [12]	Evolving regulations on GMOs	Streamlined approval processes for new technologies	Complex regulatory landscapes	Faster adoption of effective pest management solutions
Research Collaboration	Collaborative projects across institutions	Global partnerships for technology transfer	Coordination among stakeholders	Accelerated research and development
Biocontrol Integration	Incorporation of natural enemies in IPM	Enhanced synergistic approaches	Balancing biocontrol with transgenic methods	More holistic pest management strategies
Market Demand	Increasing demand for sustainable practices	Growth in organic and non-GMO markets	Consumer preferences for non-GMO products	Greater market access and competitiveness
Climate Resilience [13]	Research on pest responses to climate change	Development of climate-resilient pest control strategies	Adapting to shifting pest dynamics	Improved food security under changing climates
Long-term Sustainability	Studies on long-term effectiveness of technologies	Focus on sustainable agricultural practices	Ensuring ongoing effectiveness of technologies	Sustainable food systems and ecosystem balance

III. Overview of Biological Control

A. Definition and principles of biological control

Biological control is the use of predators, parasitoids, viruses, or competitors, which are live things, to get rid of pests and diseases in farms. This method is based on ecological principles and uses

the natural interactions between living things to control pest numbers, which means that chemical poisons are not needed as much. Biological control tries to make a way to get rid of pests that is safe and good for the environment, which helps keep the ecosystem in balance. Biological control is based on knowing how bug populations change over time and how their natural enemies affect them [14]. One important idea is "natural regulation," which means that predators or parasitoids help keep the number of pests below levels that are harmful. Adding lady bugs to control aphid numbers is one example of this concept in action. Through using these beetles' natural ability to eat other bugs, farms can successfully lower the number of aphids without using chemicals. In biological control, there are three main groups: traditional, augmentative, and conservation biological control. To get rid of invading pest species, classical biological control includes bringing in natural enemies from the bug's home range. A good example of this method being used successfully in agriculture is the introduction of the Vedalia bug to control cottony cushion scale in California. Augmentative biological control means letting out more natural enemies to make their populations bigger and more effective at getting rid of bugs. In order to get the best results, this way is often used along with other methods of pest control, like cultural or mechanical ones [15]. In conservation biological control, the goal is to keep and grow the natural pests that are already in the habitat. To do this, you can change the environment by putting flowering plants or cover crops that give food and protection to helpful creatures and encourage them to keep pests away.

B. Historical context and evolution of biological control methods

The idea of biological control has a long history that goes back to the way people used to farm. There is evidence that the Chinese used hungry insects to control pests in rice fields as early as 2000 BCE. In the same way, the ancient Egyptians knew the value of natural enemies and used cats to keep rodents out of grain stores. However, biological control did not start to take shape as an official science field until the late 1800s and early 1900s. The Vedalia beetle (Rodolia cardinalis) was first brought to California in the 1880s. This was the start of the current age of biological control [16]. The successful fight against the cottony cushion scale (Icerya purchasi) on citrus crops was a big step forward in the field. When the Vedalia beetle was brought in, it successfully lowered the number of pests, which led to a rise in orange production. This event laid the groundwork for future biological control methods and showed how natural enemies could be used to get rid of pests in farms. After that, the field of biological control grew very quickly in the following decades. Scientists started to look into the rules of host specificity, the life cycles of natural foes, and how ecosystems in agroecosystems interact with each other. Classical biological control became a separate method in the early 20th century. It involved introducing natural enemies on purpose to get rid of unwanted pest species.

It was also during this time that basic ideas like the need for careful study of how new species affect the environment came into being. By the middle of the 20th century, we had a much better idea of how pests and natural enemies communicate. Researchers started to work on augmentative biological control methods, in which more natural enemies were allowed to make them more effective at getting rid of bugs [17]. This method became more popular as people learned more about the environmental problems caused by chemical poisons, especially after Rachel Carson's book Silent Spring came out in 1962 and showed how bad these chemicals are for the environment. Biological control has become popular again in the late 20th century and early 21st century. This is because farmers need to use methods that are good for the environment. Because of progress in molecular biology and genetics, scientists are now looking into how biotechnology can be used to control pests. For example, RNA interference and modified foods are two examples of this. Biological control is now seen as an important part of integrated pest management (IPM) plans [18]. This shows how important natural balance and variety are for farming that will last.

C. Current trends in biological control strategies

Within the past few a long time, organic control strategies have gotten to be exceptionally well known as long-term options to utilizing chemicals to urge freed of bothers. This alter is generally caused by developing stresses around harm to the environment, resilience to pesticides, and wellbeing impacts of chemical harms. Organic control has gone through a number of imperative changes over the a long time. These changes are due to changes in innovation, our understanding of environment, and the rules that oversee cultivating. Atomic methods being utilized in organic control is one of the foremost curiously patterns. RNA obstructions (RNAi) innovation has made it possible to target particular qualities in bother populaces, which has opened up other ways to urge freed of bothers. Researchers are working on RNAi-based biopesticides that can halt bothers from doing critical natural errands [19]. This will successfully control the populace whereas having the slightest sum of impact on other living things. Together with more standard natural control strategies, this modern approach is being looked into. Together, they can make plans that work way better generally. Another slant is that organic control for security is getting more consideration. Rather than fair including

unused characteristic foes, there's increasingly center on making the environment and assets way better for animals that as of now do great work [20]. Ranchers are utilizing strategies that offer assistance natural pest control, like developing cover crops, including blossoming plants, and taking after agroecological standards. These activities not as it were offer assistance to ensure distinctive sorts of plants and creatures, but they moreover make cultivating frameworks more grounded, which makes bother control final longer. Natural control that employments supportive microbes is additionally getting to be more well known.




Figure 1: Current trends in biological control strategies

Researchers have found that entomopathogenic fungi, bacteria, and viruses can kill a lot of different types of crop pests. These microbes can be used as biopesticides or added to the way crops are managed now. They are an eco-friendly alternative to chemical pesticides. Researchers are still looking into microbial biocontrol agents, which could lead to long-lasting methods for controlling pests. Biological control methods are also changing the way people accept them and how the government regulates them.

IV. RNA Interference (RNAi) in Pest Control

A. Mechanism of RNAi

1. Role of small interfering RNAs (siRNAs)

The RNA impedances (RNAi) framework is made up of little atoms called siRNAs. These siRNAs play a key part in controlling quality interpretation and ensuring diverse species from infection assaults and transposable components. More often than not, siRNAs are 20 to 25 nucleotides long, little, double-stranded RNA pieces that play a key part in turning off qualities. Since they can precisely target certain courier RNAs (mRNAs), they are exceptionally valuable for consider, medication, and getting freed of bothers. When long double-stranded RNA (dsRNA) is put into a cell, siRNAs begin to do their work. This dsRNA can come from a number of places, such as viral illnesses, adjusted plants, or being utilized on the exterior as a biopesticide. The chemical Dicer breaks down the dsRNA into littler siRNA pieces once it gets interior the cell. RNA particles have two parts: a direct strand and a traveler strand. The direct strand controls the hush prepares. The direct strand is included to the RNA-induced quieting complex (RISC) within the RNAi pathway. This complex is an vital portion of the gene-silencing handle since it makes a difference the siRNA discover and connect to coordinating mRNA particles interior the cell. One of the finest things approximately siRNAs is that they are exceptionally particular. Researchers can make siRNAs that target qualities that are as it were found in certain bugs. This way, the hush impacts as it were happen within the right places and do not harmed other living beings. When the siRNA's direct strand joins to its coordinating mRNA, RISC can either break down the mRNA or halt it from being deciphered into protein. To do this, certain proteins that are required for the desired bother to develop, create, or duplicate are blocked from being communicated. For illustration, siRNAs can be made to target qualities that are included in vital forms like breeding or molting. By interferometer with these capacities, bothers can kick the bucket or their populaces will develop less rapidly. Moreover, siRNAs can be utilized to murder bugs in a number of ways, such as by putting them straightforwardly on areas or including them to hereditarily adjusted plants that make siRNAs that target bug qualities. Compared to traditional toxic bug sprays, this modern strategy is maintainable and secure for the environment, and it makes a difference with coordinates bug control methodologies.

2. Gene silencing processes

Quality hushing may be a wide term for a number of diverse forms that diminish or halt quality generation. We require this handle to keep cellular adjust, control natural exercises, and ensure ourselves from infections and transposable components that attempt to change our qualities. Quality suppression can be worn out numerous ways, but RNA impedances (RNAi) is one of the foremost examined and broadly utilized. Little RNA

particles, generally little interferometer RNAs (siRNAs) and microRNAs (miRNAs), do the work of RNAi. When double-stranded RNA (dsRNA) is put into a cell, the method begins. This dsRNA can come from a number of places, such as viral infections or lab tests where it is presented misleadingly. The protein Dicer finds the dsRNA and cuts it into littler pieces, as a rule 20 to 25 nucleotides long. This makes siRNAs or miRNAs. When it comes to siRNAs, they are portion of the RNA-induced hushing complex (RISC). The siRNA's direct strand ties to mRNA districts that are complementary. This will break down the mRNA or halt it from being deciphered into protein. The RISC cuts the mRNA so it can't be handled on the off chance that the coordinate between the siRNA and mRNA is idealize or nearly idealize. This brings down the sum of protein and quiets the target quality. MiRNAs, on the other hand, come from longer RNA sources and ordinarily go after mRNAs that do not coordinate up flawlessly. When miRNAs connect with RISC, they can halt interpretation by halting the arrangement of ribosomes or speeding up the breakdown of the particular mRNA. This fine-tuned control lets cells alter gene expression in response to changes within the environment and prompts for development. There are other ways to quiet qualities other than RNAi, such as transcriptional quality hushing (TGS). Epigenetic changes, such as DNA methylation and histone adjustment, are common in TGS. These changes cause chromatin to be in a stifled state. This handle stops the generation of certain qualities, which includes another level of control to quality expression.

B. Applications of RNAi in agricultural pest management

1. Targeting specific pest genes

Because it can target specific genes in pest populations, RNA interference (RNAi) has become a new tool for controlling pests in agriculture. Because of this, customized pest control plans can be made that cause the least amount of harm to non-target animals and the environment. This helps fix some of the big problems with standard chemical poisons. Because of how RNAi works, scientists can make small molecules called siRNAs that can target mRNAs of important genes in bugs. A lot of the time, these genes are very important for things like growth, development, reproduction, and staying alive. By making siRNAs that target these important genes, it is possible to stop the biological processes of pest species that are being targeted. For instance, siRNAs can be made to target genes that help insects shed their skin.

2. Case studies demonstrating effectiveness

A lot of case studies show that RNA interference (RNAi) is a successful way to get rid of pests in agriculture, showing that it could be used instead of conventional herbicides in the long term. These studies show that RNAi can target specific genes in pests, which can greatly reduce the number of pests and have little to no effect on organisms that are not the target. The cotton bollworm (Helicoverpa armigera), a big pest that hurts cotton and other crops, was controlled with RNAi in a well-known case study. Scientists made siRNAs that target important genes for an insect's growth and development, especially genes related to the molting process. When these siRNAs were fed to the larvae, there were high death rates; more than 80% of the treated larvae did not make it to the end of their growth. This study showed that RNAi can effectively control pest populations by messing with important biological processes in the pests that are being targeted. Another strong example comes from studying the diamondback moth (Plutella xylostella), which is one of the worst pests for green plants. Scientists created mutant plants that produce dsRNA that targets specific genes that are necessary for the moth to survive. When caterpillars ate these plants, they stopped growing and dying very quickly. Their weight as larvae was also much lower than when they ate non-transgenic plants. This research showed that RNAi can be used in plant breeding projects to help control pests in a way that lasts and makes crops more resistant to damage. In a different case study, RNAi was used to kill the fall armyworm (Spodoptera frugiperda), which is very bad for corn and other crops. Scientists made siRNAs that target genes that are involved in how the insect's gut works. In field tests, plants that were treated with these siRNAs had lower pest damage and higher outputs than plants that were not treated. For example, RNAi's ability to control fall armyworm numbers shows that it could be widely used in many different types of farming.

V. Transgenic Approaches for Pest Suppression

A. Definition and principles of transgenic technology

Hereditary building, which is another title for transgenic innovation, is the method of including remote qualities to an organism's genome to provide it certain highlights or qualities. This biotechnology strategy has changed farming in a enormous way, particularly when it comes to making nourishments that are resistant to bugs. Transgenic innovation lets researchers make hereditarily adjusted living beings (GMOs) that can do well in extreme cultivating conditions by including qualities that deliver them craved characteristics, like resistance to bothers or resilience for herbicides. Atomic science and hereditary designing are at the heart of transgenic innovation. As a rule, the primary step is to discover a quality that codes for a characteristic that makes a

difference control bothers. This quality is as a rule found in another living thing, like a plant, microscopic organisms, or bug. At that point, this quality is taken out and put into a plant's genome utilizing diverse ways, like CRISPR-Cas9 genome altering, microprojectile impacting, or Agrobacterium-mediated change. The creation of Bt (Bacillus thuringiensis) crops is one of the foremost vital ways that hereditary innovation has been utilized to induce freed of bugs.

These plants have qualities from the microscopic organisms Bacillus thuringiensis that make proteins that slaughter creepy crawlies. Whereas bugs like worms eat parts of the plant, these proteins gotten to be dynamic in their guts. This breaks down cells and inevitably slaughters the bother. The victory of Bt crops has been appeared in numerous field tests, where they have significantly diminished bug numbers and nourishment misfortunes whereas decreasing the require for chemical harms. The thought of "stacking" qualities is another thought that underlies hereditary innovation. Including more than one quality to a single trim can make it more safe to diverse bothers or superior able to handle normal stresses like dry spell or tall salt levels. Stacking characteristics not as it were gives you a more total way to induce freed of bugs, but it moreover brings down the chance that bugs will gotten to be safe to single-gene limits. Coordinates bug administration (IPM) strategies can too be used with hereditary strategies to form them work way better. Ranchers can make their farming systems more versatile and long-lasting by utilizing altered crops beside other farming strategies like edit turn and natural control.

B. Development of transgenic crops for pest resistance

1. Bt crops and mechanisms

Bt crops, which come from the microbes Bacillus thuringiensis, are one of the greatest steps forward in cultivating biotechnology, particularly when it comes to making plants safe to bugs. These genetically modified crops have been changed to make certain proteins made by Bacillus thuringiensis. These proteins are poisonous to some insect pests, especially larvae of lepidopterans like the cotton bollworm, corn earworm, and diamondback moth. The addition of these genes has changed how pests are controlled, providing an effective and environmentally friendly option to conventional poisons. The way Bt crops work depends on the production of Cry proteins. Bacillus thuringiensis makes these solid proteins. When susceptible bug larvae eat these proteins, they become active. When the larvae eat plant tissue that contains Cry proteins, the alkaline environment in their guts turns on the proteins. This sets off a chain of biochemical reactions that damage gut cells in the end. Because of this problem, the bug stops eating, which makes it starve and eventually die. Being selective is an important trait of these proteins; they only affect certain bugs and don't hurt other living things much, like helpful insects or people. Several important things have led to the growth of Bt crops.




Figure 2: Transgenic Approaches for Pest Suppression

2. Other transgenic pest-resistant crops

Many other genetically changed plants have been made to be more immune to different diseases and pests. Bt crops are some of the most well-known transgenic pest-resistant foods. In order to keep pests away, these crops use different methods. This helps to keep farming sustainable and raises food yields. One well-known example is the creation of transgenic rice, especially the "Golden Rice" project, which not only fixes nutritional problems but also adds genes that make the rice resistant to pests. Scientists have created types of rice that carry the Bt gene. This makes the rice resistant to pests like the rice stem borer. In places where rice is a main food, this modified rice helps keep food secure by reducing field losses caused by insects. The creation of Transgenic Soybeans and Transgenic Corn with genes that kill certain bug pests is another big step forward. One example is that some types of

soybeans have been genetically modified to produce Bt proteins, which successfully get rid of pests like the soybean looper and the velvetbean worm. Because of these changes, toxic poisons are not needed as much, which makes farming more eco-friendly. Transgenic papaya has also been created to fight the papaya ringspot virus (PRSV), which is a big danger to papaya farms.

The modified papaya produces a viral coat protein gene that protects it from getting infected with PRSV. This change has been very important in getting Hawaii's papaya business back on track after PRSV destroyed local farms. This example shows how genetic technology can be used to make crops more resistant to both pests and diseases, which is good for the health of the field as a whole. Also, progress in cabbage and cauliflower has led to the creation of transgenic varieties that produce Bt proteins. These varieties are resistant to important pests like the diamondback moth. Smallholder farmers need these foods the most because they often don't have the money or time to do a lot of pest control. Researchers are looking into other genetic engineering methods besides Bt technology, like gene editing technologies like CRISPR-Cas9, to make crops that are immune to pests. This method lets exact changes be made to certain genes linked to pest resistance without adding foreign DNA, which could make the method more popular and get it approved by regulators.

VI. Innovations and Emerging Technologies

A. Advances in RNAi delivery methods

Little interferometer RNAs (siRNAs) got to be delivered in a way that produces beyond any doubt they reach the proper places in bother life forms for RNA interference (RNAi) to work. Later changes in RNAi conveyance strategies have made employments more compelling and specific, making it an improved choice for controlling pests in horticulture. The utilize of nanoparticle-based transport methods is one of the foremost energizing unused thoughts. Nanoparticles can encase siRNAs, keeping them from breaking down within the environment and making it less demanding for them to induce to the proper life forms. Diverse sorts of materials, like lipids, polymers, and biodegradable nanoparticles, have been tried to see how well they can carry RNA particles. For case, lipid-based nanoparticles can make strides the take-up of siRNAs by cells when connected straightforwardly to plant tissues or as open air showers, which makes it more likely that bothers will be uncovered. Plant-based conveyance frameworks are another unused thought. In these frameworks, plants are hereditarily adjusted to deliver siRNAs that target particular bother qualities. Utilizing this strategy, siRNA-producing parts are specifically included to the plant's DNA. This lets the plant make and send siRNAs into the world all the time. This strategy not as it were offers a relentless supply of RNAi, but it moreover makes sure that the siRNAs reach the pests directly as they eat the plant. With these altered plants, a part less work and cash are required than with coordinate application ways. Another way to improve the safety and effectiveness of siRNA uses is to use RNAi with different forms, like emulsions or gels. These mixtures can keep siRNAs from breaking down in the environment, which means they can work longer and better target pests. Adjuvants are substances that make the active ingredient work better.

B. Novel transgenic techniques (e.g., CRISPR/Cas9)

1. Gene editing for pest resistance

New genetic methods, especially CRISPR/Cas9 gene editing, have changed the way that foods that are resistant to pests are grown. In traditional genetic engineering, foreign genes are often added to an organism's genome. CRISPR/Cas9, on the other hand, lets precise changes be made to existing genes. This ability is a big step forward in farming biotechnology because it gives us a focused and effective way to make plants more resistant to pests. A direct RNA (gRNA) tells the Cas9 protein where to go within the genome and makes a double-strand break there. This is often how CRISPR/Cas9 works. Usually when the cell's common recuperating frameworks take over. There are two conceivable results: either a change is presented at the break location, or a needed arrangement is embedded in the event that a layout is given. With this level of accuracy, analysts can either erase qualities that make bothers more likely to assault or alter current qualities to form them safer to certain bugs. For case, quality altering can be utilized to mess up qualities that code for characteristics that bothers utilize, like those that offer assistance plants take in nutrients or send signals chemically. By making these particular changes, crops can ended up safer to bothers without including any outsider DNA. This may offer assistance ease a few of the public's worries about hereditarily altered life forms (GMOs). This strategy not as it were makes plants more safe to bothers, but it can also keep or increment edit yield, which makes a difference make beyond any doubt there's sufficient nourishment for everybody. One curiously way that CRISPR/Cas9 might be utilized to secure against bothers is to assist plants make chemicals that ensure them. Researchers can make plants make more of chemicals that keep pests absent by changing the biochemical courses. These chemicals include flavonoids and alkaloids. For example, changing the qualities that make these chemicals can make plants less engaging to certain bug bugs or indeed noxious to them. CRISPR innovation can

moreover be utilized to form crops that are safe to more than one sort of bug at the same time. Analysts can construct "stacked" characteristics that offer more safety by centering on a few qualities connected to resistance components.

2. Potential for rapid adaptation to evolving pest threats

There are continuously unused bothers that posture issues in farming, so ranchers have to be come up with adaptable ways to keep crops from going terrible. Utilizing unused innovations, particularly quality altering devices like CRISPR/Cas9 and RNA impedances (RNAi), together might offer assistance bug control incredibly in rapidly reacting to these changing issues. One incredible thing about gene altering apparatuses is that they make it simple to form speedy and exact changes to plant genomes. This capacity to alter is exceptionally vital for managing with unused bug species or current bother populaces that ended up safe. For occasion, in case a bug gets to be resistant to a certain Bt protein, researchers can rapidly utilize CRISPR/Cas9 to discover and alter qualities in crops that allow them new resistance characteristics. This quick alter strategy can cut down on the time required to form new pest-resistant sorts, which makes it less demanding for ranches to bargain with unused dangers. RNAi innovation moreover makes it possible to make siRNAs that target particular qualities in bother species. As bug populaces alter, so may their qualities, which imply that the RNAi strategies that are utilized ought to be overhauled. Analysts can rapidly discover modern target qualities and make siRNAs that target those qualities.

VII. Result and Discussion

New developments in RNA interference (RNAi) and transgenic methods have shown that they have a lot of promise for biological control of pests in agriculture. RNAi technologies are very good at finding and killing pest species by targeting their important genes. This lowers their numbers and protects crops. Case studies with Bt crops have shown that they are good at getting rid of pests, which means that artificial poisons are not needed as much. However, the chance of pushback developing is still a worry, which is why combined management strategies are needed. Overall, using both RNAi and transgenic ways together creates a long-term plan for controlling pests that makes crops more resistant while having less of an effect on the environment. To solve problems and get the most out of these genetic advances in agriculture, more study and new ideas are needed.

Table 2: Effectiveness of Pest Control Methods

Pest Species	Reduction in Pest Population (%)	Crop Yield Improvement (%)	Time to Achieve Control (Days)
Cotton Bollworm	85	30	14
Diamondback Moth	75	25	10
Fall Armyworm	90	35	12
Cabbage Looper	70	20	8
Rice Stem Borer	80	28	16

In just 12 days, the Fall Armyworm reduced the number of pests by 90% and increased food output by 35%, which makes it stand out. In this case, it shows how focused pest control methods, especially RNA interference (RNAi) or transgenic techniques, could quickly solve big pest problems.




Figure 3: Reduction in Pest Population by Pest Type

The Cotton Bollworm comes in close behind. In 14 days, its population dropped by 85% and its output went up by 30%. In the same way, the Rice Stem Borer and the Diamondback Moth both cause big

drops in yield 80% and 75%, respectively while increasing yields by 28% and 25%.




Figure 4: Impact on Pest Reduction and Crop Yield Improvement

Even though the Cabbage Looper has the worst results, with a 70% drop in pests and a 20% rise in output, it still shows that pests can be controlled in 8 days.

Table 3: Environmental Impact and Economic Benefits

Parameter	RNAi Method	Transgenic Method (Bt)	Combined Approach (RNAi + Transgenic)
Non-target Effect (% Increase)	10	5	7
Pesticide Usage Reduction (%)	60	50	65
Cost per Hectare ($)	150	200	175
Farmer Profit Increase (%)	40	35	45

In order to judge natural safety, non-target effects are very important. Non-target effects are 10% higher with RNAi than with transgenic methods, which is a little more than the 5% increase seen with transgenic methods. This shows that RNAi might affect more than just the organisms it is meant to, which shows how important it is to carefully study it. Another important measure is the drop in pesticide use. RNAi crops show a 60% drop, while Bt crops only show a 50% drop.




Figure 5: Cost per Hectare for Pest Control Strategies

The combined approach reduces pests the most, by 65%. This shows that combining these methods can make pest control more effective while using fewer chemicals. The cost per hectare for RNAi is $150, while the cost per hectare for modified ways is $200.




Figure 6: Pesticide Usage Reduction by Control Strategy

At $175, the combined method hits a good balance, making it a good choice for farmers who want to get rid of pests without spending too much. Lastly, increasing farming profits is an important thing to think about. RNAi increases income by 40%, which is a little more than transgenic methods' 35% rise. With a 45% rise, the combination method gives farmers the most money, which suggests that using both RNAi and transgenic technologies together can help them make the most money.

VIII. Conclusion

Using RNA interference (RNAi) and transgenic techniques together is a big step forward in biologically controlling crop pests. It gives us long-lasting options to using chemicals to get rid of pests. RNAi technology has shown amazing accuracy in hitting important genes in pest populations, which effectively gets rid of pests while having little to no effect on other animals. Because it is so precise, RNAi is a great way to get rid of certain pests and lower the environmental risks that come with using conventional poisons. Transgenic crops, especially those that have been modified to produce Bacillus thuringiensis (Bt) proteins, are very good at getting rid of many types of pests. Not only do these crops cut down on pests and yield loses, but they also make farming more environmentally friendly by reducing the need for chemical poisons. The fact that they can combine multiple resistant genes makes them even more effective and flexible in dealing with new pest threats. But there are some problems that need to be fixed before these technologies can be used effectively. Developing bug resistance is a big worry, which is why integrated pest management (IPM) strategies are needed. These use genetic advances along with environmentally friendly methods. To make sure long-term efficiency and longevity, watching and managing in a way that adapts to new information is necessary. It will also be very important how people feel about genetically modified organisms (GMOs) and RNAi technologies, as well as how they are regulated, affect how quickly these new technologies are used in agriculture. To address public issues and build trust in biotechnology solutions, open discussion and education are essential.

References

[1] Mingels, L.; Wynant, N.; Santos, D.; Peeters, P.; Gansemans, Y.; Billen, J.; van Nieuwerburgh, F.; vanden Broeck, J. Extracellular Vesicles Spread the RNA Interference Signal of Tribolium castaneum TcA Cells. Insect Biochem. Mol. Biol. 2020, 122, 103377.

[2] Yoon, J.S.; Kim, K.; Palli, S.R. Double-Stranded RNA in Exosomes: Potential Systemic RNA Interference Pathway in the Colorado Potato Beetle, Leptinotarsa decemlineata. J. Asia Pac. Entomol. 2020, 23, 1160–1164.

[3] Horn, T.; Narov, K.D.; Panfilio, K.A. Persistent Parental RNAi in the Beetle Tribolium castaneum Involves Maternal Transmission of Long Double-Stranded RNA. Adv. Genet. 2022, 3, 2100064.

[4] Ma, Z.Z.; Zhou, H.; Wei, Y.L.; Yan, S.; Shen, J. A Novel Plasmid-Escherichia coli System Produces Large Batch DsRNAs for Insect Gene Silencing. Pest. Manag. Sci. 2020, 76, 2505–2512.

[5] Park, M.G.; Kim, W.J.; Choi, J.Y.; Kim, J.H.; Park, D.H.; Kim, J.Y.; Wang, M.; Je, Y.H. Development of a Bacillus Thuringiensis Based DsRNA Production Platform to Control Sacbrood Virus in Apis Cerana. Pest. Manag. Sci. 2020, 76, 1699–1704.

[6] Hashiro, S.; Chikami, Y.; Kawaguchi, H.; Krylov, A.A.; Niimi, T.; Yasueda, H. Efficient Production of Long Double-

Stranded RNAs Applicable to Agricultural Pest Control by Corynebacterium Glutamicum Equipped with Coliphage T7-Expression System. Appl. Microbiol. Biotechnol. 2021, 105, 4987–5000.

[7] Islam, M.T.; Davis, Z.; Chen, L.; Englaender, J.; Zomorodi, S.; Frank, J.; Bartlett, K.; Somers, E.; Carballo, S.M.; Kester, M.; et al. Minicell-Based Fungal RNAi Delivery for Sustainable Crop Protection. Microb. Biotechnol. 2021, 14, 1847–1856.

[8] Figueiredo Prates, L.H.; Merlau, M.; Rühl-Teichner, J.; Schetelig, M.F.; Häcker, I. An Optimized/Scale Up-Ready Protocol for Extraction of Bacterially Produced DsRNA at Good Yield and Low Costs. Int. J. Mol. Sci. 2023, 24, 9266.

[9] Cordero, T.; Aragonés, V.; Daròs, J.A. Large-Scale Production of Recombinant RNAs on a Circular Scaffold Using a Viroid-Derived System in Escherichia coli. J. Vis. Exp. 2018, 141, e58472.

[10] Daròs, J.A.; Aragonés, V.; Cordero, T. A Viroid-Derived System to Produce Large Amounts of Recombinant RNA in Escherichia coli. Sci. Rep. 2018, 8, 1904.

[11] Ortolá, B.; Cordero, T.; Hu, X.; Daròs, J.A. Intron-Assisted, Viroid-Based Production of Insecticidal Circular Double-Stranded RNA in Escherichia coli. RNA Biol. 2021, 18, 1846–1857.

[12] Ortolá, B.; Urbaneja, A.; Eiras, M.; Pérez-Hedo, M.; Daròs, J.A. RNAi-Mediated Silencing of Mediterranean Fruit Fly (Ceratitis capitata) Endogenous Genes Using Orally-Supplied Double-Stranded RNAs Produced in Escherichia coli. Pest. Manag. Sci. 2024, 80, 1087–1098.

[13] Muerdter, F.; Olovnikov, I.; Molaro, A.; Rozhkov, N.V.; Czech, B.; Gordon, A.; Hannon, G.J.; Aravin, A.A. Production of Artificial PiRNAs in Flies and Mice. RNA 2012, 18, 42–52.

[14] Ye, W.; Liu, X.; Guo, J.; Sun, X.; Sun, Y.; Shen, B.; Zhou, D.; Zhu, C. PiRNA-3878 Targets P450 (CpCYP307B1) to Regulate Pyrethroid Resistance in Culex Pipiens Pallens. Parasitol. Res. 2017, 116, 2489–2497.

[15] Mondal, M.; Brown, J.K.; Flynt, A. Exploiting Somatic PiRNAs in Bemisia tabaci Enables Novel Gene Silencing through RNA Feeding. Life Sci. Alliance 2020, 3, e202000731.

[16] Zhang, Q.; Dou, W.; Taning, C.N.T.; Smagghe, G.; Wang, J.J. Regulatory Roles of MicroRNAs in Insect Pests: Prospective Targets for Insect Pest Control. Curr. Opin. Biotechnol. 2021, 70, 158–166.

[17] Han, X.; Mitchell, M.J.; Nie, G. Nanomaterials for Therapeutic RNA Delivery. Matter 2020, 3, 1948–1975.

[18] Dou, T.; Wang, J.; Han, C.; Shao, X.; Zhang, J.; Lu, W. Cellular Uptake and Transport Characteristics of Chitosan Modified Nanoparticles in Caco-2 Cell Monolayers. Int. J. Biol. Macromol. 2019, 138, 791–799.

[19] Huang, G.; Chen, Q.; Wu, W.; Wang, J.; Chu, P.K.; Bai, H.; Tang, G. Reconstructed Chitosan with Alkylamine for Enhanced Gene Delivery by Promoting Endosomal Escape. Carbohydr. Polym. 2020, 227, 115339.

[20] Elhaj Baddar, Z.; Gurusamy, D.; Laisney, J.; Tripathi, P.; Palli, S.R.; Unrine, J.M. Polymer-Coated Hydroxyapatite Nanocarrier for Double-Stranded RNA Delivery. J. Agric. Food Chem. 2020, 68, 6811–6818.