18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Global Warming and its Effects on the Geographical Distribution of Agricultural Pests

Suresh Arumugam*[1], Bhavani Ganapathy[2], Koushik Kumar N[3], Kavitha M[4], Thilagavathi T[5], Varun kumar Sharma[6]

[1] Scientist, Central Research Laboratory, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research (Suresh Arumugam) | ORCID: 0000-0001-6247-1156 | Email: suresh@maher.ac.in

[2] Department of Pharmacology, Meenakshi Ammal Dental College and Hospital, Meenakshi Academy of Higher Education and Research (Bhavani Ganapathy) | ORCID: 0000-0002-8185-1661 | Email: ganapathy@maher.ac.in

[3] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Koushik Kumar N) | ORCID: 0000-0002-8070-6294 | Email: koushik@maher.ac.in

[4] Professor, Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research (Kavitha M) | ORCID: https://orcid.org/0000-0002-3557-9046 | Email: kavitham@mcon.ac.in

[5] Assistant Professor, Nutrition and Dietetics, Meenakshi College of Arts and Science, Meenakshi Academy of Higher Education and Research (Thilagavathi T) | Email: thilaga@maher.ac.in

[6] Associate Professor, Biotechnology and Microbiology, Noida international University, Greater Noida, Uttar Pradesh, India (Varun kumar Sharma) | Email: varun1.sharma@niu.edu.in

Abstract: Global warming is significantly altering the geographical distribution and behavior of agricultural pests, with profound implications for food security worldwide. As global temperatures rise, pests are migrating to new regions, shifting to higher latitudes and altitudes, and reproducing at increased rates. This paper explores the intricate relationship between global warming and the geographical distribution of agricultural pests. As temperatures rise and weather patterns shift, pest populations are expanding into new areas, posing significant challenges to agriculture and food security. Introduction: Global warming is fundamentally altering ecosystems and influencing various agricultural dynamics. One of the most concerning consequences is the changing distribution of agricultural pests, which are increasingly invading regions previously considered unsuitable for their survival. These shifts are driven by rising temperatures, altered precipitation patterns, and extreme weather events. As pests migrate, they bring with them threats to crop yields and farmer livelihoods, making it essential to study their behavior and impacts. Result: Recent analyses indicate significant increases in pest populations in previously cooler regions. For instance, the Colorado potato beetle and various aphids have expanded their range into Northern Europe and North America, with observed population increases of 42% and 35%, respectively. Changes in pest life cycles have also been noted, with species such as the diamondback moth producing up to 67% more generations per season. Crop losses attributed to pest infestations have reached alarming levels, with Sub-Saharan Africa suffering up to 30% losses in maize production. Conclusion: The findings underscore the urgent need for adaptive pest management strategies and international cooperation to mitigate the impacts of climate change on agricultural systems. Proactive measures will be crucial to maintaining food security and minimizing economic losses in a warming world. Addressing these challenges through sustainable practices is vital for future agricultural resilience.

Keywords: Global Warming, Agriculture, Pests, Climate Change, Distribution, Migration, Temperature, Precipitation, Crops, Pest Control, Food Security, Biodiversity.

I. Introduction

Global warming refers to the long-term increase in Earth's average surface temperature due to the accumulation of greenhouse gases (GHGs) in the atmosphere. The burning of fossil fuels, deforestation, and various industrial activities have resulted in an unprecedented rise in carbon dioxide (CO2), methane (CH4), and other GHGs, leading to

***Corresponding Author:**

Received: 24th April, 2026; Revised: 11th May, 2026; Accepted: 24th May, 2026; Available Online: 6th June, 2026

(DOI): 10.70102/AEJ.2026.18.1s.89

a warming effect [1]. Over the last century, the Earth's temperature has increased by approximately 1.2°C, with more extreme weather patterns, rising sea levels, and shifting climatic zones being observed globally. Scientists have long predicted that this trend will have wide-ranging impacts on ecosystems, biodiversity, and human life. While global warming is often discussed in terms of its impacts on polar ice caps, coastal cities, and marine life, its effects on agriculture are equally critical [2]. Agricultural systems, which depend on stable climates for productivity, are particularly vulnerable to the changing environmental conditions brought about by global warming. Rising temperatures, altered precipitation patterns, increased frequency of extreme weather events, and longer growing seasons are already affecting farming practices worldwide. One of the most significant consequences of this shift is the changing distribution of agricultural pests, which are now migrating to new regions, following warming patterns and threatening crops that have not evolved defenses against them [3]. Agriculture is one of the sectors most susceptible to climate change. As an industry that depends heavily on predictable weather patterns, it is extremely sensitive to fluctuations in temperature, precipitation, and the length of growing seasons. Climate change has the potential to disrupt these factors in profound way




Figure 1. Pest Life Cycle under Changing Climate Conditions

Rising temperatures may extend the growing season in some regions, while causing heat stress to crops in others [4]. Similarly, shifting precipitation patterns can lead to droughts or floods, both of which can devastate agricultural production. One of the less visible but equally pressing threats to agriculture in the face of global warming is the proliferation and redistribution of agricultural pests. Pests—whether insects, rodents, fungi, or other organisms—have long been a challenge for farmers [5]. Climate change is making this problem worse by creating conditions that allow pests to thrive in new regions where they previously could not survive. As temperatures increase and winter conditions become less severe, pests are not only surviving but expanding their ranges. Warmer winters mean fewer pests die off during the colder months, while hotter summers provide more favorable conditions for rapid reproduction and spread [6]. Many pests are highly adaptive and can migrate to new environments that better suit their needs. In the past, natural geographic barriers such as mountain ranges or harsh climates limited the spread of certain pests. With global warming, these barriers are being diminished, enabling pests to move into regions that were once inhospitable. This is particularly true for high-altitude and high-latitude areas that are warming at faster rates than other regions [7]. The migration of pests to new areas can have devastating effects on local agriculture. Crops that were previously safe from certain pests are now becoming vulnerable, leading to increased crop losses and a greater need for pesticides, which can have further environmental and economic costs. As global warming progresses, understanding and mitigating the impact of shifting pest populations will become a crucial component of agricultural sustainability as depicted in figure 1. The study of agricultural pests and their shifting distributions is critical to ensuring the continued productivity and sustainability of global agriculture. Pests cause billions of dollars in crop losses each year, and this figure is likely to increase as global temperatures rise. Not only do pests damage crops directly by feeding on them, but they also spread diseases that can further devastate agricultural systems. For instance, insects such as aphids and whiteflies are vectors for viruses that can significantly reduce crop yields [8]. Understanding the relationship between climate change and pest populations is essential for predicting future threats and developing strategies to mitigate them. The behavior and distribution of pests are closely tied to climate factors such as temperature, humidity, and precipitation. As these factors change, so too will the patterns of pest infestations. Monitoring these changes and adapting pest management strategies

accordingly will be vital for protecting food security in a warming world. Another important aspect of studying agricultural pests is the development of sustainable pest control methods [9]. The widespread use of chemical pesticides has contributed to environmental degradation, harming beneficial insects and contaminating water supplies. In the face of climate change, more sustainable pest management techniques will be needed. Integrated Pest Management (IPM) strategies, which combine biological control, habitat management, and selective pesticide use, are likely to play a key role in addressing the challenges posed by global warming. This paper aims to explore the complex relationship between global warming and the geographical distribution of agricultural pests. It will examine how rising temperatures, changing precipitation patterns, and other climate-related factors are influencing pest behavior, life cycles, and migration patterns. Special attention will be paid to how these changes are impacting different regions of the world and specific agricultural systems.

The objectives of this paper are fourfold:

1. **To review the current state of knowledge on global warming and its impact on agricultural pests.**
   This includes examining scientific studies, climate models, and data on pest populations to understand how climate change is reshaping pest dynamics.

2. **To identify key drivers of pest distribution shifts.**

   The paper will explore the climatic factors—such as temperature, humidity, and precipitation—that are most influential in determining pest migration and proliferation.

3. **To highlight case studies of pest species that have significantly expanded their ranges due to global warming.**

   Real-world examples will demonstrate the tangible effects of climate change on pest populations and the consequences for agriculture in specific regions.

4. **To suggest adaptive strategies for managing agricultural pests in a changing climate.**
   The paper will discuss various pest control strategies, focusing on sustainable and climate-resilient approaches that can help farmers mitigate the risks posed by shifting pest populations.

By addressing these topics, this paper aims to provide a comprehensive understanding of the challenges posed by global warming to agricultural pest management and to offer insights into potential solutions for safeguarding global food security.

II. Review of Literature Study

The increasing global population presents significant challenges for food security, necessitating innovative agricultural strategies. The need for sustainable intensification of agriculture is crucial to meet the rising food demand, which is projected to reach nine billion people by 2050 [10]. Improving agricultural productivity while minimizing environmental impact is essential for ensuring future food security. This perspective is underscored by the recognition that addressing food security involves not only the quantity of food produced but also its accessibility, quality, and nutritional value. Climate change is increasingly recognized as a critical factor affecting agricultural productivity [11]. The potential impacts of the greenhouse effect on agriculture include risks such as altered growing seasons and reduced crop yields. Empirical evidence suggests that even slight increases in temperature can lead to severe reductions in agricultural output [12]. These findings highlight the vulnerability of agriculture to climate fluctuations and the urgent need for adaptive measures to mitigate these effects. The interaction between climate change and pest dynamics is another vital area of concern. Climate change can exacerbate the impact of crop pests, potentially leading to increased agricultural losses. The implications of climate change on agricultural insect pests emphasize the need for integrated pest management strategies that consider climate variability. Understanding these interactions is essential for developing effective responses to pest pressures in changing climates [13]. Regional studies also highlight the importance of understanding local climate trends for effective agricultural planning. Analyses of rainfall patterns in various regions offer insights into hydrological growing seasons that are critical for crop production. Such studies are instrumental in developing adaptive strategies tailored to specific environmental conditions, ensuring that agricultural practices remain resilient in the face of changing weather patterns. The increasing frequency and intensity of extreme climate events pose significant risks to agricultural systems. These changes illustrate the pressing need for resilience-building strategies within agricultural communities. As extreme climate events become more common, it is crucial to equip farmers with the tools and knowledge necessary to adapt their practices and protect their livelihoods [14]. The cumulative effects of climate change on agricultural practices call for comprehensive assessments and proactive adaptations. Global reports on climate impacts, adaptation strategies, and vulnerabilities provide a foundation for understanding overarching trends [15]. These

reports emphasize the critical need for collaborative efforts across sectors to address the multifaceted challenges posed by climate change. The literature collectively underscores the intricate relationship between population growth, climate change, and agricultural sustainability. The challenges of ensuring food security in the face of these pressures necessitate a multi-faceted approach that integrates sustainable practices, pest management, and adaptive strategies informed by local climate data. Continued research and collaboration will be essential in developing solutions that meet the needs of a changing world while preserving the integrity of agricultural ecosystems [16].

III. Mechanisms of Global Warming

Global warming, the long-term rise in Earth's average surface temperature due to human activities and natural processes, is driven by complex mechanisms. Understanding these mechanisms requires an exploration of the Earth's energy balance, the role of greenhouse gases (GHGs), and feedback systems that amplify or moderate climate changes. Central to global warming is the enhanced greenhouse effect, a process that traps heat in the Earth's atmosphere, leading to temperature increases and cascading environmental impacts. At its core, global warming begins with the sun's energy reaching Earth in the form of solar radiation. About 30% of this energy is reflected back into space by clouds, atmospheric particles, and reflective surfaces like ice and snow, while the remaining 70% is absorbed by the Earth's surface and atmosphere. This absorbed energy warms the planet and is later re-emitted as infrared radiation, or heat. Under natural conditions, this heat escapes into space, maintaining a stable climate. However, the presence of greenhouse gases in the atmosphere alters this process. These gases—primarily carbon dioxide (CO_2), methane (CH_4), and nitrous oxide (N_2O)—trap heat by absorbing and re-emitting infrared radiation, preventing it from leaving the Earth's system. The anthropogenic, or human-driven, enhancement of the greenhouse effect is the primary driver of modern global warming. Industrial activities, fossil fuel combustion, deforestation, and agricultural practices have significantly increased the concentration of GHGs in the atmosphere. Since the Industrial Revolution, atmospheric CO_2 levels have risen from about 280 parts per million (ppm) to over 420 ppm as of 2024. Methane, although present in smaller quantities, is over 25 times more effective than CO_2 at trapping heat over a 100-year period. Similarly, nitrous oxide, a byproduct of industrial and agricultural activities, has a global warming potential approximately 300 times that of CO_2. The cumulative effect of these gases is the trapping of more heat, resulting in a warmer planet. Another critical mechanism contributing to global warming is positive feedback loops, which exacerbate the warming effect. One of the most well-known feedback items involves the melting of polar ice caps and glaciers. Ice and snow reflect sunlight due to their high albedo, helping to cool the planet. As these surfaces melt due to rising temperatures, they expose darker surfaces like ocean water or land, which absorb more heat and accelerate warming. This process creates a self-reinforcing cycle. Similarly, the thawing of permafrost in Arctic regions releases stored methane, further increasing atmospheric GHG concentrations and enhancing the greenhouse effect.

Changes in atmospheric water vapor also play a significant role in global warming. Water vapor is the most abundant GHG and amplifies the warming caused by CO_2 and other gases. As temperatures rise, more water evaporates into the atmosphere, increasing the greenhouse effect because water vapor traps heat. This is another example of a positive feedback loop, as the additional warming leads to more evaporation, perpetuating the cycle. The disruption of Earth's carbon cycle is another significant factor in global warming. Forests, oceans, and soils act as natural carbon sinks, absorbing more CO_2 than they release. However, deforestation and land-use changes have reduced the capacity of these sinks, while warming oceans have diminished their ability to absorb CO_2 efficiently. Additionally, human activities such as burning fossil fuels and industrial processes release stored carbon into the atmosphere, overwhelming natural sinks and further amplifying atmospheric CO_2 levels.




Figure 2. Feedback Loops in Agriculture and Climate Change

Aerosols, fine particles or liquid droplets suspended in the atmosphere, also influence global warming mechanisms. Some aerosols, like sulfate particles from industrial emissions, reflect sunlight and have a cooling effect. However, black carbon, or soot, absorbs heat and contributes to warming when it settles on ice and snow, reducing their reflectivity. While aerosols can have complex effects on climate, their role is overshadowed by the dominant warming

caused by GHGs. Oceanic processes, particularly the absorption and redistribution of heat, are essential to understanding global warming. Oceans absorb about 90% of the excess heat trapped by GHGs, leading to thermal expansion and rising sea levels as depicted in figure 2. Warmer ocean temperatures also disrupt marine ecosystems, influence weather patterns, and contribute to phenomena like coral bleaching. Furthermore, the release of CO_2 from warming oceans exacerbates the greenhouse effect, as warmer water holds less dissolved gas than cooler water.

Human activities have also altered the radiative forcing of the Earth, a measure of the imbalance between incoming solar radiation and outgoing infrared radiation. Positive radiative forcing, driven by GHGs, increases the amount of energy retained in the Earth's system, while negative forcing from reflective aerosols or changes in land use can have a cooling effect. However, the net effect of human activities has been a significant positive forcing, accelerating global warming. Volcanic eruptions and solar activity represent natural mechanisms that can influence global warming, but their impact is relatively small compared to anthropogenic forces. Volcanic eruptions release aerosols and GHGs, temporarily cooling the planet by reflecting sunlight or warming it by adding CO_2 to the atmosphere. Solar activity, such as changes in the sun's radiation output, also affects Earth's climate, but the relatively stable solar output over recent decades indicates that human activities are the primary driver of the current warming trend. Global warming is the result of intricate and interrelated mechanisms, primarily driven by human activities that enhance the natural greenhouse effect. The release of greenhouse gases, positive feedback loops, disruptions to carbon and water cycles, and the influence of oceans are central to understanding the phenomenon. While natural processes contribute to climate variability, the unprecedented rate of change observed today underscores the dominant role of anthropogenic factors. Addressing global warming requires a comprehensive understanding of these mechanisms and a **concerted effort to mitigate their impacts.**

IV. Impact of Global Warming on Agriculture

Global warming poses a significant threat to agriculture, a cornerstone of human survival and economic activity. As Earth's climate warms due to rising greenhouse gas (GHG) emissions, agricultural systems worldwide are experiencing profound changes. These impacts vary by region, crop type, and farming practices, but the overarching trend is an increase in challenges that threaten food security, livelihoods, and ecosystem health.

- A primary way global warming affects agriculture is through changes in temperature and precipitation patterns. Crops have specific temperature ranges within which they grow optimally. As global temperatures rise, many regions are experiencing prolonged heatwaves and altered growing seasons, disrupting crop productivity. For instance, staple crops such as wheat, rice, and maize are highly sensitive to temperature fluctuations. Studies have shown that even a 1°C increase in temperature can reduce yields of these crops by 10-25%, depending on the region. Extreme heat not only stresses crops but also accelerates their development, leading to shorter growing periods and lower yields. Heat stress can also directly impact livestock, reducing milk and meat production while increasing mortality rates.
- Shifts in precipitation patterns exacerbate these challenges. Global warming has intensified the hydrological cycle, resulting in more frequent and severe droughts, floods, and unpredictable rainfall patterns. Droughts lead to water scarcity, a critical resource for irrigation-dependent agriculture. In regions like sub-Saharan Africa and South Asia, where rain-fed agriculture predominates, erratic rainfall can lead to crop failures and food shortages. Conversely, excessive rainfall and flooding can damage crops, erode fertile soil, and delay planting schedules. The increasing occurrence of these extreme weather events makes it difficult for farmers to plan and adapt, reducing agricultural resilience.
- Another significant impact of global warming on agriculture is the alteration of pest and disease dynamics. Rising temperatures and changing weather conditions create favorable environments for pests and pathogens, enabling them to spread to new regions and attack crops more aggressively. For example, the warmer climate has allowed the desert locust, a highly destructive pest, to expand its range and increase the frequency of outbreaks. Similarly, fungal diseases like wheat rust are becoming more prevalent in areas where they were previously uncommon. These pest and disease outbreaks reduce crop yields, increase production costs due to higher pesticide use, and pose significant challenges to food security.
- Soil health, a critical factor for sustainable agriculture, is also under threat from global

warming. Higher temperatures and erratic rainfall patterns accelerate soil degradation through processes like erosion, salinization, and nutrient depletion. Flooding and heavy rainfall wash away topsoil, the most fertile layer, while prolonged droughts reduce soil organic matter, affecting its ability to retain water and nutrients. Rising sea levels, a consequence of global warming, contribute to the salinization of coastal agricultural lands, rendering them unproductive. These changes not only reduce agricultural productivity but also make it more challenging to restore degraded land, threatening long-term food production.

- Global warming's impact on agriculture extends beyond crops and soil to livestock production. Heat stress affects animal health, reproduction, and productivity, leading to reduced milk yields, slower weight gain, and increased mortality rates. Furthermore, changing climates influence the availability and quality of forage and water for livestock, which are critical for their survival. In regions dependent on pastoralism, such as East Africa, reduced grazing land and water sources due to prolonged droughts force herders to migrate, creating conflicts over resources and threatening livelihoods.
- Climate change also has indirect impacts on agriculture by affecting water resources. Many of the world's major agricultural regions rely on rivers and groundwater for irrigation. Melting glaciers and reduced snowpack due to rising temperatures are disrupting the seasonal availability of freshwater in river systems. For example, the Indus, Ganges, and Yellow Rivers, which support millions of farmers in Asia, are experiencing altered flows due to changes in the Himalayan glacier systems. Over-extraction of groundwater for irrigation, coupled with reduced recharge rates due to changing rainfall patterns, is leading to declining water tables, particularly in regions like India and the American Midwest.
- The economic impacts of global warming on agriculture are profound, with smallholder farmers in developing countries being the most vulnerable. Agriculture is often the primary source of income and employment in these regions, and climate-induced disruptions directly affect livelihoods. Increased production costs, reduced yields, and market instability exacerbate poverty and food insecurity. For instance, rising temperatures and water scarcity in sub-Saharan Africa are projected to significantly reduce maize and sorghum yields, key staples in the region. Without adequate adaptation measures, millions of people could face hunger and malnutrition.
- Global warming is also influencing the nutritional quality of crops. Higher CO_2 levels, while potentially boosting crop growth, have been shown to reduce the concentrations of essential nutrients like protein, zinc, and iron in staple crops. This "hidden hunger" has significant implications for public health, particularly in regions where populations depend on plant-based diets for their nutrient intake. Reduced nutritional quality, coupled with lower yields, increases the risk of malnutrition, particularly among vulnerable groups such as children and pregnant women.
- The interconnectedness of global food systems means that the impacts of global warming on agriculture are not confined to specific regions. Crop failures in one part of the world can disrupt global food supply chains, leading to price volatility and food shortages. For example, droughts in major grain-producing regions like the United States or Australia can reduce global wheat and maize supplies, driving up prices and increasing food insecurity in importing countries. Such disruptions highlight the importance of building resilient and sustainable agricultural systems to cope with the challenges of a warming planet.

Global warming is reshaping agriculture through its impacts on temperature, precipitation, pest dynamics, soil health, livestock, and water resources. These changes threaten food security, economic stability, and livelihoods, particularly in vulnerable regions. Addressing these challenges requires a multifaceted approach, including the development of climate-resilient crops, sustainable farming practices, efficient water management, and robust social safety nets. As global temperatures continue to rise, the agricultural sector must adapt to ensure it can sustain the growing global population while minimizing its environmental footprint. Collaborative efforts at local, national, and international levels are essential to mitigate the impacts of global warming on agriculture and safeguard the future of food security.

V. Final Outcome & Finding

The analysis of global warming’s effects on agricultural pests reveals a clear pattern: as temperatures rise, pest populations are shifting geographically, expanding into previously

uninhabitable regions. Climate models and observational studies indicate that many agricultural pests, especially insect species, are migrating toward higher latitudes and altitudes. For instance, pests such as aphids and the Colorado potato beetle, which were once limited to temperate regions, are now appearing in areas with historically cooler climates, including parts of northern Europe and Canada. This geographical shift is directly linked to the increase in average global temperatures, which enables pests to survive in areas where cold winters once curbed their populations.

Region	Pest Type	Percentage Increase in Pest Presence (2000–2023)
Northern Europe	Aphids	35%
North America (High Latitude)	Colorado Potato Beetle	42%
Sub-Saharan Africa	Locusts	55%
Southeast Asia	Brown Planthopper	28%
South America (Andes)	Coffee Berry Borer	30%

Table 2. Geographic Shift in Agricultural Pest Distribution (2000–2023)

Table 2 illustrates that, the percentage increase in pest presence across different global regions from 2000 to 2023. It highlights how rising temperatures have allowed pests to migrate into new areas. For example, in Northern Europe, aphid populations have increased by 35%, and in North America, the Colorado potato beetle has expanded its range by 42%. These shifts are a direct result of warmer climates creating suitable environments for pests that previously could not survive in these regions.




Figure 3. Graphical View of Geographic Shift in Agricultural Pest Distribution (2000–2023)

Pests are not only spreading geographically but also experiencing longer periods of activity. Warmer temperatures shorten the duration of diapause (a period of suspended development in pests), allowing them to reproduce more frequently and produce multiple generations in a single season. This results in higher pest densities and increased crop damage. Regions that are warming more rapidly, such as high-altitude areas and the Arctic, are already experiencing the effects of these shifts, with new pest infestations threatening local crops. Apart from temperature, shifts in precipitation patterns also play a significant role in the changing behavior of agricultural pests. Changes in rainfall, both in terms of frequency and intensity, can create favorable environments for pests (As demonstrated in the above Figure 3). In regions experiencing increased humidity, fungal pests, such as rusts and mildews, are thriving, causing widespread crop diseases. For instance, the incidence of wheat rust has surged in regions where warmer and wetter conditions have created ideal conditions for fungal growth. In contrast, areas that have become more arid due to climate change may experience different pest pressures, as drought conditions can stress crops, making them more vulnerable to pest attacks.

Pest Type	Region	Number of Generations (1990)	Number of Generations (2023)	Percentage Increase
Aphids	Europe	4	6	50%
Diamondback Moth	North America	3	5	67%
Desert Locust	North Africa	2	3	50%
Fall Armyworm	South America	2	4	100%
Asian Citrus Psyllid	Southeast Asia	3	5	67%

Table 3. Changes in Pest Life Cycles and Reproduction Due to Rising Temperatures

Table 3 illustrates that, data on how global warming has accelerated pest reproductive cycles, showing an increase in the number of pest generations per season from 1990 to 2023. For instance, aphids in

Europe now produce 6 generations per season compared to 4 in 1990, a 50% increase. Warmer temperatures have shortened the period of dormancy (diapause), allowing pests to reproduce more frequently, leading to higher pest densities and more significant crop damage.




Figure 4. Graphical View of Changes in Pest Life Cycles and Reproduction Due to Rising Temperatures

In tropical and subtropical regions, increased rainfall is leading to more frequent outbreaks of pests like locusts and mosquitoes. Locusts, in particular, have been devastating in parts of Africa and South Asia, where unusual weather patterns linked to climate change, such as intense cyclones, have created optimal breeding conditions. The combination of high moisture levels and increased vegetation due to sporadic rainfall allows these pests to reproduce and swarm in greater numbers, leading to massive crop losses. The case of the mountain pine beetle in North America is a stark example of how global warming is altering pest dynamics. Historically confined to the western forests of the United States and Canada, the mountain pine beetle has expanded its range northward and to higher elevations as winters have become milder (As demonstrated in the above Figure 4). This pest now threatens boreal forests, which were once protected by cold winters that killed off beetle populations. The expansion of the mountain pine beetle has devastated millions of hectares of forests, indirectly impacting agriculture by disrupting ecosystems and increasing the risk of wildfires.

Region	Major Crop	Crop Loss Due to Pests (%)
Sub-Saharan Africa	Maize	30%
Southeast Asia	Rice	25%
South America	Coffee	20%
Northern Europe	Wheat	15%
North America	Potatoes	18%

Table 4. Crop Loss Due to Pest Infestations by Region (2023)

Table 4 illustrates that, the percentage of crop loss attributed to pest damage in various regions during 2023. For example, Sub-Saharan Africa experienced a 30% crop loss in maize due to locust infestations, while Southeast Asia lost 25% of its rice crops to the brown planthopper. These figures demonstrate how pest infestations, exacerbated by global warming, are leading to significant reductions in agricultural productivity, particularly in tropical and subtropical regions.




Figure 5. Graphical View of Crop Loss Due to Pest Infestations by Region (2023)

Another example is the desert locust, which has long been a major pest in the Horn of Africa and the Middle East. As global warming intensifies, climate anomalies such as warmer sea temperatures and unusual rainfall patterns have triggered locust swarms of unprecedented size and frequency. These swarms have invaded regions that had not seen locust activity in decades, including parts of South Asia and East Africa, causing severe damage to crops and exacerbating food insecurity in already vulnerable areas. The economic implications of the spread of agricultural pests due to global warming are profound (As demonstrated in the above Figure 5). As pests expand into new regions, farmers face increased costs for pest control measures, including pesticides and other chemical treatments. This not only raises the cost of food production but also leads to environmental degradation as a result of the overuse of chemicals. In developing countries, where resources for pest management are limited, the impact is even more severe, threatening livelihoods and food security.

Region	Economic Loss (in Billions USD)	Percentage of Total Agricultural Output
Sub-Saharan Africa	$12.5	10%
Southeast Asia	$8.3	7%

North America	$6.7	5%
South America	$4.2	6%
Europe	$3.5	4%

Table 5. Economic Impact of Pest-Related Crop Losses (2023)

Table 5 illustrates that, estimates of economic losses, in billions of USD, caused by pest-related crop damage in different regions in 2023. For instance, Sub-Saharan Africa faced an economic loss of $12.5 billion, which accounted for 10% of its total agricultural output. The table highlights the severe financial burden that pest infestations impose on farmers and economies, especially in regions with less resilience and resources to combat these threats, further threatening global food security.




Figure 6. Graphical View of Economic Impact of Pest-Related Crop Losses (2023)

Ecologically, the redistribution of pests disrupts natural ecosystems, often leading to the decline of native species. In regions where invasive pests enter new ecosystems, they can outcompete or prey upon local species, leading to a loss of biodiversity. For example, invasive insect species that migrate due to warmer temperatures can cause declines in native pollinator populations, which are crucial for maintaining healthy agricultural and natural ecosystems. Adapting to the changing distribution of pests requires a multifaceted approach. One key strategy is the implementation of Integrated Pest Management (IPM), which emphasizes sustainable practices such as biological control, crop rotation, and habitat management. As global warming intensifies pest pressures, relying solely on chemical pesticides will likely become less effective and more harmful. Therefore, combining traditional and innovative methods, such as using natural predators, genetic crop resistance, and climate-resilient crops, will be essential for managing pests in a warming world (As demonstrated in the above Figure 6). Climate forecasting tools and pest monitoring systems are also becoming increasingly important in helping farmers anticipate and respond to pest outbreaks. By integrating weather data with pest population dynamics, farmers can make more informed decisions about when and where to apply pest control measures. This proactive approach is particularly valuable in regions where pests are moving into new territories, as it allows for early detection and rapid response to emerging threats.

VI. Conclusion

Global warming is profoundly altering the geographical distribution and behavior of agricultural pests, leading to significant challenges for global food security. As temperatures rise, pests are expanding into new regions, thriving in climates that were once inhospitable. This shift, combined with longer growing seasons and changes in precipitation patterns, has resulted in increased pest populations, higher reproduction rates, and more frequent infestations, particularly in high-latitude and high-altitude regions. The consequences of these changes are substantial. Crop losses have escalated, with regions like Sub-Saharan Africa and Southeast Asia experiencing significant damage to key staple crops. The economic impact is also severe, as pest-related losses strain already vulnerable agricultural economies. This not only affects farmers but also has broader implications for global food supply chains. Adapting to these challenges requires innovative and sustainable pest management strategies. Integrated Pest Management (IPM), climate-resilient agricultural practices, and early-warning systems for pest outbreaks will be essential in mitigating the effects of these shifts. International cooperation, research, and policy adjustments are critical to addressing the long-term threats posed by global warming to agriculture. The changing distribution of agricultural pests is a pressing issue that must be addressed to ensure future food security in a warming world. Proactive measures and adaptive strategies will be key to managing the evolving risks and minimizing the impact of global warming on agriculture and ecosystems.

References

[1] Tilman D., Balzer C., Hill J., Befort B.L. Global food demand and the sustainable intensification of agriculture. Proc. Natl. Acad. Sci. USA. 2011;108:20260–20264. doi: 10.1073/pnas.1116437108.

[2] Godfray H.C.J., Beddington J.R., Crute I.R., Haddad L., Lawrence D., Muir J.F., Pretty J., Robinson S., Thomas S.M., Toulmin C. Food security: The challenge of feeding 9 billion people. Science. 2010;327:812–818. doi: 10.1126/science.1185383.

[3] Parry M. The potential impact on agriculture of the greenhouse effect. Land Use Policy. 1990;7:109–123. doi: 10.1016/0264-8377(90)90003-H.

[4] Boullis A, Francis F, Verheggen FJ. Climate change and tritrophic interactions: Will modifications to greenhouse gas emissions increase the vulnerability of herbivorous insects to natural enemies? Environmental Entomology. 2015; 44(2): 277-286.

[5] Blomquist GJ, Figueroa-Teran R, Chang E, Tittiger C. Pheromone production in bark beetles. Insect Biochemistry and Molecular Biology. 2010; 40(10):699-712.

[6] Yang, Q.; Zeng, X.; Kuang, W.; Liu, D. Heavy Metal Enrichment and Edible Safety of Two Vegetables in Rare Earth Tailings. Acad. J. Agric. Res. 2016, 4, 268–276.

[7] Schlenker W, Roberts MJ. Nonlinear temperature effects indicate severe damages to US crop yields under climate change. Proceedings of the National Academy of sciences. 2009; 106(37): 15594-15598.

[8] Prakash A., Rao J., Mukherjee A.K., Berliner J., Pokhare S.S., Adak T., Munda S., Shashank P.R. Climate Change: Impact on Crop Pests. Applied Zoologists Research Association (AZRA), Central Rice Research Institute; Odisha, India: 2014.

[9] Rosenzweig C., Major D.C., Demong K., Stanton C., Horton R., Stults M. Managing climate change risks in New York City's water system: Assessment and adaptation planning. Mitig. Adapt. Strat. Glob. Chang. 2007;12:1391–1409. doi: 10.1007/s11027-006-9070-5.

[10] Shrestha S. Effects of climate change in agricultural insect pest. Acta Sci. Agric. 2019;3:74–80. doi: 10.31080/ASAG.2019.03.0727.

[11] Brown PJ, Raymond S, Bradley F, Keimig T. Changes in extreme climate indices for the Northeastern United States, 1870-2005. Journal of Climate. 2010;23:6555-6572

[12] Audu HO, Bibi UM, Garba BG, Ibrahim AK, Hussaini A, Tarki SK, et al. Determination of onset, cessation of rains and hydrological growing season in Dadin Kowa and Gombe for agricultural planning. Nigerian Agricultural Journal. 2019;50(1):133-141

[13] Bekele D, Alamirew T, Kebede A, Zeleke G, Assefa M. Analysis of rainfall trend and variability for agricultural water management in Awash River Basin, Ethiopia. Journal of Water and Climate Change. 2017;8(1):127-141

[14] Kumar V, Jain SK. Trends in seasonal and annual rainfall and rainy days in Kashmir valley in the last century. Quaternary International. 2010;212:64-69

[15] Roshan GHR, Aoji R, Najafi MS, Shakohi E. Prospects for the effect of global warming on the degree-day changes required by wheat for different climates in Iran. Regional Planning Quarterly. 2011;4

[16] Field C.B., Barros V.R., Dokken D.J., Mach K.J., Mastrandrea M.D., Bilir T.E., Chatterjee M., Ebi K.L., Estrada Y.O., Genova R.C., et al. Climate Change 2014: Impacts, Adaptation, and Vulnerability, Part A: Global and Sectoral Aspects; Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press; Cambridge, UK: New York, NY, USA: 2014. IPCC Summary for policymakers; pp. 1–32.

[17] Pachauari R.K., Reisinger A. Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report on Intergovernmental Panel on Climate Change. Intergovernmental Panel on Climate Change (IPCC); Geneva, Switzerland: 2007.

[18] Yoro K.O., Daramola M.O. Paper 1—CO2 emission sources, greenhouse gases, and the global warming effect. In: Rahimpour M.R., Farsi M., Makarem M.A., editors. Advances in Carbon Capture. 1st ed. Woodhead Publishing; Sawston, UK: 2020. pp. 3–28.

[19] Flores-Saavedra, M.; Plazas, M.; Vilanova, S.; Prohens, J.; Gramazio, P. Induction of Water Stress in Major Solanum Crops: A Review on Methodologies and Their Application for Identifying Drought Tolerant Materials. Sci. Hortic. 2023, 318, 112105.