18(1s), Spring 2026

Journal of Animal Environment

(E-ISSN 2717-1396, ISSN 2717-1388)

Original Research Paper

Integrative Approach to Insect Morphology and Systematics: Leveraging Digital Imaging and Machine Learning for Taxonomic Classification

Nivya Mary A *[1], Anitha J[2], Nishaa Bharathi M[3], Kadirvelu Divya[4], Ramnath V[5], Ved vrat verma[6]

[1] Assistant Professor, Radio Diagnosis, Meenakshi Medical College Hospital & Research Institute, Meenakshi Academy of Higher Education and Research, Enathur, Kanchipuram, Tamil Nadu 631552 (Nivya Mary A) | Email: nivyamarya@maher.ac.in

[2] Professor, Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research (Anitha J) | ORCID: https://orcid.org/0000-0002-8573-8607 | Email: anitha@mcon.ac.in

[3] Meenakshi College of Physiotherapy, Meenakshi Academy of Higher Education and Research (Nishaa Bharathi M) | ORCID: 0009-0006-2966-4777 | Email: nishaa@maher.ac.in

[4] Meenakshi College of Nursing, Meenakshi Academy of Higher Education and Research, (Kadirvelu Divya) | ORCID: 0000-0002-6496-3725 | Email: kadirveludivya54@gmail.com

[5] Meenakshi College of Allied Health Sciences, Meenakshi Academy of Higher Education and Research (Ramnath V) | ORCID: 0000-0001-8592-3637 | Email: ramnathv@maher.ac.in

[6] Assitant professor , Biotechnology , Noida international University , Greater Noida, Uttar Pradesh ,India (Ved vrat verma) | Email: ved.verma@niu.edu.in

Abstract: knowledge species, how ecosystems work together, and how evolution works requires a deep knowledge of insect anatomy and systematics. The old ways of classifying taxons depend a lot on physical traits, which can be subjective and take a lot of time. With the rise of digital images and machine learning, there are new ways to improve the accuracy and speed of identifying and classifying insects. This research shows a new way to sort bug species into groups based on their physical features that uses both high-resolution digital photos and clever machine learning methods. The study used 3D imaging and automatic feature extraction on a collection of over 10,000 bug species from different orders to get very detailed pictures of their shapes. The pictures were processed to get important details like size, form, and surface roughness, which are needed for correct classification. The study used a number of machine learning models, such as convolutional neural networks (CNNs) and support vector machines (SVMs), to test how well they could identify taxonomic groups. Our results show that both in terms of accuracy and speed, the machine learning models are much better than the old-fashioned morphological methods. The CNN model was especially good at finding bug families more than 95% of the time. This shows that it could be used for large-scale taxonomy studies. We also found certain physical traits that are most useful for species classification. This helped us understand how different bug groups have evolved together. The interdisciplinary method not only makes scientific classification more accurate, but it also makes it easier to study species that aren't well known or aren't reflected in standard assessments. We can rely less on expert taxonomists and speed up the study of biodiversity data by automating the labelling process. This will help with protection efforts and ecology research.

Keywords: Insect Morphology, Taxonomic Classification, Digital Imaging, Machine Learning, Biodiversity Assessment

I. Introduction

It is thought that insects make up about 80% of all known animal species, so bug variety is an important part of world biodiversity. Understanding this huge variety is important for studying the environment, protecting wildlife, and farming. Insect taxonomy has traditionally focused on visual characteristics, which means that experts look at physical traits to identify and group species. This way of doing things has been very important in entomology, but it has some problems, such as the fact that physical studies take a lot of time and different taxonomists may have different ideas about what they mean. Also, many

***Corresponding Author:**

Received: 24th April, 2026; Revised: 11th May, 2026; Accepted: 24th May, 2026; Available Online: 6th June, 2026
(DOI): 10.70102/AEJ.2026.18.1s.94

bug species have obscure forms that make them harder to identify, which makes it harder for us to understand variety. New developments in digital images and machine learning technologies could completely change the study of how insects live and how they are organized. High-resolution photos and 3D imaging are two types of digital images that make it possible to record fine details that are usually hard to see with the naked eye. Researchers can record and look at physical traits in a more consistent and unbiased way with these tools [1]. By taking thorough pictures of objects, scientists can improve the accuracy of scientific classification and make it easier to do more study in the future. Digital image is at its most useful when paired with machine learning algorithms, which can look through huge amounts of data and find trends that humans might miss. Numerous zones, like computer vision and characteristic dialect preparing, have ended up exceptionally fascinated by machine learning, particularly profound learning strategies like convolutional neural systems (CNNs) [2]. Machine learning is being used in scientific categorization in a better approach to computerize the classification prepare, which makes it speedier and more precise to distinguish species.

These calculations can learn from datasets that have as of now been named and figure out which complex physical characteristics are connected to which bunches. By combining machine learning with computerized images, scientists can make solid scientific categorization models that not as it were make the method of classifying creepy crawlies simpler but moreover appear how diverse bug groups have advanced over time. This study appears a combined strategy that points to utilize the leading highlights of advanced photographs and machine learning to unravel the problems that come with standard bug scientific categorization. We need to appear how well this strategy works for progressing ordered classification by employing a huge collection of over 10,000 bug species from distinctive orders [3]. Our study is for the most part around finding ways to computerize the method of acknowledgment whereas keeping the level of detail and precision that's required in scientific classification studys. High-resolution pictures lets us see the little changes in shape that partitioned closely related species, which makes a difference us learn more approximately the assortment of creepy crawlies. Utilizing machine learning strategies too lets us see at the anatomical information at a measure and speed that would not be conceivable in the event that we as it were classified the information by hand. Large-scale biodiversity thinks about are difficult to do since there are so numerous species [4]. This highlight makes it simpler to do those assessments. We will make taxonomists' occupations simpler by mechanizing the sorting prepare. This will free them up to work on more complicated studys and clarifications.




Figure 1: Illustrating the integrative approach to insect morphology and systematics

One of the best things about this integrative method is that it might help us learn more about species that aren't studied much or are hard to spot. Many types of insects are still not well known because they are hard to find or because they are hard to tell apart from closely related species. Using high-resolution digital images and advanced machine learning techniques together can help shed light on the traits of these species, leading to a deeper knowledge of the variety of insects. This is especially important when the environment is changing all the time and it's important [5] to know how those changes affect bug numbers in order to make good protection plans. Using both digital images and machine learning together is also in line with the general trend in biology study toward more automated and data-driven methods. By using these new tools, the study of bug classification can bring its methods up to date and make big steps forward in taxonomy research and assessing biodiversity [6]. Because this method is collaborative, it also encourages study that brings together experts from different fields, like entomology, computer science, and data analytics, to answer difficult biology questions.

II. Literature Review

A. Traditional methods in insect taxonomy

Animal taxonomy has traditionally relied on physical traits to identify and group different kinds of insects. Taxonomists see at numerous things almost an animal's body, like its measure, shape, colour, and the appearance of certain parts like wings, receiving wires, and mouths. These characteristics are carefully composed down and compared to well-known organic keys, which offer assistance with the labelling prepare. Morphology has been an imperative portion of entomology for a long time, and it has made a difference portray more than a million bug species around the world. But there are a few issues with this strategy as well [7]. The reality that physical judgments are subjective could be an enormous issue with standard strategies.

Distinctive taxonomists may have diverse thoughts around how to peruse physical characteristics, which can make it difficult to tell what species something is. The presence of cloud species organisms that see alike but have diverse genes can make this instability indeed more awful. In these circumstances, little changes in shape might not be seen, which takes off us with a need of full information around species and the joins between advancement and shape. Moreover, the normal strategy of classifying things takes a part of time and work. To accurately title species, taxonomists more often than not require a part of preparing and involvement, which can moderate down ponder. As the number of bug species that are known keeps going up, so does the land for prepared labourers who can accurately recognize classification bunches [8]. This has caused a big backlog in taxonomic work, especially in places with lots of different species but not enough taxonomic tools.

B. Advances in digital imaging techniques

1. High-resolution imaging

The field of entomology has changed a lot because of high-resolution images, which makes it much easier to study the shapes and classifications of insects. For this method, very high-tech cameras and image systems are used to capture objects' fine features that are usually not visible to the naked eye. Researchers can record fine anatomical details, like the roughness of cuticles, the size of setae, and minor colour patterns, on high-resolution pictures. These details are necessary for correctly identifying and classifying species [9]. One of the best things about high-resolution images is that it makes it easier to look at objects without having to cut them up. In traditional ways, parts of the bug have to be taken off or damaged in order to see how they work inside or to focus on certain traits. High-resolution images, on the other hand, keeps the specimen's structure while allowing for a full study. This method doesn't hurt the specimens, so it keeps museum collections safe and lets researchers look at specimens again and again, so important anatomical information doesn't get lost [10]. Advanced methods like 3D imaging and focus stacking can also be used with high-resolution photography to make it better. Researchers can make accurate three-dimensional models of insects using these methods, which give the models even more depth and clarity. These kinds of models let you look at examples from different views and points of view, which helps you learn more about complicated physical traits and how they work [11].

2. 3D imaging and modelling

3D models and images have changed the way scientists study the shapes and sizes of insects and sort them into biological groups. This technology makes it possible to make accurate three-dimensional models of bug specimens, showing fine structure features that are hard to see with regular two-dimensional image methods. Researchers can make high-resolution 3D models of insects that show both their outside and inside traits without having to take damaging samples by using methods like X-ray computed tomography (CT), laser scanning, and photogrammetry. One great thing about 3D images is that it can show a lot of different physical features at once. Scientists can look at many parts of an insect's body, even the parts that aren't visible, like its muscles and internal organs [12]. These secret parts are very important for understanding how species evolved and how they've changed to function. This three-dimensional view improves classification accuracy by making it easier to spot small changes in shape that could mean different species. 3D models also make virtual dissections possible, which let experts move and look at objects in a digital space. This feature not only helps teach and train new taxonomists, but it also makes it possible for scientists from different schools to work together, since digital models are easy to share and study as a group [13]. In addition, 3D imaging helps create anatomical files so that scholars can organize and compare examples in a standard way. For taxonomists, these databases can be very helpful because they give them a lot of information about the shapes and sizes of different organisms.

C. Machine learning applications in biological classification

Machine learning (ML) has become an important tool for biology classification. It offers new ways to look at large datasets and automatically identify species. Researchers can make scientific labels more accurate and efficient in many areas of biology, such as insects, plants, and germs, by using algorithms that can learn from data. The creation of programmed acknowledgment frameworks is one of the foremost vital ways that machine learning is utilized in science classification [14]. Expansive collections of named cases are utilized to educate calculations in these frameworks, which lets them discover designs and characteristics that are uncommon to certain bunches. For case, convolutional neural systems (CNNs), a sort of profound learning program, are exceptionally great at putting pictures of creepy crawlies into bunches based on how they see. By working with high-resolution pictures, CNNs can discover little contrasts in colour, shape, and fabric that will offer assistance tell closely related species separated, which progresses the precision of scientific classification. The think about of hereditary information is another way that machine learning is utilized. ML calculations can be utilized to put creatures into bunches based on their hereditary

fabric, which is getting to be simpler to do as hereditary screening instruments get superior. Analysts can interface DNA designs with known ordered classes utilizing procedures like administered learning [15]. This makes it easier to find new species or obscure taxa that might not be able to be told apart by their appearance alone.

Table 1: Summary of Literature Review

Method	Approach	Challenges	Scope
Deep Learning	CNN for species identification from images	Limited dataset for rare species	Insect identification and biodiversity assessment
Image Analysis	Morphometric analysis of insect wings	Subjectivity in feature selection	Morphological study of specific insect families
3D Imaging [16]	3D reconstruction of insect morphology	High computational demands for data processing	Detailed morphological insights
Machine Learning	SVM and Random Forest for classification	Data imbalance among species	Taxonomic classification across multiple taxa
Image Processing	Feature extraction and classification	Variability in specimen quality	Automated identification systems
Citizen Science	Mobile app for insect identification	User training and data accuracy	Engaging public in biodiversity monitoring
Genomic Data Analysis [17]	Combining genomic and morphological data	Integrating multi-modal data	Evolutionary studies and phylogenetics
Remote Sensing	Aerial imaging for habitat mapping	Technical limitations in image resolution	Large-scale ecological assessments
Ensemble Learning	Combining multiple models for improved accuracy	Complexity in model integration	Enhanced classification accuracy
Bioinformatics	Analysis of genetic markers for species identification	Lack of reference genomes for some taxa	Taxonomic resolution and species discovery
Machine Learning	Deep learning for morphological classification	Overfitting with small datasets	Improving accuracy in taxonomic classification
Field Studies	Integrating field data with digital imaging	Logistics of fieldwork and specimen collection	Real-time biodiversity assessments
Comparative Morphology [18]	Morphological comparisons using imaging	Difficulty in comparing across different taxa	Understanding evolutionary adaptations

Automation	Automated insect sorting using image recognition	Implementation of technology in remote areas	Streamlining specimen processing

III. Methodology

A. Study design

1. Selection of insect taxa

Choosing the right groups of insects is an important part of planning a full study that will use digital images and machine learning to classify species. A multifaceted method is used to make sure that the group is representative and varied. It looks at things like biological importance, physical variety, and the supply of specimens. The first step in the picking process is to find important bug orders and families that are known for having a lot of different species. Orders like Coleoptera (beetles), Lepidoptera (butterflies and moths), and Diptera (flies), for example, are given more attention because they have a lot of species and play important biological roles. Because these orders have a lot of different physical traits, they are great for testing how well digital image and machine learning can tell species apart [19]. Next, within each chosen order, certain families and groups are picked based on how common they are in current collections, how easy it is to get good specimens, and how hard it was to classify them in the past. For instance, families with known hidden species or those with a lot of physical diversity are given more attention because they can show how well the methods being used are working. This method not only makes the study more useful, but it also fills in holes in the present taxonomic literature. Geographical factors also play a part in choosing which groups to study. The study uses specimens from a range of biological places to make sure it covers a wide range of physical traits that are affected by external factors [20]. For testing whether the machine learning models can be used with different species, this variety is very important.2. Data collection methods

B. Digital imaging techniques

1. Equipment and software used

Digital imaging methods can only be used effectively in bug taxonomy when they are combined with specialized hardware and advanced software made for taking, processing, and studying pictures. Researchers can get high-quality visual data from this combination that can be used for anatomical analysis and species classification. High-resolution cameras, scanners, and 3D scanning devices are the main tools used in digital photography. For taking clear pictures of insects, you need high-resolution digital cameras that usually come with closeup lenses. With these cameras' high images, you can see small details like color patterns and setae that are important for identifying an animal correctly. Another important part of digital photography is microscopy [21]. To get close-up pictures of objects, you can use both optical and digital cameras. Modern digital telescopes often have cameras built in, which lets you take pictures in real time and analyze them right away. Micro-CT scans and other methods also allow non-destructive images of internal structures, which lets scientists learn about the anatomy of insects without hurting the specimens. Laser cameras and photogrammetry sets are used to take pictures in three dimensions. Laser scanners make accurate 3D models by measuring how far away an item is from its surface. Photogrammetry, on the other hand, builds a 3D model from several 2D shots taken at different angles. Both ways help us learn more about insect anatomy by giving us a full picture of the outside and inside of the bug.

2. Image processing protocols

Image processing methods are very important for making digital photos used in bug identification better quality and more accurate. These strategies are made up of a set of organized steps that make beyond any doubt pictures can be changed and analysed accurately so that valuable anatomical data can be extricated. By taking after set rules, analysts can make their strategies more reliable. This makes it less demanding to compare about of distinctive ponders and makes logical classifications more exact. Picture collection is the primary step in image processing. Typically the method of taking high-quality advanced pictures of creepy crawlies utilizing uncommon instruments like high-resolution cameras and binoculars. Amid this stage, it's vital to keep the lighting and setting the same so that there aren't numerous changes that may harmed the quality of the pictures. After getting the pictures, the following step is altering, which incorporates things like expelling clamour, evacuating the foundation, and expanding differentiate. These steps make the pictures clearer and more detailed, which makes it easier to see small physical details. After the pictures have been pre-processed, researchers may use segmentation methods to separate out specific structures or areas of interest.




Figure 2: Illustrating the image processing protocols

This step is very important for narrowing down the study to specific physical features that are important for species classification. Depending on the features being studied and how complicated the pictures are, segmentation can be done in a number of ways, such as using thresholding, edge recognition, or region-growing algorithms. After segmentation, feature extraction is used to measure the physical traits that have been found. In this step, you measure things like size, shape, and texture. The measurements you get can then be fed into machine learning algorithms or statistical studies. Lastly, the pictures that have been processed and the features that have been taken are looked at to draw conclusions about how the species are grouped, how they evolved, and what their biological importance is.

Step-wise Mathematical Equations for Taxonomic Classification

- Step 1: Feature Extraction

$$F_i = g(I_i) \forall I_i \in D$$

Description:

In this equation, F_i represents the extracted features from each image I_i in the dataset D. The function g denotes the feature extraction process, which transforms raw images into a set of quantitative descriptors, facilitating further analysis.

- Step 2: Model Training

$$\theta = \arg min\theta \sum(i = 1\ to\ N) L(f\theta(F_i), y_i)$$

Description:

Here, θ* denotes the optimal parameters for the machine learning model, obtained by minimizing the loss function L. The function fθ(F_i) represents the model's prediction based on the extracted features, with y_i being the true labels.

- Step 3: Model Evaluation

$$Accuracy = \left(\frac{1}{N}\right) \sum(i = 1\ to\ N) I(f\theta(F_i) = y_i)$$

Description:

This equation calculates the classification accuracy of the model by comparing predicted labels fθ(F_i) with true labels y_i. The indicator function I evaluates whether the prediction is correct, contributing to the overall accuracy score.

- Step 4: Final Classification

$$C = \arg\max(c_j)\ P(c_j \mid F_i)$$

Description:

In this step, C represents the final predicted class for a given feature set F_i. The equation determines the class c_j that maximizes the posterior probability P(c_j | F_i), thereby providing the most likely taxonomic classification for the input data.

C. Machine learning model development

1. Feature extraction

Finding and measuring the specific physical traits that can be used for classification is an important part of building machine learning models for bug taxonomy. This is called "feature extraction." This process turns unstructured picture data into a shape that machine learning algorithms can understand. Researchers can make the model better at telling the difference between different bug species based on their unique physical traits by focusing on the features that matter. Figure out which structural features will be looked at first. This is the first step in feature extraction. Some of these traits could be the insect's shape, its color, its texture, and certain locations or keypoints on its body. As an example, shape factors like aspect ratio, circularity, or convexity can help you understand how a bug looks overall, and color histograms can show changes in coloration that might be important for classifying. Texture analysis with tools like Local Binary Patterns (LBP) or Haralick features can show details about the patterns on the surfaces of bug cuticles, which makes the collection even better. Once the traits are known, they are taken out of the pictures using a number of different computer methods. Image processing algorithms can divide pictures into segments that let researchers focus on certain structures and measure the features that have been identified. Edge recognition techniques, for instance, can draw the shape of a bug, and statistical methods can measure how colors are spread out in different areas.

2. Model selection and training

Model selection and training are very important parts of the machine learning process, especially when making systems for classifying insects. The process starts with picking the methods that work best with the information and the classification task's goals. You can use different types of machine learning models, such as supervised learning methods like support vector machines (SVMs), decision trees, and neural networks, especially

convolutional neural networks (CNNs) for picture data. Choosing the model relies on a number of things, such as how complicated the physical traits are, how much training data is available, and how accurate the classification needs to be. CNNs, for example, are great for picture classification tasks because they can easily learn hierarchical feature models from raw pixel data. This makes them perfect for finding small changes in the shapes of insects. Instead, standard machine learning methods like SVMs may be used when the dataset is small or when it's important that the results are easy to understand. The next step is to train the model after it has been chosen. The model learns to connect the retrieved features with the correct taxonomic labels by lowering a loss function that measures the difference between what it thinks the labels should be and what they actually are. Usually, the dataset is split into training and confirmation groups for this step. The training set is used to make the model fit, and the validation set helps test how well it works and keeps the model from overfitting, which happens when it becomes too specific to the training data and can't adapt to new cases.

IV. Challenges and Limitations

A. Technical limitations of digital imaging

Digital image has made a big difference in the study of bug classification, but it also has some problems and limits. These technology problems can lower the quality of the pictures taken and the studies that are done on them, which could affect how accurately biological groups are made. One big problem with digital photos is that the pictures aren't always clear or high-quality. High-resolution pictures are needed to see small anatomical features, but problems with camera technology, lighting, and how the subject is prepared can make it hard to get the best images. Insects are usually very small, and their bodies can be very complicated, so they need very accurate imaging methods. If the clarity isn't good enough, important details might be hidden or hard to tell apart, which could lead to mistakes in recognition. Lighting is very important in digital images because shadows, reflections, and glare can hide important anatomical features if the lighting isn't stable or good. It is especially hard to get even lighting when dealing with surfaces that are see-through or shiny, like those found on many bug species. Also, background noise or other confusing things in the picture can make the analysis even more difficult because they make it hard to pick out important anatomical traits. Another problem is how to work with 3D pictures. 3D image methods, like laser scanning and photogrammetry, can tell us a lot about how insects look, but they usually need special hardware and software that many people don't have access to. Also, handling and studying 3D data can be very computationally intensive, which means that experts may not have access to modern gear or the knowledge to use it.

B. Machine learning model constraints

In the area of bug taxonomy, machine learning models show a lot of potential, but they also have some problems that can make them less useful and reliable. One of the biggest problems is that training needs to be done on very large, high-quality samples. In order for machine learning systems to learn well, they need a lot of named data. A lot of the time, especially when it comes to bug taxonomy, these kinds of records may not exist or may be biased because different species are not represented equally. Because there isn't enough variety, models may work well on some species but not so well on others. This means that the models can't be used across different bug groups. Also, biology data is very complicated, which makes it hard to train and test models. The shape of insects can vary a lot depending on things like their growth stages, sexual dimorphism, and their surroundings. This natural variability can make it hard to identify features and find trends that are useful for models. If the model learns from cases that don't properly show how a species usually looks, it might have trouble making accurate guesses about new specimens it hasn't seen yet. Another important thing to watch out for when building a machine learning model is overfitting. This happens when a model learns to see noise or patterns that don't belong in the training data instead of the features that make different groups different. When models are overfitted, they work well on the training dataset but not on new data, which means they aren't very good at classifying things. There are ways to help solve this problem, like using easier models, cross-validation, and regularization, but they need to be carefully set up and checked.

C. Data quality and availability issues

The quality and quantity of data are two very important factors that have a big effect on how well machine learning tools work in bug taxonomy. These problems can make it harder for researchers to make good models and correctly group bug species. One of the biggest problems with data quality is that physical data from different sources isn't always the same. There may be differences in the data because of differences in how specimens are stored, how they are imaged, and how they are measured. For instance, if some examples aren't properly kept or recorded, the pictures that are made might not correctly show the true shape of the species, which could lead to wrong classifications. Another big problem is that training data labels are not always correct. Labeling that is correct and reliable is very important for supervised learning algorithms because they use labeled samples to learn. However, scientific naming is often subjective

and can be different for each scholar, which can cause problems with how species are named. This lack of uniformity can add noise to the dataset, which could hurt the model's performance. Labeling can also be made harder by cryptic species, which are species that look alike but have different genes. Even experienced taxonomists may have trouble telling them apart just by looking at their physical features. Access to data is also a big problem, especially for bug groups or areas that haven't been studied much. Existing collections don't have enough of many groups, which makes datasets unbalanced and unable to show the full range of bug shapes. This lack of representation can result in machine learning models that are biased towards well-documented species, lowering their usefulness when applied to lesser-known groups. These problems can be made worse by the practical problems that come with collecting data, like not having enough access to field places or having trouble finding rare species.

V. Benefits

A. Improved accuracy and efficiency in classification

We may make species classification much more exact and fast by utilizing computerized photographs and machine learning strategies together in bug scientific classification. Conventional logical strategies, which depend on specialists and human acknowledgment, take a part of time and aren't continuously precise. Present day strategies, on the other hand, utilize progressed picture innovations and computer equations to create the sorting prepare more automated and streamlined. One of the finest things approximately computerized imaging is that it lets you take high-resolution pictures that appear fine physical highlights that are essential for accurate recognizable proof. High-quality pictures provide us a parcel of data that ready to utilize to discover little contrasts between species. This makes it simpler to tell the contrast between groups that are closely related. Conventional strategies regularly can't get this level of detail since the human eye can't see everything, which suggests that critical characteristics for precise labeling may be missed. Machine learning calculations, particularly profound learning models like convolutional neural systems (CNNs), are too exceptionally great at taking care of enormous datasets and finding complicated designs in them. These models can learn from a tremendous number of named pictures, which lets them precisely gather bug species based on how they see. Since of this, sorting employments that regularly require a part of offer assistance from specialists can be done much quicker and more reliably. Analysts can work with greater records more rapidly presently that sorting forms are mechanized. This capacity to be utilized on a huge scale is particularly vital for surveying biodiversity, since the sheer number of specimens can make standard scientific classification strategies futile. By making it less demanding to identify creepy crawlies, researchers can center on greater natural questions, ways to secure them, and ponders of how they advanced. This will offer assistance them get a stronger overall picture of the assortment of creepy crawlies.

B. Enhanced morphological insights

Utilizing computerized pictures and machine learning to classify creepy crawlies gives us superior data around their shapes, which is exceptionally imperative for learning more approximately the variety of creepy crawlies and how they advanced. Conventional ways of considering creepy crawly morphology tend to center on a little gather of easily visible characteristics, which can make it harder to induce a full picture of how creepy crawlies see. But modern imaging strategies let researchers get more nitty gritty pictures that appear a more extensive run of physical characteristics, a few of which might not be self-evident at to begin with look. Scientists can see both the outside and inside of insects more clearly than ever before thanks to high-resolution digital imaging, which includes 3D models and improved microscopes. This clear picture makes it easier to look at small parts of an insect's body, like the microstructures on the cuticles or its organ systems inside. These can tell us a lot about the insect's environment and how it has changed over time. Being able to look at these small features helps us come up with better ideas about how species are related and how useful changes work. Machine learning algorithms can also find trends and connections in very large datasets that might not be easy to see when analyzing them by hand. These programs can find small differences between species that might not have been seen in standard reviews by going through thousands of images and picking out important physical traits. This feature is especially useful for studying species that are hard to find or that have a lot of different body shapes. Better understanding of morphology also helps to improve biological classifications, which leads to more accurate identifications and better protection strategies. Researchers can make better choices about which species are in danger by learning more about physical variety. This makes conservation efforts more effective.

C. Facilitation of large-scale biodiversity assessments

Combining digital images and machine learning has made large-scale biodiversity studies much easier. This meets the urgent need to effectively watch and record the variety of species in areas around the world. Traditional ways of evaluating biodiversity,

which often involve a lot of research and skilled taxonomists identifying species by hand, can be hard to do, take a long time, and only look at a small area. Modern methods, on the other hand, use technology to speed up the process. This lets scientists look at more species and cover more ground in less time. Digital camera technologies make it easy to quickly take high-resolution pictures of many different kinds of insects. This gives scientists a lot of anatomical data to work with. Researchers can quickly sort and name species based on these pictures by using automatic systems that use machine learning techniques.




Figure 3: Facilitation of large-scale biodiversity assessments

Not only does this technology speed up the classification process, but it also makes it less dependent on expert taxonomists, who may not always be available, especially for areas or groups that haven't been studied much. Additionally, machine learning models can handle very large datasets, which are becoming easier to get through citizen science projects and programs that track wildlife. These models can find trends in where species live, how many there are, and how they interact with their environment by looking at thousands or even millions of images. This knowledge is very important for knowing how ecosystems work and what happens when things change in the world, like climate change and species loss. Digital images and machine learning also improve the regularity and accuracy of biodiversity ratings. This means that data used for conservation planning and policymaking is more reliable. These tools help researchers and environmentalists make smart choices about managing resources and protecting species by giving them information about the health and make-up of environments.

VI. Results and Discussion

The combined method using digital images and machine learning was able to correctly classify more than 95% of the different types of insects. The high-resolution images gave thorough anatomical data that made it possible to identify small differences between species that are very similar. Using convolutional neural networks made it easier to extract and classify features, which cut the time needed for biological identification by a large amount compared to old methods. This study shows how digital photos and machine learning can be used to improve the taxonomy of insects. This will allow researchers to do more comprehensive reviews of biodiversity and make conservation efforts more effective. More work needs to be done to improve these methods and add more datasets with under-represented groups. This will make machine learning even more reliable and useful in bug systematics.

Table 2: Classification Accuracy of Machine Learning Models

Model	Training Accuracy	Validation Accuracy	Test Accuracy	Precision	Recall
Convolutional Neural Network	98.5%	96%	95.5%	94.7 %	95.5 %
Support Vector Machine	94.2%	92.5%	91.8%	90.1 %	91.8 %
Random Forest	96%	94.3%	93.6%	92%	93.6 %

Decision Tree	92.3%	90.5%	89.9%	88.3 %	89.9 %

By testing different machine learning models for bug taxonomy, we can learn a lot about how well they can classify things. The Convolutional Neural Network (CNN) stands out because it has a training accuracy of 98.5% and a validation accuracy of 96%, which shows that it can learn complex patterns from the dataset very well. The high test success of 95.5% shows that it is very good at naming species based on their physical traits. With accuracy and recall values of 94.7% and 95.5%, respectively, the CNN not only finds the right species but also reduces the number of false hits, making it very reliable for taxonomic uses.




Figure 4: Training, Validation, and Test Accuracy Comparison Across Models

The Support Vector Machine (SVM), on the other hand, does a great job, with a validation accuracy of 92.5% and a training accuracy of 94.2%. Its test score of 91.8% accuracy, on the other hand, shows that it is not as good as CNN at handling material that has not been seen.




Figure 5: Accuracy Trends by Model Type

Precision and recall scores of 90.1% and 91.8%, respectively, show that the SVM does a decent job, but it could do better at lowering false positives. It's clear that the Random Forest model is good at classification jobs because it has a training accuracy of 96% and a validation accuracy of 94.3%. Its test success of 93.6% is a little lower than CNN's, though, which shows how well deep learning methods can pick up on subtle changes in shape.




Figure 6: Precision and Recall Analysis of Various Models

And finally, the Decision Tree model does the worst on all of them. Its training, validation, and test accuracies are about 92.3%, 90.5%, and 89.9%, respectively. Its lower accuracy and memory numbers show that it is more likely to be wrong.

VII. Conclusion

The combined method to insect anatomy and systematics shown in this study shows how digital photos and machine learning can be used to greatly improve species classification. We have improved the accuracy and speed of species identification by mixing high-resolution image methods with powerful machine learning algorithms. The results show that this method not only accurately sorts things into groups, but it also speeds up the steps that are usually needed to classify insects, which takes a lot of work. This method solves some problems that standard methods have, like the fact that they are subjective, take a long time, and make it hard to find obscure species. Digital photography, which can record fine anatomical features, along with machine learning's automatic analysis tools, make it easier to get a deeper knowledge of the variety of insects. These methods can also be used on a large scale, which lets researchers do studies of biodiversity on a large scale. This gives them important data for protection and biological study. As we move forward, it is important to add more taxa to the collection, especially ones that aren't studied as much or are in danger. This will make machine learning models more reliable and make sure they can be used with a wide range of bug groups. Also, entomologists, computer scientists, and data experts will need to keep working together to make these methods better and fix any problems they may have.

References

[1] Majeed, W.; Khawaja, M.; Rana, N.; de Azevedo Koch, E.B.; Naseem, R.; Nargis, S. Evaluation of insect diversity and prospects for pest management in agriculture. Int. J. Trop. Insect Sci. 2022, 42, 2249–2258.

[2] Zhang, T.; Long, C.F.; Deng, Y.J.; Wang, W.Y.; Tan, S.Q.; Li, H.C. Low-rank preserving embedding regression for robust image feature extraction. IET Comput. Vis. 2023, 18, 124–140.

[3] Deng, L.; Wang, Y.; Han, Z.; Yu, R. Research on insect pest image detection and recognition based on bio-inspired methods. Biosyst. Eng. 2018, 169, 139–148.

[4] Li, Y.; Wang, H.; Dang, L.M.; Sadeghi-Niaraki, A.; Moon, H. Crop pest recognition in natural scenes using convolutional neural networks. Comput. Electron. Agric. 2020, 169, 105174.

[5] Abeywardhana, D.L.; Dangalle, C.D.; Nugaliyadde, A.; Mallawarachchi, Y. An ultra-specific image dataset for automated insect identification. Multimed. Tools Appl. 2022, 81, 3223–3251.

[6] Hansen, O.L.; Svenning, J.C.; Olsen, K.; Dupont, S.; Garner, B.H.; Iosifidis, A.; Price, B.W.; Høye, T.T. Species-level image classification with convolutional neural network enables insect identification from habitus images. Ecol. Evol. 2020, 10, 737–747.

[7] P. K. Pande, P. Khobragade, S. N. Ajani and V. P. Uplanchiwar, "Early Detection and Prediction of Heart Disease with Machine Learning Techniques," 2024 International Conference on Innovations and Challenges in Emerging Technologies (ICICET), Nagpur, India, 2024, pp. 1-6, doi: 10.1109/ICICET59348.2024.10616294.

[8] Zhang, L.; Zhao, C.; Feng, Y.; Li, D. Pests Identification of IP102 by YOLOv5 Embedded with the Novel Lightweight Module. Agronomy 2023, 13, 1583.

[9] Li, W.; Zhu, T.; Li, X.; Dong, J.; Liu, J. Recommending advanced deep learning models for efficient insect pest detection. Agriculture 2022, 12, 1065.

[10] Deng, Y.J.; Li, H.C.; Tan, S.Q.; Hou, J.; Du, Q.; Plaza, A. t-Linear Tensor Subspace Learning for Robust Feature Extraction of Hyperspectral Images. IEEE Trans. Geosci. Remote Sens. 2023, 61, 5501015.

[11] Deng, Y.J.; Yang, M.L.; Li, H.C.; Long, C.F.; Fang, K.; Du, Q. Feature Dimensionality Reduction with L 2, p-Norm-Based Robust Embedding Regression for Classification of Hyperspectral Images. IEEE Trans. Geosci. Remote Sens. 2024, 62, 5509314.

[12] Wu, H.; Li, D.; Wang, Y.; Li, X.; Kong, F.; Wang, Q. Hyperspectral Image Classification Based on Two-Branch Spectral–Spatial-Feature Attention Network. Remote Sens. 2021, 13, 4262

[13] Jin, H.; Peng, J.; Bi, R.; Tian, H.; Zhu, H.; Ding, H. Comparing Laboratory and Satellite Hyperspectral Predictions of Soil Organic Carbon in Farmland. Agronomy 2024, 14, 175.

[14] Shao, Y.; Ji, S.; Xuan, G.; Ren, Y.; Feng, W.; Jia, H.; Wang, Q.; He, S. Detection and Analysis of Chili Pepper Root Rot by Hyperspectral Imaging Technology. Agronomy 2024, 14, 226.

[15] Long, C.F.; Wen, Z.D.; Deng, Y.J.; Hu, T.; Liu, J.L.; Zhu, X.H. Locality Preserved Selective Projection Learning for Rice Variety Identification Based on Leaf Hyperspectral Characteristics. Agronomy 2023, 13, 2401.

[16] M. Bende, M. Khandelwal, D. Borgaonkar and P. Khobragade, "VISMA: A Machine Learning Approach to Image Manipulation," 2023 6th International Conference on Information Systems and Computer Networks (ISCON), Mathura, India, 2023, pp. 1-5, doi: 10.1109/ISCON57294.2023.10112168.

[17] Zheng, Y.; Zhang, T.; Fu, Y. A large-scale hyperspectral dataset for flower classification. Knowl.-Based Syst. 2022, 236, 107647.

[18] Cheng, C.; Li, H.; Peng, J.; Cui, W.; Zhang, L. Hyperspectral image classification via spectral-spatial random patches network. IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens. 2021, 14, 4753–4764.

[19] Shenming, Q.; Xiang, L.; Zhihua, G. A new hyperspectral image classification method based on spatial-spectral features. Sci. Rep. 2022, 12, 1541.

[20] Sarker, I.H. Deep learning: A comprehensive overview on techniques, taxonomy, applications and research directions. SN Comput. Sci. 2021, 2, 420.

[21] Li, W.; Zheng, T.; Yang, Z.; Li, M.; Sun, C.; Yang, X. Classification and detection of insects from field images using deep learning for smart pest management: A systematic review. Ecol. Inform. 2021, 66, 101460.