



Original Research Paper/Review Paper

Deep Learning Prediction of Reptile Diversity in Semi-Arid Ecosystems

Dr. Priya Paul^{1*}, Dr. Pokhraj Sahu²

^{1*}Assistant Professor, Kalinga University, Naya Raipur, Chhattisgarh, India.
Email: ku.priyapaul@kalingauniversity.ac.in, ORCID: <https://orcid.org/0009-0008-7934-9307>

²Assistant Professor, Kalinga University, Naya Raipur, Chhattisgarh, India.
Email: ku.pokhraj_sahu@kalingauniversity.ac.in, ORCID: <https://orcid.org/0009-0007-7535-0848>

Key Words

Abstract

Semi-arid ecosystem, Reptile diversity, Deep learning, Species distribution modeling, Habitat suitability, Herpetofauna.

Introduction: Semi-arid environments host an unusually diverse range of specialized reptiles that thrive in environments with extreme temperatures, vegetation and variable precipitation, but quantifying this biodiversity remains a challenging task due to the elusive nature of reptiles, uneven distributions and high cost associated with sampling on the arid environment. Methodology: In this paper, the utilization of deep learning (DL) for predicting reptile biodiversity in semi-arid environments is theoretically synthesized. DL models based on image-based convolutional neural networks, deep regression for species richness prediction using bioclimatic variables and species distribution modeling approaches are discussed, and an analytical framework is proposed. Conclusion: Deep learning has provided a conceptual framework that can be used for reptile diversity prediction across semi-arid ecosystems, provided that occurrence data are sufficiently available, geographical bias is addressed, and appropriate aridity-related covariates are included in the model. Results: From the analysis, we note that deep learning models that have been trained on camera trap and citizen science images have the potential to provide high accuracy species identification for snakes and lizards, that non-linear richness gradients can be captured by deep regression and convolution models when used with bioclimatic and remote sensing covariates, and that model performance is heavily dependent on aridity-related covariates.

* Corresponding Author's email: ku.priyapaul@kalingauniversity.ac.in

Received: 25th Aug, 2026 | Revised: 1st Sep, 2026 | Accepted: 5th Sep, 2026 | Available Online: 14th Sep, 2026

(DOI): 10.70102/AEJ.2026.18.5s.3

Introduction

Semi-arid systems, which occur over broad swaths of North Africa, the Arabian Peninsula, the Sahel, and similar arid areas across the globe, house reptile faunas that are relatively rich for the region when compared to the rest of the region's vertebrates. The low water availability, large thermal variation, and open vegetation create environments that are suitable for ectotherms that can burrow and are nocturnal. Due to the nature of reptiles being cryptic, small, and thermally specific, these animals are often overlooked when compared with birds and mammals, and current efforts to survey are unable to match the scale of semi-arid environments that are vulnerable to climate change and land-use pressures. This paper provides a potential means to bridging the assessment gap through species recognition via image processing and modeling of how diversity patterns correspond to bioclimatic and remote-sensing covariates in large-scale areas. The paper is entirely theoretical and conceptual, and thus it will not contain any empirical information, modeling, nor estimation of any parameters in equations. It simply integrates current knowledge about the ecology of reptiles in arid zones and the use of deep learning for species recognition and modeling of their distribution into one conceptual framework (Aldosari & Huda, 2024).

Review of Literature

A great deal of foundational research exists regarding the influence of climate and habitat variables on reptile distributions in arid zones. The study by (El-Gabbas et al., 2016) utilized

species distribution models (Maxent) for the projections of current and future distributions of seventy-five Egyptian reptile species in the conditions of climate change and determined that the contraction of ranges could be expected for a significant portion of the studied assemblage. (Sow et al., 2014) biogeography of Atlantic Sahara reptiles revealed that aridity and temperature variables are the main correlates of the distribution and vulnerability of the reptiles to climate change. (Alatawi et al., 2020) developed the first ever models of the richness and habitat suitability of terrestrial reptile species for Saudi Arabia.

Another related set of research studies has been done in the context of deep learning for herpetological identification and, more broadly, species distribution modeling (Binta et al., 2023) that showed convolutional neural networks can be used to classify snakes, lizards, and toads from camera trap images with class imbalances in natural data. (Ahmed et al., 2024; Othman et al., 2021) have proven that convolutional neural networks can be used to get high accuracy in fine-grained, image-based identification of snake species, which is quite a challenging task for non-specialists because of the very subtle differences in morphological characteristics between species. (Botella et al., 2018) have shown that using neural networks with deep architecture, convolution, and multiple labels can help surpass classical methods of environmental niche modeling for species distribution forecasting, especially when the realized niche in environmental space is not regular (Selvam & Devi, 2025; Brugere et al., 2023) have shown that

neural network and random forest modeling can surpass classical generalized linear models for predicting species richness based on environmental variables. As a whole, however, this literature suggests that the climate-controlled distribution of desert reptiles is relatively well documented, and that deep-learning techniques have become increasingly proficient at automatically identifying reptile species as well as predicting their richness; yet what is relatively underdeveloped at the moment is an integrative theoretical approach to this problem.

Paper Organization and Key Contributions

The rest of this research paper is structured as follows. The Methodology chapter explains the methodology that combines an environmental and presence dataset with deep learning techniques and diversity outcomes. This analysis is shown through a schematic diagram without any equations. The Results and Discussion chapter reviews the strengths and limitations of the major deep learning methodologies employed in this area and shows their comparative summary table, as well as explains the relationship between the aridity covariates and occurrence data and the performance of models.

The major contributions of this study include: (i) integration of arid lands herpetological and deep learning literature into a unified framework for predicting reptile diversity; (ii) a straightforward analysis process that does not require any formulae and is relevant to empirical research on linking bioclimatic variables with occurrences or images in future; (iii) a

comparative review of deep learning image-based, regression-based, and transfer learning methods in this field; and (iv) determination of sampling bias and covariate selection in semi-arid regions as the most important issues to consider in future model-based assessments of reptile diversity.

Methodology

The present study utilizes a theory-driven, literature review-based approach in which no data collection takes place and no statistical or mathematical modeling has been done. The process flowchart generated based on the conceptual framework presented in the literature divides the analysis process into four phases, as presented in Figure 1 below. In the first phase, environmental data for each site of interest are collected including information on temperature and rainfall patterns, the index of aridity, vegetation coverage, and soil type that define the semi-arid gradient. In the second phase, occurrence data are collected which includes camera-trap data, museum records, citizen-science records, and transect records.

The third stage is featuring construction, which uses the bioclimatic and habitat descriptors from the environmental data and occurrence and remote sensing data respectively to provide input for modeling. The fourth stage employs the deep learning approach, leveraging the classes discussed in the literature review which include image based convolutional neural networks, deep regression networks, deep species distribution models, and transfer learning methods to provide species level identification

and outputs related to diversity including species richness maps and habitat suitability maps. This being a theoretical paper, no particular neural network is trained in this paper.




Figure 1: Conceptual Workflow for Deep-Learning-Based Prediction of Reptile Diversity

Results and Discussion

Since this paper has an academic approach, results are not obtained through experimentation; therefore, the results provided below are a combination of results obtained from existing

research literature. Table 1 below shows the type of input data used by various categories of deep learning approaches, together with their strengths and weaknesses in prediction of diversity of reptiles in semi-arid regions.

Table 1: Comparative Overview of Deep Learning Approaches Reported for Reptile Diversity Prediction

Approach	Typical Input Data	Reported Strength	Reported Limitation
Convolutional neural networks (image-based)	Camera-trap or field photographs of snakes and lizards	High species-level accuracy on large labelled image sets	Struggles with cryptic, camouflaged, or rare desert species
Deep regression networks	Bioclimatic and remote-sensing covariates per grid cell	Captures non-linear richness-aridity relationships	Needs dense environmental covariate coverage
Deep species distribution models	Occurrence records with environmental predictors	Improves on classical Maxent for irregular niches	Sensitive to sampling bias in sparse arid records
Transfer learning approaches	Pre-trained image or environmental feature models	Reduces labelled-data requirements for rare species	Source-domain mismatch can limit desert applicability

From the discussion above, several themes have been derived. First, there is high accuracy in image-based convolutional neural networks when used to identify animal species if they have been fed a large enough dataset, although they have challenges identifying cryptic, camouflaged, or nocturnal desert animals. Additionally, deep neural network-based methods can model complex richness-aridity relationships, as compared to classical linear and

generalized additive models, by fitting non-linearities in bioclimatic and remote sensing environmental covariates, which is especially important considering the existence of extreme environmental gradients in semi-arid environments; however, this is contingent on dense sampling of environmental covariates. Third, transfer learning methods lessen the need for labeled data for endangered or range-restricted species through transfer of features

obtained from other tasks, although their effectiveness is limited when the source domain is very different from the arid zone.

Sampling bias further exacerbates the above trade-offs. Occurrence data on reptiles in semi-arid environments tends to be aggregated along accessible road networks, settlements, and conservation sites, suggesting that naïve modeling on such data would amount to sampling effort rather than habitat suitability. Also, evidence provided in (El-Gabbas et al., 2016; Sow et al., 2014) indicates that species richness and distribution are based on covariates relating to certain climatic conditions, for instance, seasonal temperature variation and vegetation, and not generic dryness measures, implying that covariate selection should be done based on the ecological gradient considered.

Conclusion

The diversity of reptiles in the semi-arid ecosystem is influenced by the presence of significant climatic and habitat gradients, and comprehensive evaluation of such a diversity has, for long, been limited by the secretive and fragmented distribution of reptiles in the arid zones. Ecological studies of reptiles in the arid zones and the development of deep learning-based methods for species classification and distribution modeling are, to a large extent, independent areas of research. This paper, however, on a theoretical basis, proposes the unification of both within one workflow. The realization of this potential through research will involve addressing issues of sampling bias within occurrence data, the choice of covariates that

reflect aridity, and having sufficient labelled imagery for rare or cryptic species. These are all natural topics for empirical investigation in the future based on the current framework proposed.

References

- [1] Ahmed, Kareem, Mai A. Gad, and Amal Elsayed Aboutabl. "Snake species classification using deep learning techniques." *Multimedia Tools and Applications* 83, no. 12 (2024): 35117-35158.
- [2] Alatawi, Abdulaziz S., Francis Gilbert, and Tom Reader. "Modelling terrestrial reptile species richness, distributions and habitat suitability in Saudi Arabia." *Journal of Arid Environments* 178 (2020): 104153.
- [3] Aldosari, Huda Mohammed. "An Expert Model Using Deep Learning for Image-based Pest Identification with the TSLM Approach for Enhancing Precision Farming." *J. Wirel. Mob. Networks Ubiquitous Comput. Dependable Appl.* 15, no. 3 (2024): 160-183. <https://doi.org/10.58346/JOWUA.2024.13.012>
- [4] Binta Islam, Sazida, Damian Valles, Toby J. Hibbitts, Wade A. Ryberg, Danielle K. Walkup, and Michael RJ Forstner. "Animal species recognition with deep convolutional neural networks from ecological camera trap images." *Animals* 13, no. 9 (2023): 1526.
- [5] Botella, Christophe, Alexis Joly, Pierre Bonnet, Pascal Monestiez, and François

- Munoz. "A deep learning approach to species distribution modelling." In *Multimedia tools and applications for environmental & biodiversity informatics*, pp. 169-199. Cham: Springer International Publishing, 2018.
- [6] Brugere, Lian, Youngsang Kwon, Amy E. Frazier, and Peter Kedron. "Improved prediction of tree species richness and interpretability of environmental drivers using a machine learning approach." *Forest Ecology and Management* 539 (2023): 120972.
- [7] El-Gabbas, Ahmed, Sherif Baha El Din, Samy Zalat, and Francis Gilbert. "Conserving Egypt's reptiles under climate change." *Journal of Arid Environments* 127 (2016): 211-221.
- [8] Othman, Zainab, Nur Nabilah Abu Mansor, Nur Farhani Azmi, Nurul Hidayah Mat Zain, Khyrina Airin Fariza Abu Samah, Ismassabah Ismai, and Khairul Adilah Ahmad. "Snake species identification using digital image processing." In *2021 6th IEEE International Conference on Recent Advances and Innovations in Engineering (ICRAIE)*, vol. 6, pp. 1-6. IEEE, 2021.
- [9] Selvam, R. P., and R. Devi. "A Comparative Stratification of Fish Species Using Transfer Learning on Pre-Trained Deep Learning Networks Juxtaposed with Shuffleres-A Hybrid Deep Network Classifier." *Archives for Technical Sciences/Arhiv za Tehnicke Nauke* 34 (2025): 157. <https://doi.org/10.70102/afts.2025.1834.157>
- [10] Sow, Andack Saad, Fernando Martínez-Freiría, Hamidou Dieng, Soumia Fahd, and José Carlos Brito. "Biogeographical analysis of the Atlantic Sahara reptiles: Environmental correlates of species distribution and vulnerability to climate change." *Journal of Arid Environments* 109 (2014): 65-73.