



Original Research Paper

Integrated Approaches to Increase Potatoes' Productivity, Quality and Disease Resistance Under Variable Climate Conditions - A Review

Lokendra Kumar¹, Dr. Avadhesh Kumar Koshal², Dr. Deepali Singh³, Samiksha Parmar⁴, Harshit Sharma⁵, Dr. Prem Kumar⁶, Dr. Ram Ashish⁷, Priyanka Kumari⁸, Durgesh Kumar Maurya⁹, Sanya Bhasin¹⁰

¹Assistant Professor, Faculty of Agriculture, Motherhood University, Roorkee, Uttarakhand, India

²Professor, Faculty of Science, Motherhood University, Roorkee, Haridwar, Uttarakhand, India

³Assistant Professor, Faculty of Agriculture, Motherhood University, Roorkee, Uttarakhand, India

⁴Research Scholar, Department of Botany, Motherhood University, Roorkee, Haridwar, Uttarakhand, India

⁵Assistant Professor, Department of Agriculture, Bhagwant Global University, Kotdwar, India

⁶Teaching Associate, College of Horticulture, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, India

⁷Assistant Professor, School of Agriculture Science and Engineering, IFTM University, Moradabad, Uttar Pradesh - 244102, India

⁸Ph.D. Scholar, Department of Horticulture and Post-Harvest Technology, Institute of Agriculture, Visva-Bharati, Sriniketan-731236, West Bengal, India

⁹SMS Agronomy, KVK, Santkabirnagar (ANDUAT, Kumarganj, Ayodhya), India

¹⁰Department of Agriculture, Lab Technician (Genetics and Plant Breeding), Maharishi Markandeshwar Deemed to be University, Mullana - Ambala, Haryana, India

Abstract

The potato (*Solanum tuberosum* L.) is among the most important foods produced around the world and an essential source of carbohydrate, vitamins, minerals, and incomes for millions of farmers. Nonetheless, potato yields, tubers' quality, and disease resistance are increasingly at risk due to climate variability, such as increased temperatures, erratic precipitation, droughts, floods, and pests/pathogens development. To counteract these problems, it is necessary to implement climate-resilient technologies for integrated potato production that will secure its cultivation and ensure food security across the globe. This review highlights recent achievements in integrated technology used to increase potato productivity, tubers' quality, and resistance to diseases due to climate changes. The review focuses on climate-resilient and improved potato varieties, nutrient management, irrigation, conservation agriculture, crop rotations, integrated pest and disease management (IPDM), biological agents of protection, and the use of digital agriculture technologies such as remote sensing, artificial intelligence, and precision agriculture. There is special attention to the role of genomics, marker-assisted selection, genome editing, and micro biome technology in development of resilient potato varieties with improved tolerance to biotic and abiotic stress factors. Besides, there is also attention to post-harvest handling, seed quality improvement and soil health practices as essential components in sustaining the quality of tubers and minimizing production losses. It is noted that integration of agronomy, horticulture, plant breeding, plant pathology, soil sciences, and climate-smart agriculture practices is more beneficial than implementation of technologies separately. Such integration helps not only increase resource-use efficiency, yield stability, nutritional quality and sustainability, but also makes potato production systems more resilient to climatic changes. Future research should be aimed at multidisciplinary approach, decision-support systems and innovations for farmers. The review offers a number of important insights for researchers, extension agents, policymakers, and potato growers interested in sustainable use of climate-resilient technologies in potato production.

Keywords Potato (*Solanum tuberosum* L.); Climate change; Abiotic stress; Biotic stress; Integrated crop management; Precision farming; Disease resistance; Late blight; Early blight; IDM; Plant breeding; Molecular breeding; Biotechnology; Soil health; Nutrient management; Water management; Plant growth promoting rhizobacteria (PGPR).

1. Introduction

Potato (*Solanum tuberosum* L.), the third most important food crop in the world following rice and wheat, provides a rich source of carbohydrates, vitamins, minerals, dietary fiber, and antioxidants for millions of people worldwide. This crop is grown in over 150 countries in temperate, subtropical, and tropical zones. Potato is an essential crop because of its high yield potential and crop period in commercial as well as subsistence farming systems. In countries like India, China, Russia, Ukraine, the United States, and some European countries, potato growing is a crucial component of increasing income of farmers,

employment generation, and agro-industrialization. Nonetheless, potato cultivation has faced threats due to the changing climatic conditions that include high temperatures, erratic rainfall, drought periods, heat waves, floods, and weather extremes. Such climatic changes directly affect crop growth, photosynthesis, initiation, bulking, and maturation, causing yield reduction and reduced marketability of tubers. Increased temperatures cause high plant respiration rates, low tuber set, and more physiological problems, like heat sprouts, hollow heart, and secondary growth. Water stress, irrespective of being due to either drought or excess rain, affects nutrient uptake, root growth, and tuber uniformity.

*Corresponding Author:

Received: 25th July, 2026 | Revised: 26th August, 2026 | Accepted: 18th September, 2026 | Available Online: 24th September, 2026

(DOI): 10.70102/AEJ.2026.18.7s.36

Climatic variability also leads to increased chances of occurrence and spread of pathogens and insects. Some of the important diseases include late blight (*Phytophthora infestans*), early blight (*Alternaria solani*), black scurf (*Rhizoctonia solani*), common scab (*Streptomyces scabies*), bacterial wilt (*Ralstonia solanacearum*), soft rot, viruses, and nematodes. At the same time, insects and pathogens are adapting to the increased temperatures, making control measures difficult and costly, involving increased use of chemical pesticides. Such climatic changes directly affect crop growth, photosynthesis, initiation, bulking, and maturation, causing yield reduction and reduced marketability of tubers. Increased temperatures cause high plant respiration rates, low tuber set, and more physiological problems, like heat sprouts, hollow heart, and secondary growth. Water stress, irrespective of being due to either drought or excess rain, affects nutrient uptake, root growth, and tuber uniformity. Climatic variability also leads to increased chances of occurrence and spread of pathogens and insects. Some of the important diseases include late blight (*Phytophthora infestans*), early blight (*Alternaria solani*), black scurf (*Rhizoctonia solani*), common scab (*Streptomyces scabies*), bacterial wilt (*Ralstonia solanacearum*), soft rot, viruses, and nematodes. At the same time, insects and pathogens are adapting to the increased temperatures, making control measures difficult and costly, involving increased use of chemical pesticides. In order to solve these issues, holistic approaches based on advanced agronomical practices, climate-smart agriculture, proper irrigation and nutrient management, precision farming techniques, cultivation of resistant varieties, use of biological control agents, IDM and advanced breeding methods are assuming great importance. The recent development in genomics, marker assisted selection, genome editing, remote sensing, artificial intelligence and digital agriculture have opened up new possibilities for improving the productivity and quality of potatoes as well as resistance to biotic and abiotic stress. Moreover, sustainable soil management, conservation agriculture, diversification of crops, bio-fertilization and plant growth promoting microorganisms play important role in keeping crop resilient to changing climates. This review is intended to provide a thorough analysis of the advancements made in an integrated approach towards increasing the yield, quality, and disease resistance of potatoes in varying climatic conditions. It discusses the importance of climate-smart agronomy, genetics, precision farming, biology, integrated disease management, and technology in creating sustainable and resilient potato production systems. This review also discusses the gaps that exist in the current research along with the future prospects for making potato production resilient to climate changes while maintaining productivity and sustainability. The present review seeks to critically analyze the recent developments in integrated approach towards enhancement of productivity, quality, and disease resistance of potatoes under varying climatic conditions. It emphasizes the importance of climate-smart agronomy, genetic modification, precision agriculture, biological approaches, disease management, and innovative technologies for creating a sustainable and resilient potato production system. The review also pinpoints the existing gaps in the field of research and discusses the future perspectives of increasing climate resilience with productivity and sustainability of the potato production system.

2. Worldwide Importance of Potato Cultivation

The potato (*Solanum tuberosum* L.), which is the third most important food crop in the world, after rice and wheat, in terms of

Journal of Animal Environment, Vol. 18, No. 7s, Spring 2026

human consumption, is essential in terms of global food security and nutrition. The potato is cultivated in more than 150 countries in temperate, subtropical, and tropical areas, occupying 17–19 million hectares of land annually and producing over 370 million tonnes of potatoes. Thanks to high productivity per hectare and short cycle of cultivation, potato became one of the most productive crops for feeding the growing world population. Potato is rich in carbohydrates, dietary fiber, vitamin C, vitamin B6, potassium, and antioxidants. Potato gives more energy and protein from a unit volume of water compared to other cereal crops. Therefore, it is one of the key elements of sustainable agriculture. Besides being a staple crop, potatoes are widely processed into chips, French fries, flakes, starch, flour, and alcohol. The crop is of considerable economic value as well. Millions of subsistence and large scale farmers earn their livelihood from potato farming as it has high market value and also gives quick economic returns. The leading nations where potato is grown include China, India, Ukraine, Russia, USA, Germany, Bangladesh, France, Pakistan, and the Netherlands. India is the second largest producer of potatoes in the world. In spite of its importance in the world, growing potatoes faces several problems that include changing climatic conditions like increasing temperature, unpredictable rains, drought, floods, pests, and diseases. This poses threat to potato yields, their quality, storage period, and their profitability. Hence, increasing the productivity of potatoes, improving their nutrition value, and disease resistance have become a major area of research in the world. Production of potatoes in the future is going to rely on climate-smart technologies, new breeding techniques, water and nutrient management, and disease management strategies. These will make potatoes more resilient to changes in environmental conditions, and at the same time guarantee sustainable production, better food security, and greater farmer revenues. Potatoes can thus be considered an essential crop for ensuring sustainable agriculture on a global scale.

3. Effects of Climate Variability on Potato Production and Quality

Potato (*Solanum tuberosum* L.) is one of the most important food crops worldwide; however, potato productivity and tuber quality are very vulnerable to changes in the climate. Shifts in temperature, rainfall patterns, humidity, solar radiation, and occurrence of climatic extremes have had a great impact on potato production in temperate and tropical areas. Climate variability impacts all stages of crop development, from sprouting to germination, vegetative development, tuber set, tuber growth, and maturation, ultimately resulting in reduced yield and quality. Among climatic factors that affect potato production, temperature is the most significant one. Optimal temperature regime for tuberization ranges between 15–20°C; however, temperatures exceeding 25°C prevent tuber initiation and stimulate excessive vegetative growth. Heat stress reduces crop development time, causes premature plant senescence, lowers photosynthesis efficiency, and impairs carbohydrate transfer from the leaves to tubers. This results in smaller-sized tubers, non-uniform in shape, with reduced dry matter content. Moreover, heat stress contributes to physiological disorders like heat necrosis, hollow heart, and secondary growth, lowering potato process ability. Water scarcity and drought remain key constraints in potato farming. Insufficient water during tuber initiation and bulking phases results in reduced cell enlargement and therefore production of few tubers. Water scarcity results in nutrient deficiency, inhibition of photosynthesis process, and low water use efficiency, which greatly reduce yield. On the other

hand, excess rains coupled with waterlogged soils create anaerobic conditions in the soils that limit root respiration and nutrient uptake as well as tuber rot infections. Waterlogged soils encourage growth of soil borne diseases such as black scurf and bacterial soft rot. Weather fluctuations further affect the occurrence and level of infection from insects and diseases. High temperatures and humidity provide good conditions for pathogens such as late blight (*Phytophthora infestans*), early blight (*Alternaria solani*), and *Rhizoctonia solani*. Some insects including aphids, whiteflies, and potato tuber moth develop higher number of generations in warm weather conditions. The quality of the potato tubers is further compromised due to climatic variability. Heat and drought lead to reduced starch concentration, low dry matter, and high reducing sugar content in potatoes which compromise their fry quality and dark color of chips. Excessively wet climate can cause low specific gravity, higher cracking, and bruising. Climate stress will further reduce the concentration of vitamin C and composition of nutrients in tubers. Climate variability can cause further degradation of the soil health through increased erosion, nutrient leaching, soil salinization, and soil moisture variability. This would reduce fertilizer use efficiency and further cause harm to root growth and microbial processes. This makes maintenance of soil fertility and moisture level increasingly difficult in varying climatic conditions. In order to reduce adverse effects of climate variability on potato production, there should be adoption of climate-smart farming practices. These include use of heat, drought and disease-resistant cultivars, efficient irrigation techniques like drip irrigation, mulching, proper nutrient management, crop rotation, precision farming, weather-based forecasting of diseases and pests, and integrated disease and pest management. The use of conservation agriculture and resilient cultivars bred via conventional and molecular techniques could improve adaptation to varying climatic conditions.

4. Determinants of Potatoes Productivity and Quality Productivity and quality of potato (*Solanum tuberosum* L.) tubers are controlled by various genetic, environmental, agronomic and biotic determinants. Optimum production of potato tubers that have excellent quality necessitates the optimization of all these determinants at every stage of growth. The importance of each determinant changes with the genotype of potatoes, the period of growth, soil types and climate conditions.

A. Genetic Factors-Genetics of a potato variety determines its productivity level, tuber size and shape, color, dry matter content, starch level, disease and pest resistance level. Improved varieties of potatoes with high productivity potential, heat and drought tolerance and disease resistance like late blight and viruses have higher productivity and quality of tubers. Processing potato varieties have high dry matter and low reducing sugar content for making chips and French fries.

B. Climatic Factors-Potato is a crop that thrives well in cool climatic conditions within 15-20°C for tuber formation. Higher temperature inhibits tuber formation, increases respiration rate, and reduces dry matter accumulation, resulting in poor quality and low yields. Frost, erratic rains, droughts, and heat waves are also unfriendly climatic factors affecting growth, photosynthesis, and formation of tubers. Solar radiation and humidity are needed for maximum productivity.

C. Soil Factors-Potatoes thrive well in well-drained sandy loam to loamy soils with a pH of 5.5-6.5. The fertility level, structure, moisture holding capacity, aeration of the soil and organic matter content greatly affects the development of roots and tubers. In compacted and waterlogged soils, the roots fail to expand hence the incidence of diseases and low marketable yield.

D. Nutrient

Management-Fertilizer application plays an important role in ensuring high productivity and quality of the tubers produced. Excess nitrogen results in vegetative growth but it delays the maturity of the tubers and reduces dry matter content. Phosphorus is needed for the development of the roots and tubers at an early stage. Potassium increases starch content and enhances size of the tubers, resistance to diseases and storage quality.

E. Water Management-Appropriate irrigation scheduling is very important all through the growth cycle of the crop, especially during the initiation and bulking phases. Water shortage as well as excess moisture leads to decrease in tuber size and quality. Water shortage causes reduction in tuber size and an increase in tuber deformities. Excess water results in water logging, leaching of nutrients, and an increased risk of diseases.

F. Agricultural Practices - Optimal farming practices such as optimal planting time, plant spacing, size of seed tubers, planting depth, weed control, earthing up, crop rotation, and timely harvesting affect potato yield and quality. Certification of the seed tubers and precise farming practices improve yield and quality of potatoes.

G. Biotic Factors -Diseases such as late blight, early blight, black scurf, bacterial wilt, and viruses decrease potato yield and marketability. Pests such as aphids, potato tuber moth, white grubs, and cutworms destroy the foliage and tubers. Integrated management practices are required for minimizing the loss of production.

H. Agronomic Practices - Optimum practices, such as planting time, recommended spacing between plants, size of seed tuber, planting depth, weeding, earthing up, crop rotation, and harvest time, affect the yield of potatoes. Use of certified seed tubers and precision agriculture practices increase yield and quality of potatoes.

I. Biotic Factors - The occurrence of several diseases, including late blight, early blight, black scurf, bacterial wilt, and virus infections, causes poor yield and low marketable nature of potatoes. Pest infestations from insect pests like aphids, potato tuber moth, white grubs, and cutworms cause leaf and tuber damage. It is important to adopt pest and disease management practices for increasing yield and quality.

J. Physiological and Post-harvest Factors -Physiological problems such as hollow heart, greening, cracking, growth cracks, and black spot decrease the market value of potatoes. Recommended practices of harvest, curing, grading, packaging, and proper storage conditions are required to maintain potato quality, reduce weight loss, and increase shelf life.

5. Integrated Agronomic Techniques for Enhancing Potato Yields

Integrated agronomic techniques refer to the integration of improved crop management practices for achieving maximum potato yield along with soil health, efficient utilization of resources, and sustainability of environment. For attaining optimum potato yield, quality, and reducing risk factors under varying climatic conditions, it is necessary to adopt an integrated approach in agronomy.

Variety Selection-Selection of varieties having high yield, resistance to various diseases, and capable of adapting themselves to the prevailing climatic conditions becomes imperative for obtaining increased productivity from potatoes. Varieties which can withstand heat, drought, and various diseases, including late blight, have greater chances of survival under varying climatic conditions. Use of healthy seed potatoes helps ensure successful cultivation.

Optimum Planting and Seed Management-Planting at optimum times ensures that the crops make best use of favorable temperatures and moisture levels for their optimal growth. Seed potatoes that are healthy and of medium size (30-50 grams) and have been treated with

proper fungicides or biological agents before planting help enhance germination and increase the vigor of plants. **Balance Nutrient Management-** It is imperative to apply balanced proportions of macro-nutrients (Nitrogen, Phosphorus, Potassium) as well as secondary and micro-nutrients for achieving high yields of potato and maintaining its quality standards. The practice of INM includes the use of chemical fertilizers, farm yard manure (FYM), compost, green manures and biofertilizers. **Effective Water Management-**Potatoes need sufficient amounts of moisture in the soil especially during tuber initiation and bulking. Irrigation can be planned on the basis of water requirements of the crop as well as the status of soil moisture. Practices of drip irrigation, sprinkler irrigation and mulching increase the efficiency of water utilization. **Soil Health Management-**Healthy soil is very crucial in sustainable production of potatoes. Organic matter, crop residues, compost and biochar increase the healthiness of the soil as they improve its structure, water-holding capacity, biological activities and nutrient levels. Rotation of crops with legumes and other non-host crops also decreases soil diseases and increases fertility. **Weed Management-**Weeds compete with the crop for nutrients, moisture, light and space. This results in reduction of the crop productivity. Integrated weed management includes use of cultural practices, mechanical weeding, mulching and selective herbicides. **Precision and Climate-Smart Agriculture-**Modern precision technologies like nutrient management, moisture sensors in soils, remote sensing techniques, GPS-controlled equipment, and decision support systems improve efficiency in using inputs and minimize costs. Climate-smart agricultural practices enable farmers to adapt to changes in climatic conditions and increase productivity with minimum environmental degradation. **Integrated Pest and Disease Management (IPDM)-**The integration of pest and disease-resistant cultivars, crop rotation, biological pest control methods, fertilization, sanitation, and rational use of chemical pesticides is an effective way of managing pests and diseases. IPDM decreases the losses due to pests and diseases, delays the development of pesticide resistance, and guarantees environmentally sustainable potato production. **Mechanization and Better Crop Management-** Mechanization of potato production through the use of advanced equipment for tillage, planting, hilling, watering, spraying, and harvesting minimizes labor inputs, decreases crop losses, and increases efficiency of fieldwork.

6. Role of Horticulture Practices in Improving Potato Tuber Quality

Horticulture practices have an important role in improving the physical, nutritional, processing, and storage qualities of potato tubers. Effective management of crop during its planting and harvesting makes possible production of uniform, healthy, and marketable potato tubers having high consumer acceptability. Implementation of scientific horticulture practices not only increases yield but also improves tuber quality in terms of size, shape, dry matter, starch content, and shelf-life, especially in varying climatic conditions.

a. Use of Good Quality Seed Tubers - Use of good quality, certified and disease free seed tubers is an important step in quality potato production. Health seed tubers having proper physiological age and proper size (30-50 g) make possible uniform germination, strong plant growth, and high marketable yield. Seed tubers can be treated with fungicides or biocontrol agents to prevent seed borne diseases and crop establishment. **b. Variety Selection-** Variety selection according to local agro-climatic conditions and market

Journal of Animal Environment, Vol. 18, No. 7s, Spring 2026 demand plays an important role in determining tuber quality. Varieties having high dry matter content, cooking quality, disease resistance and heat/drought tolerance are the best for obtaining quality tubers. **c. Proper Planting Techniques-**On time planting with proper spacing and depth results in proper growth of the plants and proper development of tubers. Proper spacing avoids competition for nutrition, water, and sunlight among the plants which helps in growing large and uniform size of tubers. Appropriate depth of planting is also important as it helps in preventing greening of the tubers. **d. Balanced Nutrient Management-** Balanced application of the nutrients is an essential component for growing quality tubers. Nitrogen favours vegetative growth but its excess application results in reduced dry matter production and delay in maturation process. Phosphorus favors root development and tuber formation whereas potassium increases the starch level, tuber hardness, skin quality, storage quality and resistance against physiological disorders. Organic manures and bio-fertilizers add to the soil fertility and increase nutrient use efficiency. **e. Irrigation Management-**Maintenance of optimum soil moisture is essential at the time of tuber initiation and tuber bulking stage for the production of high quality potatoes. Drought results in small, misshapen, and cracked tubers. On the other hand excess irrigation may result in waterlogging and diseases infection along with poor storage quality of the tubers. **f. Earthing Up and Mulching-**Earthing up avoids the exposure of the tubers to sunlight, prevents their greening and solanine production. Other benefits include better soil aeration, tuber expansion and weeding. Organic and plastic mulching conserves soil moisture, regulates soil temperature, controls weed development, and increases tuber cleanliness and quality. **g. Integrated Pest and Disease Management-**It is crucial to manage diseases such as late blight, early blight, bacterial wilt, and common scab, and also insects such as aphids and potato tuber moth in order to ensure the best quality of tubers. Integrated Pest and Disease Management (IPDM), through the use of disease-resistant varieties, crop rotation, biological control agents, field sanitation, and proper pesticide use, lowers crop damage and decreases pesticide residues. **h. Harvesting and Post-Harvest Management-**Harvesting of the tubers during physiological maturity results in maximized tuber size, dry matter and storability of the tubers. Post-harvest losses, nutrient levels, and shelf-life of potatoes can be lowered by careful harvesting that prevents mechanical damage. Curing, grading, packing and storing of the tubers under optimum temperature (4–10 °C according to the end purpose) and relative humidity (90–95%). **i. Climate-Smart Horticultural Management Techniques-**The use of climate-smart horticulture practices like conservation tillage, precision irrigation, nutrient management, crop rotation, and decision making based on weather patterns improves the quality of potatoes as per changes in the climate.

7. Year-wise Advances in Potato Breeding for Climate Resilience in India

S. No	Year/Period	Important Breeding Achievement	Climate Resilience Characteristic	Cultivar/Technological Development/variety	Impact
1	195	Commenc	Suitabilit	Kufri Jyoti,	Establish

	8 - 1970	ement of scientific breeding of potatoes by ICAR-CPRI	y to Indian Agro climatic conditions	Kufri Chandramukhi	ed a base for potato breeding in India
2	1971 - 1990	Disease resistant varieties	Late Blight resistance and suitability	Kufri Badshah, Kufri Sindhuri	Benefited from improved yield stability across different environments.
3	1991 - 2000	High yielders and processable varieties	Early maturity and processability	Kufri Pukhraj, Kufri Chipsona-1	Facilitated increased yield and suitability for processing industries.
4	2001 - 2010	Multi trait breeding	Disease resistance and increased dry matter content	Kufri Chipsona-3, Kufri Frysona	Improved industrial use and quality.
5	2011 - 2015	Introduction of climate-adaptation in breeding programs	Heat resistance and multiple stresses resistance	Kufri Surya, Kufri Khyati	Cultivation in warm potato growing areas
6	201	Increased	Heat	Kufri Mohan,	Increase

	6-2020	efforts on climate resilient breeding programs	resistance , drought adaptation, and late blight resistance	Kufri Lima	d resilience and processing qualities due to climatic changes
7	2021-2024	Development of future generation of climate smart cultivars	Heat resistance , stability of yield, insect resistance	Kufri Bhaskar	Increased stability of yield under heat stress and resistance to mites and hopper burn
8	2025	Greater breeding efforts and collaboration with other countries	Climate smart breeding and seed system development	ICAR-CPRI and International Potato Center (CIP) collaboration	Greater efforts in breeding and climatically resilient potato research in India.
9	2026	Announcement of new climate-smart potatoes	Heat tolerance, disease resistance , yield, and processing qualities	Kufri Tejas, Kufri Ratan, Kufri Chipbharat-1, Kufri Chipbharat-2, Kufri Abhedya, Kufri Ruby	Breeding achievements that aid in climate-smartness, potato processing

					g industrie s, and farmer incomes
--	--	--	--	--	---

8. Integrated Disease Management (IDM) in Potato

Integrated Disease Management (IDM) in potato refers to a comprehensive and sustainable system that involves cultural, genetic, biological, physical, and chemical measures to limit disease occurrence and reduce pesticide usage. Potatoes are vulnerable to several diseases caused by fungi, bacteria, viruses, nematodes, and phytoplasma, including late blight, early blight, black scurf, bacterial wilt, common scab, potato viruses (PVY, PVX, PLRV) and potato cyst nematodes. Proper IDM starts with the selection of healthy seed potatoes and disease-resistant/tolerant cultivars. Use of disease-free seed eliminates pathogens' entry to the field and provides good crop establishment. The application of cultural practices plays an important role in disease control. Crop rotation with non-host crops for 2–3 years assists in limiting soil-borne pathogens like *Rhizoctonia solani*, *Streptomyces scabies* and *Ralstonia solanacearum*. The practice of field sanitation and removal of diseased plant parts, volunteer potatoes and weeds is necessary for eliminating sources of inoculum. Well-drained soils, balanced fertilization, appropriate spacing and irrigation decrease humidity and development of late and early blight. The biological control measures are being considered in IDM of potatoes. The beneficial microorganisms include *Trichoderma harzianum*, *Trichoderma viride*, *Pseudomonas fluorescens*, and *Bacillus subtilis*, which help in combating soil-borne pathogens through competition, antibiosis, and induction of systemic resistance. Seed inoculation and application of these biocontrol agents increase plant vitality and decrease the risk of infection in a sustainable manner. Chemical control continues to play a vital role along with other methods of management. Potatoes can be treated with the right fungicides before planting to avoid seedborne and soil-borne infections. Prophylactic fungicide applications through spraying with contact fungicides including mancozeb or chlorothalonil, followed by systemics such as metalaxyl, cymoxanil, mandipropamid, or fluopicolide, help in preventing late blight under favorable weather conditions. These fungicides must be alternated depending on their various modes of action to overcome the possibility of developing resistance. Insecticides may also be needed to control the insect vectors, which spread viruses. Disease monitoring and prediction systems are great aids for quick interventions. Disease symptoms can be detected through field scouting while the disease prediction models based on weather forecast the onset of certain diseases like late blight. Technologies for precision farming, remote sensing, UAVs, and decision support systems add to the capabilities of disease monitoring and efficient application of pesticides. The successful IDM program for potatoes utilizes resistant cultivars, seed health, crop rotation, proper nutrition, biocontrol agents, timely fungicides, vector management, and regular disease monitoring. Apart from the reduction of crop loss and reduced costs of production, an IDM program results in the improvement of potato quality, reduction of the development of pathogen resistance, environmental safety, and sustainability of potatoes cultivation.

9. Role of Biological Control and Microbial Technologies in Potato Disease Control

The application of


Journal of Animal Environment, Vol. 18, No. 7s, Spring 2026

biological control methods and microbial technology has become an important aspect of sustainable potato farming as a means of ensuring an environmentally friendly alternative to indiscriminate use of chemicals. These techniques involve the use of beneficial microorganisms and their metabolites to control diseases, promote plant growth, and improve soil quality. The biological control agents (BCAs) function through such methods as competition for nutrients and space, antibiosis, mycoparasitism, secretion of hydrolytic enzymes, and induction of systemic resistance in plants. Their incorporation in potato farming reduces the prevalence of major diseases like late blight, early blight, black scurf, common scab, bacterial wilt, and soft rot, while limiting environmental pollution and emergence of resistance to pesticides. Some of the commonly used biological fungi in potato farming include *Trichoderma harzianum*, *Trichoderma viride*, and *Trichoderma asperellum*. These organisms help to control soil borne pathogens such as *Rhizoctonia solani*, *Fusarium* spp., and *Sclerotium rolfsii* by mycoparasitism, enzymatic action, and competition. Besides, they improve the growth of roots and facilitate nutrient absorption, producing healthy plants and increasing tuber yield. In addition to beneficial fungi, bacteria have also proven helpful in the control of potato diseases. The bacteria *Pseudomonas fluorescens*, *Bacillus subtilis*, *Bacillus amyloliquefaciens*, and *Bacillus velezensis* are capable of producing antimicrobial agents, siderophores, and lytic enzymes that suppress fungal and bacterial pathogens. Most of the aforementioned bacteria have plant growth-promoting rhizobacteria (PGPR) properties, increasing nutrients uptake, root growth, and environmental stress tolerance. Treatment of seeds with PGPR has been reported to increase plant vigor and suppress *Ralstonia solanacearum* and *Streptomyces scabies*. Microbial technologies have greatly contributed to biological control of diseases. Microbial consortiums, consisting of many beneficial microorganisms, are more efficient in disease suppression than single strains. New bioformulations have increased storage life of inoculum, such as liquid formulations, encapsulation of microbes, and polymer-coated inoculants. With the help of genomics, metagenomics, and microbiome engineering new insights into beneficial microbes of potato rhizosphere were achieved, resulting in improvement of the efficacy of bioinoculants. In addition, nanotechnology-based microbial delivery systems are being developed for protection of microbial viability and colonization in the field.

Journal of Animal Environment, Vol. 18, No. 7s, Spring 2026

chemical fertilizers, organic manures, crop residues, bio-fertilizers, and others is made together so that the soil is maintained fertile, nutrients are utilized efficiently, and potato crops give high yields. The potato plant is a nutrient-exhausting crop because it has shallow root system and grows rapidly. This crop is highly productive too. Thus, an INM approach is necessary in order to increase the yield, quality of tubers, and their pest, disease, and abiotic stress tolerance. Use of organic nutrients in the form of farmyard manure (FYM), compost, vermicompost, poultry manure, and green manures improves the soil properties, both physically and chemically. The application of organic materials increases the soil organic carbon content and makes the soil structure better by increasing its water holding capacity and stimulating beneficial microorganisms. It gradually releases nutrients in the soil throughout the crop growth period. Application of well-decomposed FYM is widely recommended in the rate of 20–25 t ha⁻¹ before planting. Chemical fertilizers supply the essential nutrients needed at the critical growth periods of potato plant. Nitrogen (N) favors vegetative growth and leaf canopy, phosphorus (P) stimulates the growth of roots and tuber initiation, while potassium (K) favors tuber bulking, starch formation, disease resistance, and tuber storage. Nutrient application based on soil analysis helps to avoid nutrient deficiencies and excessive use of fertilizers, hence reducing environmental pollution. Nitrogen split application together with phosphorus and potassium basally is an efficient way of utilizing the nutrients without causing their loss by leaching or volatilization. The use of biofertilizers in INM programs is increasing to enhance the availability of nutrients and minimize the use of chemical fertilizers. Nitrogen-fixing bacteria like *Azotobacter* and *Azospirillum* provide nitrogen fixed biologically, while phosphate-solubilizing bacteria and phosphate-solubilizing fungi increase phosphorus availability from insoluble forms in the soil. Arbuscular mycorrhizal fungi enhance phosphorus uptake, water uptake ability, and drought and disease tolerance in plants. Combining the use of biofertilizers with the other sources of nutrients increases nutrient use efficiency. Equally important is the role of micronutrients in potato growth and tuber quality. Zinc, boron, magnesium, sulfur, iron, and manganese are involved in enzyme activation, chlorophyll formation, carbohydrate utilization, and tuber formation. Their deficiency can cause reduced yield, poor tuber quality, and increased vulnerability to diseases. Soil and plant analysis-assisted site-specific application of micronutrients guarantees adequate nutrition and top crop performance. The latest INM developments are precision nutrient management, sensor-assisted fertilizer recommendation, fertigation using drip irrigation, controlled-release fertilizers, nanofertilizers, and digital decision support systems. These innovations optimize fertilizer use efficiency, decrease losses of nutrients and environmental impact, and reduce costs of fertilizer use. The tools of precision agriculture such as remote sensing and GIS soil fertility mapping allow farmers to use nutrients according to needs of the crop and the field. An effective INM strategy should include soil testing, proper fertilization, organic manure application, biofertilizer use, micronutrient application, and proper nutrient management. Such an integrated approach helps increase potato yields, size and quality of the potatoes, their dry matter content and processing quality, increases resistance to diseases and adverse environment, as well as soil health maintenance.


11. Integrated Pest Management (IPM) in Potato
Integrated Pest Management (IPM) is a sustainable and eco-friendly practice that utilizes a combination of cultural, biological, mechanical, physical, genetic, and chemical measures for controlling insect pests in potatoes while reducing the use of



(a)

(b)

(c)



Biological Control and Microbial

Technologies

10. INM for Potato Crop(Integrated Nutrient Management)
INM technique is the sustainable system wherein the use of

pesticides and their adverse effects on the environment. There are a number of economically important insect pests that attack potatoes, which include aphids (*Myzus persicae* and *Aphis gossypii*), potato tuber moth (*Phthorimaea operculella*), whiteflies (*Bemisia tabaci*), cutworms (*Agrotis ipsilon*), leafhoppers, thrips, white grubs, and termites. In addition to causing damage by feeding on the crop, some of these insect pests act as vectors of viral diseases like Potato virus Y (PVY) and Potato leafroll virus (PLRV), thus causing losses in potato production. Prevention using cultural practices forms the cornerstone of IPM. Farmers should use certified and pest-free seed potatoes, practice crop rotation with non-host crops, field sanitation involving removal of volunteer potatoes and weed hosts, and timely planting of potato crops to avoid peak levels of insect pests. Earthing up helps prevent potato tuber moth attacks, while proper fertilization and irrigation make the potato crop healthy and less prone to insect infestations. Monitoring and surveillance of pest population on regular basis are necessary for implementing IPM strategies. Pest population must be detected on fields in early stages through frequent scouting. Yellow sticky traps are helpful in detecting aphids and whitefly populations while pheromone traps help in detecting and trapping potato tuber moth population. Control methods should only be implemented when pest population exceeds the Economic Threshold Level (ETL). Biological control is a very important component of sustainable potato pest control. Natural predators like *Coccinella septempunctata* (ladybird beetle), *Chrysoperla carnea* (green lacewing), and hoverflies control aphids naturally. *Trichogramma* wasps are natural parasitoids of eggs of potato tuber moths and many other lepidopteran pests. Entomopathogenic fungi like *Beauveria bassiana* and *Metarhizium anisopliae*, as well as bacteria like *Bt* (*Bacillus thuringiensis*) can successfully control different insect pests without causing harm to beneficials. Mechanical and physical means additionally decrease the numbers of the pests. Infected parts of the plants, deep summer ploughing to reveal the larvae, proper storage sanitation, building of an appropriate storage facility, usage of light and sticky traps are among the factors that help to efficiently manage pests. While storing potatoes, maintenance of recommended temperature and humidity levels prevents the potato tuber moth and storage pests' infestations. The application of insecticides must be reasonable and only if the other means do not work. Selective insecticides must be used according to the pest incidence and economic threshold level while changing the insecticides with various mechanisms of action in order to prevent development of insecticide resistance. Insecticide treatment of the seeds might help to protect the plants from pests during the growing season. Strict observance of recommended doses, intervals of spraying and preharvest intervals helps to minimize the presence of pesticides and to ensure safety for humans and environment. Recent developments in integrated potato pest management include precise pest monitoring, remote sensing, pest monitoring via drones, decision support systems, pest forecasting using weather, RNA interference technology, semiochemicals pest management and host plant resistance developed with traditional breeding and biotechnology methods. IPM system incorporates the use of pest-resistant varieties, disease-free planting materials, cultural management, biological control agents, monitoring, mechanical control, and chemicals on as-needed basis. Such an approach will help keep pest numbers to levels that do not cause any economical damage, reduce production costs, delay the development of insecticide resistance, conserve the environment, enhance tuber yield, and make the production profitable.

12. Climate-Smart Technologies for Potato Farming Sustainability

Climate-smart technologies refer to innovative techniques that aim at enhancing potato productivity, improving resistance to climate change and minimizing the environmental impact of potato cultivation. Potato farming is vulnerable to variations in temperatures, rainfall, droughts, flooding, and other weather-related problems due to its sensitivity. This makes potato plants suffer from reduced productivity and yield. The combination of advanced technologies, sustainable farming practices, and efficient management of natural resources plays an integral part in the implementation of Climate-Smart Agriculture (CSA). One of the most powerful climate-smart strategies includes the breeding and cultivation of climate-resistant potato varieties. Modern breeding techniques, such as conventional breeding, marker-assisted selection, genomic selection, and gene-editing technologies, enable to cultivate potato varieties tolerant to heat, drought, salinity, frost and resistant to late blight and viruses. Precision agriculture plays a very crucial role in climate smart potato production. Technologies such as GPS, GIS, remote sensing, UAVs, satellite images, soil sensors, and IoT based monitoring systems can be used to monitor crop condition, soil moisture content, nutrients in soil and any outbreak of diseases. Variable rate application of fertilizers, water and pesticides helps in improving the use of inputs, minimizing production cost and decreasing environmental pollution. Efficient water management technologies are important as potatoes are very susceptible to both drought and excess water in the field. Drip irrigation, sprinkler irrigation, fertigation, automatic irrigation schedule, rainwater harvesting and soil moisture sensors will help in making efficient water use and having appropriate soil moisture content for proper development of tubers. Mulching of the crop using organic residues or biodegradable plastic films. Climate-smart nutrient management focuses on balancing the fertilization using Integrated Nutrient Management (INM) involving the use of organic manures, compost, bio-fertilizers, crop residues, and chemical fertilizers after doing soil tests. Controlled release of fertilizers, nano-fertilizers, precision fertigation, and decision support systems contribute to increased nutrient-use efficiency, reduction of greenhouse gas emission, and soil quality improvement. Conservation agriculture measures such as minimum tillage, crop residues, and crop rotation are more beneficial to the soils in terms of organic carbon content, moisture conservation, and soil fertility. Another major climate-smart technology involves Integrated Pest and Disease Management (IPM and IDM) with the help of weather forecasting systems. Early warning systems, disease prediction models, mobile technologies, and advisory systems assist the farmer in making timely decisions concerning diseases like late blight and insect pests, leading to reduced application of pesticides. Bio agents and disease-resistant varieties add on to sustainable pest and disease management. Other modern technologies that can help potatoes to become more resilient to climate change include cold storages equipped with automated systems of temperature and humidity regulation, controlled atmosphere (CA) storage, renewable energy-driven storage, and enhanced ventilation. Irrigation systems powered by solar energy, renewable energy technologies, and energy-efficient farm machinery help to minimize the use of fossil fuels and limit greenhouse gas emissions. Advances in AI, machine learning, big data analytics, digital platforms for agriculture, climate models, and smart mobile applications give farmers detailed recommendations concerning irrigation, fertilization, pest and disease management, and harvesting. These technologies assist in improving decision-making processes, optimizing the use of resources, and enhancing climate change resilience. To sum up, climate-smart technologies offer a comprehensive approach to

sustainable potato production by including improved varieties, precision agriculture, advanced water and nutrients management, IPDM, conservation agriculture, renewable energy, and digital technologies. The use of such technologies increases potato yield, improves tubers quality, saves natural resources, minimizes greenhouse gas emissions, and increases climate change resilience of potato farming systems.

13. Post-Harvest Management for the Maintenance of Potato Quality

Post-harvest management is an essential aspect of potato production that seeks to maintain tuber quality, prevent quantitative and qualitative losses, and prolong the storage life from harvest up to consumption or processing. Poor post-harvest management causes damage to the potato due to bruising, loss of weight, sprouting, occurrence of diseases, and poor processing quality. Proper post-harvest management includes proper harvesting, curing, grading, storage, transportation, and packing of potatoes to maintain their physical, physiological, and nutritional quality. Tuber harvesting should be done when tubers attain physiological maturity and the haulms have naturally dried or haulms have been destroyed. Tubers should be harvested from dry soil to avoid mechanical damage and soil contamination. Mechanically damaged tubers suffer from cuts, bruises, and skinning and are therefore vulnerable to diseases and rots during storage. Tubers should be harvested in cool weather and protection from direct sunlight is important for tuber quality. Immediately after harvest, the tubers need to be cured for 10-14 days at 15-20 °C with relative humidity of 85-95 %. The process of curing aids the wound healing by suberization and formation of protective cork covering which prevents water loss and infection by fungi like *Fusarium* spp., *Pectobacterium* spp. and *Rhizoctonia solani*. Diseased, damaged, green and deformed tubers have to be discarded during grading and sorting. Proper storage is necessary for the maintenance of quality of potatoes. Fresh table potatoes are stored at 4-7°C whereas the seed potatoes are kept at 2-4°C for the preservation of physiological quality. The potatoes meant for processing into chips and French fries are stored at 8-12°C to prevent cold induced sweetening which results in high sugar content causing black coloring during frying. Relative humidity should be 90-95 % in order to prevent water loss and shrinkage along with good ventilation. Another vital component of post-harvest management involves sprout control. Sprouts accelerate respiration and moisture loss as well as reduce nutrients, and thus they make potatoes less marketable and decrease their quality during the process of processing. Currently, modern sprout inhibitors, low temperatures and improved ventilation help to delay sprouting while retaining the quality of tubers. Regular observation of the storage environment is required to avoid physiological disorders like blackheart, pressure bruising, chilling injury and excessive weight loss. During packaging and transportation, the main objective is to prevent mechanical injury and exposure of potatoes to unfavorable environment. Compared to sacks, ventilated plastic containers, mesh bags and corrugated fiberboard cartons provide better protection for tubers. Gently moving potatoes when loading, unloading and transporting helps to prevent bruises and minimizes the entry of storage pathogens. Diseases control in post-harvest includes sanitization of storage facilities, removal of affected tubers, and maintenance of proper temperature and humidity. Hygiene, disinfection of storage facilities, and inspection can prevent diseases like dry rot (*Fusarium* spp.), soft rot (*Pectobacterium* spp.), silver scurf (*Helminthosporium solani*), and late blight tuber rot. Emerging technologies that have transformed the field of post-harvest management include automated grading machines, optical sorting, non-destructive

Journal of Animal Environment, Vol. 18, No. 7s, Spring 2026
quality assessment with hyperspectral imaging, IoT-based storage management, smart sensing technologies, traceability through RFID, AI-based quality prediction models, and cold storage powered by renewable energy sources. These innovations help in improving efficiency of storage, reducing losses, maintaining quality, and increasing the value of potatoes. To sum up, post-harvest management involves efficient harvesting, curing, grading, scientific storage, sprout control, disease prevention, packaging, and use of modern storage monitoring technologies. All these measures greatly reduce post-harvest losses, improve quality, increase shelf-life, enhance processing qualities, and increase profitability and sustainability of potato farming.

Conclusion

It is very critical to consider integrated strategies that will ensure sustainable production of potatoes amidst all the changing climatic conditions. Combining improved agronomy, climate-smart varieties, INM, IPM, IDM, biotechnology, precision farming, water management, and advanced post-harvest handling techniques will be a complete package to improve potato production, tuber quality, and its resistance to diseases. Instead of applying just one management technique, an integration of various management technologies will enhance resource use efficiency, increase crop resilience, and stabilize yields despite various climatic conditions. The changes brought about by climate change have increased heat stress, drought, erratic rainfall patterns, and emergence of new pests and diseases that have made traditional production practices unreliable. Thus, climate-smart technologies like precision irrigation, sensor-based nutrient management, disease prediction based on weather conditions, remote sensing, AI, IoT, drones, and decision support systems are becoming increasingly vital. Certified seed tubers free from diseases, resistance varieties, crop rotation, fertilizer balance, conservation agriculture, beneficial microorganisms, as well as pesticide and fertilizer use based on need can greatly help in reduction of pest and disease burden and increase soil health and biodiversity. In addition to that, scientific advances in molecular breeding, genomics, gene editing, microbiome engineering, nanotechnology-based agriculture, and precision farming are offering new possibilities in designing high-yielding and climate-resilient potato production systems. Post-harvest management is crucial in preservation of tuber quality, loss reduction, long-term storage, and increasing market value. Harvesting techniques, curing, grading, improvement of storage facilities, cold chain and monitoring systems contribute to maintenance of processable quality and reduction of food waste; therefore, contributing to increase of farmers' income and availability of quality potatoes throughout the year. Future research should concentrate on integration of modern breeding technologies with digital agriculture, biological advancements and sustainable resource management as a solution to problems caused by climate change. Special attention should also be paid to location-specific management practices, farmers' education, policies and technologies transfer. Enhancement of cooperation between researchers, extension service, policymakers, and farmers will contribute to faster development and distribution of innovative technologies. In summary, integrated techniques are the best option when it comes to increasing potato yield, improving quality, disease resistance, and adaptation to climate variability. These techniques can help achieve sustainable agriculture, better livelihoods of farmers, food and nutrition security, natural resource conservation, and overall environmental sustainability. With climate change influencing agricultural practices, integrated and climate smart potato production will be essential in ensuring profitable production of potatoes.

References

REFERENCES

1. Adekanmbi T., Wang X., Basheer S., Nawaz R.A., Pang T., Hu Y., Liu S. Assessing Future Climate Change Impacts on Potato Yields - A Case Study for Prince Edward Island, Canada. *Foods*. 2023, 12(6), 1176. doi:10.3390/foods12061176.
2. Alyokhin A. - Colorado potato beetle management on potatoes: Current challenges and future prospects. *Fruit, Vegetable, and Cereal Science and Biotechnology*, 2007, 3 (special issue 1), 10-19.
3. Anonymous. - The measurement of potato blight. *Trans. Brit. Mycological Society* 31, U.K. 1947, pp. 140-141
4. Bucurean E., Bunta G. - Control of wireworms (*Agriotes* spp.) by chemical seed treatment. *Analele Universității din Oradea. Fascicula Protecția Mediului*, 2011, Vol. XVII, 23-26.
5. Čepl J., Kasal P., Svobodova A. - Potato weed management in condition of technology of destoned ridges. 48th Symposium "Actual Tasks on Agricultural Engineering", Zagreb, Croatia, 2021, 361-369 Downloaded from <https://cabidigitallibrary.org> by 82.208.136.135, on 10/28/24.
6. Cingel A., Savić J., Lazarević J., Ćosić T., Raspor M., Smigocki A., Ninković S. - Extraordinary Adaptive Plasticity of Colorado Potato Beetle: "Ten-Striped Spearman" in the Era of Biotechnological Warfare. *International Journal of Molecular Sciences*, 2016, 17(9), 1538. <https://doi.org/10.3390/ijms17091538>.
7. Cohen Y., Cisi, U., Mosinger E. - Systemic resistance of potato plants against *Phytophthora* infestans induced by unsaturated fatty acids. *Physiol., Molecular, Plant Pathol.*, 1991, 38, 255 – 263.
8. Ferro D.N. - Biological control of the Colorado potato beetle. In: Zehnder GW, Jansson RK, Powelson ML, Raman KV (Eds.) *Advances in Potato Pest Biology and Management*, 1994, APS Press, St. Paul, 357-37.
9. Haverkort A.J., Boonekamp P.M., Rutten R., Jacobsen E., Lotz L.A.P., Kessel G.J.T., Visser R.G.F., van der Vossen E.A.G. - Societal costs of late blight in potato and prospects of durable resistance through cisgenic modification. *Potato Res.*, 2008, 51, 47–57. doi:10.1007/s11540-008-9089-y.
10. Jennings S.A., Koehler A.K., Nicklin K.J., Deva C., Sait S.M., Challinor A.J. Global Potato Yields Increase Under Climate Change with Adaptation and CO2 Fertilisation. *Front. Sustain. Food Syst.* 2020;4:519324. doi: 10.3389/fsufs.2020.519324.
11. Menari M., Dadashi M.R., Sheikh F., Ajamnoruzi H. - Evaluation of drought stress tolerance in some faba bean genotypes using drought tolerance indices. *Romanian Agricultural Research*. 2022. no. 39, 35-43.
12. Meno L., Escuredo O., Rodríguez-Flores M.S., Seijo M.C. - Modification of the TOMCAST model with aerobiological data for management of potato Early blight. *Agronomy*, 2020, 10 (12), 1872. doi:10.3390/agronomy10121872.
13. Meno L., Escuredo O., Rodríguez-Flores M.S., Seijo M.C. - Looking for a sustainable potato crop. *Field Journal of Animal Environment*, Vol. 18, No. 7s, Spring 2026 assessment of early blight management. *Agricultural and Forest Meteorology*, 2021, Volumes 308–309, <https://doi.org/10.1016/j.agrformet.2021.108617>.
14. Mohamed S.P.- Integrated Pest Management Strategies for Sustainable Agriculture: A Review of Current Practices and Future Directions. *Asian Journal of Advances in Research*, 2023, Volume 6, Issue 1, 475-483.
15. Nikoukar A., Rashed A. - Integrated Pest Management of Wireworms (Coleoptera: Elateridae) and the Rhizosphere in Agroecosystems. *Insects*. 2022, 13(9),769. <https://doi.org/10.3390/insects13090769>.
16. Poggi S., Le Cointe R., Lehmhus J., Plantegenest M., Furlan L. Alternative strategies for controlling wireworms in field crops: A review. *Agriculture*. 2021, 11(5), 436 <https://doi.org/10.3390/agriculture11050436>.
17. Runno-Paurson E., Loit K., Hansen M., Tein B., Williams I. H., Mänd M. - Early blight destroys potato foliage in the northern Baltic region. *Acta Agriculturae Scandinavica, Section B — Soil & Plant Science*, 2015, 65(5), 422–432. <https://doi.org/10.1080/09064710.2015.1017003>.
18. Skendžić S. Zovko M., Živković I.P., Lešić V., Lemić D. The Impact of Climate Change on Agricultural Insect Pests. *Insects*. 2021, 12(5), 440. <https://doi.org/10.3390/insects12050440>.
19. Skiba D., Sawicka B., Pszczółkowski P., Barbaś P., Krochmal-Marczak B. The impact of cultivation management and weed control systems of very early potato on weed infestation, biodiversity, and health safety of tubers. *Life (Basel)*. 2021, 11(8), 826. doi: 10.3390/life11080826.
20. Struik P.C., Wiersema S.G. - Seed potato technology. Wageningen Pers., Wageningen, 1990, 51.
21. Tiwari I., Shah K.K., Tripathi S., Modi B., Subedi S., Jiban Shrestha J. - Late blight of potato and its management through the application of different fungicides and organic amendments: a review. *Journal of Agriculture and Natural Resources*, 2021, 4(1), 301-320 doi: <https://doi.org/10.3126/janr.v4i1.33374>.
22. van Herk W.G., Vernon R.S., Goudis L., Mitchell T. - Protection of potatoes and mortality of wireworms (*Agriotes obscurus*) with various application methods of broflanilide, a novel meta-diamide insecticide. *Journal of Economic Entomology*, 2022, Volume 115, Issue 6, 1930–1946, <https://doi.org/10.1093/jee/toac152>.
23. Vernon R.S., van Herk W.G. - Wireworms as pests of potato. In: Giordanengo P., Vincent C., and Alyokhin A. (ed.), *Insect pests of potato: global perspectives on biology and management*, 2nd ed. Amsterdam: Academic, 2022, 103–148.
24. Weng L., Tang Z., Sardar M.F., Yu Y., Ai K., Liang S., Alkahtani J., Lyv D. - Unveiling the frontiers of potato disease research through bibliometric analysis. *Front. Microbiol.*, 2024, 15:1430066. doi: 10.3389/fmicb.2024.1430066.
25. Yellareddy S.K.R., Taylor R.J., Julie S. Pasche, Gudmestad N.C. - Quantifying control efficacy of fungicides commonly applied for potato early blight management. *Plant Disease*, 2019, vol 103, no. 11, 2821-2824 <https://doi.org/10.1094/PDIS-03-19-0670-RE>.

26. Zarzecka K., Gugala M., Sikorska A., Grzywacz K., Niewęglowski M. - Marketable yield of potato, and its quantitative parameters after application of herbicides and biostimulants. Agriculture. 2020, 10, 49. doi: 10.3390/agriculture10020049.